# 3D visualization of intraoperative stimulation test results in deep brain stimulation.

A. Shah<sup>1</sup>, F. Alonso<sup>2</sup>, J.J. Lemaire<sup>3</sup>, D. Pison<sup>1</sup>, J. Coste<sup>3</sup>, K. Wårdell<sup>2</sup>, E. Schkommodau and S. Hemm<sup>1,2,\*</sup>

<sup>1</sup>Institute for Medical and Analytical Technologies, School of Life Sciences, University of Applied Sciences and Arts Northwestern Switzerland, Muttenz, Switzerland

<sup>2</sup>Department of Biomedical Engineering, Linköping University, Linköping, Sweden

<sup>3</sup>Centre Hospitalier Universitaire de Clermont-Ferrand, Image-Guided Clinical Neurosciences and Connectomics (EA 7282, IGCNC),

Université d'Auvergne, Clermont-Ferrand, France

### I. Introduction

Deep brain stimulation (DBS) is a treatment for movement disorders such as Essential Tremor (ET). The implant position of the electrode is crucial. After preoperative surgical planning and exhaustive intraoperative tests, the collected information is only "mentally" visualized and analyzed to decide on the optimal implant position of the DBS lead. We propose a method to present this multitude of information for surgical decision making using patient-specific simulations of electric field (EF) distribution in combination with intraoperative accelerometry based tremor evaluation and direct-targeting technique of DBS.

#### п. Метнор

Five ET patients participating in a clinical study (written informed consent) where included in the presented protocol. The procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2000 and 2008. After preoperative selection of the anatomical target position, two microelectrodes were inserted per hemisphere on parallel trajectories. Stimulation tests were performed on 5 to 10 positions per trajectory and several stimulation amplitudes per position to evaluate side effects and changes in tremor using an accelerometer fixed to the patient's wrist. To estimate the extension of the stimulation, electric field simulations were performed for 2 to 3 amplitudes per position resulting in different tremor improvements. To identify the optimal implant position for chronic stimulation, we summarized all data in "Improvement Maps" by assigning each voxel in the stimulation test region the highest improvement in tremor and by visualizing them in relation to patient-specific anatomy. Postoperatively the optimal implant positions for the 5 patients (10 hemispheres) were identified.

## III. RESULTS

The clinical teams were able to identify the optimal implant position for the 5 patients with more ease and in less time compared to the routine discussion using pen and paper. Additionally, for 7 of the 9 improvement maps, the highest improvement region was found to be in the posterior subthalamic area inferior and posterior to the ventro-intermediate (VIM) nucleus of the thalamus (routine clinical target).

# IV. CONCLUSION

Visual analysis of the results of intraoperative stimulation tests in form of improvement maps assists the clinicians in determining the optimal implant location of the chronic DBS lead and in comparing results between patients.

## V. CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

#### VI. REFERENCES

1. Zerroug A, Gabrillargues J, Coll G et al., (2016).

Personalized mapping of the deep brain with a white matter attenuated inversion recovery (WAIR) sequence at 1.5-tesla: Experience based on a series of 156 patients. Neurochirugie; 5014:183–231. DOI 10.1016/j.neuchi.2016.01.009.

- 2. Shah A, Coste J, Lemaire J-J et al., (2016). Intraoperative acceleration measurements to quantify improvement in tremor during deep brain stimulation surgery. Med Biol Eng Comput. DOI 10.1007/s11517-016-1559-9.
- 3. Hemm S, Pison D, Alonso F et al., (2016). Patient-Specific Electric Field Simulations and Acceleration Measurements for Objective Analysis of Intraoperative Stimulation Tests in the Thalamus. Front. Hum. Neurosci.; 10:577(1-14). DOI 10.3389/fnhum.2016.00577

Author: Prof. Dr. Simone Hemm-Ode

Institute: Institute of Medical and Analytical Technologies, University of Applied Sciences and Arts Northwestern Switzerland.

Street: Gruendestrasse 40 City: 4132 Muttenz Country: Switzerland

Email: simone.hemm@fhnw.ch