



**HAL**  
open science

## Zeroes of the derivative of a p-adic meromorphic function and applications

Kamal Boussaf, Jacqueline Ojeda, Alain Escassut

► **To cite this version:**

Kamal Boussaf, Jacqueline Ojeda, Alain Escassut. Zeroes of the derivative of a p-adic meromorphic function and applications. Bulletin of the Belgian Mathematical Society - Simon Stevin, 2013. hal-01907041

**HAL Id: hal-01907041**

**<https://uca.hal.science/hal-01907041v1>**

Submitted on 28 Oct 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Zeroes of the derivative of a p-adic meromorphic function and applications.

by Kamal Boussaf, Jacqueline Ojeda and Alain Escassut

**Abstract** Let  $K$  be an algebraically closed field of characteristic 0, complete with respect to an ultrametric absolute value. We show that if the Wronskian of two entire functions in  $K$  is a polynomial, then both functions are polynomials. As a consequence, if a meromorphic function  $f$  on all  $K$  is transcendental and has finitely many multiple poles, then  $f'$  takes all values in  $K$  infinitely many times. We then study applications to a meromorphic function  $f$  such that  $f' + bf^2$  has finitely many zeroes, a problem linked to the Hayman conjecture on a p-adic field.

2000 *Mathematics subject classification*: Primary 12J25 Secondary 46S10

*Keywords*: zeroes of p-adic meromorphic functions, derivative

Partially funded by the research project CONICYT "Inserción de nuevos investigadores en la academia" N°79090014 from the Chilean Government.

## Notation and Definitions.

Let  $K$  be an algebraically closed field of characteristic 0, complete with respect to an ultrametric absolute value  $|\cdot|$ . Given  $\alpha \in K$  and  $R \in \mathbb{R}_+^*$ , we denote by  $d(\alpha, R)$  the disk  $\{x \in K \mid |x - \alpha| \leq R\}$  and by  $d(\alpha, R^-)$  the disk  $\{x \in K \mid |x - \alpha| < R\}$ , by  $\mathcal{A}(K)$  the  $K$ -algebra of analytic functions in  $K$  (i.e. the set of power series with an infinite radius of convergence), by  $\mathcal{M}(K)$  the field of meromorphic functions in  $K$  and by  $K(x)$  the field of rational functions. Given  $f, g \in \mathcal{A}(K)$ , we denote by  $W(f, g)$  the Wronskian  $f'g - fg'$ .

We know that any non-constant entire function  $f \in \mathcal{A}(K)$  takes all values in  $K$ . More precisely, a function  $f \in \mathcal{A}(K)$  other than a polynomial takes all values in  $K$  infinitely many times (see [5], [8], [9]). Next, a non-constant meromorphic function  $f \in \mathcal{M}(K)$  takes every value in  $K$ , except at most one value. And more precisely, a meromorphic function  $f \in \mathcal{M}(K) \setminus K(x)$  takes every value in  $K$  infinitely many times except at most one value (see [5], [9]).

Many previous studies were made on Picard's exceptional values for complex and p-adic functions and their derivatives (see [1], [3], [6], [7], [8]). Here we mean to examine precisely whether the derivative of a transcendental meromorphic function in  $K$  having finitely many multiple poles, may admit a value that is taken finitely many times and then we will look for applications to Hayman's problem when  $m = 2$ .

From Theorem 4 [6], we can state the following Theorem A:

**Theorem A:** *Let  $h, l \in \mathcal{A}(K)$  satisfy  $W(h, l) = c \in K$ , with  $h$  non-affine. Then  $c = 0$  and  $\frac{h}{l}$  is a constant.*

Now we can improve Theorem A:

**Theorem 1:** *Let  $f, g \in \mathcal{A}(K)$  be such that  $W(f, g)$  is a non-identically zero polynomial. Then both  $f, g$  are polynomials.*

**Remark:** Theorem 1 does not hold in characteristic  $p \neq 0$ . Indeed, suppose the characteristic of  $K$  is  $p \neq 0$ . Let  $\psi \in \mathcal{A}(K)$ . Let  $f = x(\psi)^p$  and let  $g = (x + 1)\psi^{-p}$ . Since  $p \neq 0$ , we have  $f' = (\psi)^p$ ,  $g' = \psi^{-p}$  hence  $W(f, g) = 1$  and this is true for any function  $\psi \in \mathcal{A}(K)$ .

**Theorem 2:** *Let  $f \in \mathcal{M}(K) \setminus K(x)$  have finitely many multiple poles. Then  $f'$  takes every value  $b \in K$  infinitely many times.*

We can easily show Corollary 2.1 from Theorem 2, though it is possible to get it through an expansion in simple elements.

**Corollary 2.1:** *Let  $f \in \mathcal{M}(K) \setminus K(x)$ . Then  $f'$  belongs to  $\mathcal{M}(K) \setminus K(x)$ .*

**Open question:** Do exist transcendental meromorphic functions  $f$  such that  $f'$  has finitely many zeroes? By Theorem 2, such functions should have infinitely many multiple poles.

Now, we can look for some applications to Hayman's problem in a p-adic field. Let  $f \in \mathcal{M}(K)$ . Recall that in [9], [10] it was shown that if  $m$  is an integer  $\geq 5$  or  $m = 1$ , then  $f' + f^m$  has infinitely many zeroes that are not zeroes of  $f$ . If  $m = 3$  or  $m = 4$ , for many functions  $f \in \mathcal{M}(K)$ ,  $f' + f^m$  has infinitely many zeroes that are not zeroes of  $f$  (see [2], [10]) but there remain some cases where it is impossible to conclude, except when the field has residue characteristic equal to zero (see [10]). When  $m = 2$ , few results are known. Recall also that as far as complex meromorphic functions  $f$  are concerned,  $f' + f^m$  has infinitely many zeroes that are not zeroes of  $f$  for every  $m \geq 3$ , but obvious counter-examples show this is wrong for  $m = 1$  (with  $f(x) = e^x$ ) and for  $m = 2$  (with  $f(x) = \tan(-x)$ ). Here we will particularly examine functions  $f' + bf^2$ , with  $b \in K^*$ .

**Theorem 3:** *Let  $b \in K^*$  and let  $f \in \mathcal{M}(K)$  have finitely many zeroes and finitely many residues at its simple poles equal to  $\frac{1}{b}$  and be such that  $f' + bf^2$  has finitely many zeroes. Then  $f$  belongs to  $K(x)$ .*

**Remark:** If  $f(x) = \frac{1}{x}$ , the function  $f' + bf^2$  has no zero whenever  $b \neq 1$ .

**Theorem 4:** *Let  $f \in \mathcal{M}(K) \setminus K(x)$  have finitely many multiple zeroes and let  $b \in K$ . Then  $\frac{f'}{f^2} + b$  has infinitely many zeroes. Moreover, if  $b \neq 0$ , every zero  $\alpha$  of  $\frac{f'}{f^2} + b$  that is not a zero of  $f' + bf^2$  is a simple pole of  $f$  such that the residue of  $f$  at  $\alpha$  is equal to  $\frac{1}{b}$ .*

**Corollary 4.1 :** *Let  $b \in K^*$  and let  $f \in \mathcal{M}(K) \setminus K(x)$  have finitely many multiple zeroes and finitely many simple poles. Then  $f' + bf^2$  has infinitely many zeroes that are not zeroes of  $f$ .*

**Remark:** In Archimedean analysis, the typical example of a meromorphic function  $f$  such that  $f' + f^2$  has no zero is  $\tan(-x)$  and its residue is 1 at each pole of  $f$ . Here we find the same implication but we can't find an example satisfying such properties.

**Acknowledgement:** The authors are very grateful to Jean-Paul Bezin for an important improvement concerning Theorem 1.

## The Proofs.

**Notation:** Given  $f \in \mathcal{A}(K)$  and  $r > 0$ , we denote by  $|f|(r)$  the norm of uniform convergence on the disk  $d(0, r)$ . This norm is known to be multiplicative (see [4], [5]).

Lemma 1 is well known (see Theorem 13.5 [4]) :

**Lemma 1:** *Let  $f \in \mathcal{M}(K)$ . Then  $|f^{(k-1)}|(r) \leq \frac{|f|(r)}{r^{k-1}} \forall r > 0, \forall k \in \mathbb{N}^*$ .*

**Proof of Theorem 1:** First, by Theorem A, we check that the claim is satisfied when  $W(f, g)$  is a polynomial of degree 0. Now, suppose the claim holds when  $W(f, g)$  is a polynomial of certain degree  $d$ . We will show it for  $d + 1$ . Let  $f, g \in \mathcal{A}(K)$  be such that  $W(f, g)$  is a non-identically zero polynomial  $P$  of degree  $d + 1$ .

By hypothesis, we have  $f'g - fg' = P$ , hence  $f''g - fg'' = P'$ . We can extract  $g'$  and get  $g' = \frac{f'g - P}{f}$ . Now, consider the function  $Q = f''g' - f'g''$  and replace  $g'$  by what we just found: we can get  $Q = f'(\frac{f''g - fg''}{f}) - \frac{Pf''}{f}$ .

Now, we can replace  $f''g - fg''$  by  $P'$  and obtain  $Q = \frac{f'P' - Pf''}{f}$ . Thus, in that expression of  $Q$ , we can write  $|Q|(R) \leq \frac{|f|(R)|P|(R)}{R^2|f|(R)}$ , hence  $|Q|(R) \leq \frac{|P|(R)}{R^2} \forall R > 0$ . But by definition,  $Q$  belongs to  $\mathcal{A}(K)$  and further,  $\deg(Q) \leq \deg(P) - 2$ . Consequently,  $Q$  is a polynomial of degree at most  $d - 2$ .

Now, suppose  $Q$  is not identically zero. Since  $Q = W(f', g')$  and since  $\deg(Q) < d$ , by induction  $f'$  and  $g'$  are polynomials and so are  $f$  and  $g$ . Finally, suppose  $Q = 0$ . Then  $P'f' - Pf'' = 0$  and therefore  $f'$  and  $P$  are two solutions of the differential equation of order 1 for meromorphic functions in  $K : (\mathcal{E}) y' = \psi y$  with  $\psi = \frac{P'}{P}$ , whereas  $y$  belongs to  $\mathcal{A}(K)$ . The space of solutions of  $(\mathcal{E})$  is known to be of dimension 0 or 1 (see for instance Lemma 60.1 in [4]). Consequently, there exists  $\lambda \in K$  such that  $f' = \lambda P$ , hence  $f$  is a polynomial. The same holds for  $g$ .

**Proof of Theorem 2:** Suppose  $f'$  has finitely many zeroes. By classical results (see [4], [5]) we can write  $f$  in the form  $\frac{h}{l}$  with  $h, l \in \mathcal{A}(K)$ , having no common zero. Consequently, each zero of  $W(h, l)$  is a zero of  $f'$  except if it is a multiple zero of  $l$ . But since  $l$  only has finitely many multiple zeroes, it appears that  $W(h, l)$  has finitely many zeroes and therefore is a polynomial. Consequently, by Theorem 1, both  $h$  and  $l$  are polynomials, a contradiction because  $f$  does not belong to  $K(x)$ .

Now, consider  $f' - b$  with  $b \in K$ . It is the derivative of  $f - bx$  whose poles are exactly those of  $f$ , taking multiplicity into account. Consequently,  $f' - b$  also has infinitely many zeroes.

**Notation:** Given  $f \in \mathcal{M}(K)$ , we will denote by  $\text{res}_a(f)$  the residue of  $f$  at  $a$ .

**Lemma 2:** Let  $f = \frac{h}{l} \in \mathcal{M}(K)$  with  $h, l \in \mathcal{A}(K)$  having no common zero, let  $b \in K^*$  and let  $a \in K$  be a zero of  $h'l - hl' + bh^2$  that is not a zero of  $f' + bf^2$ . Then  $a$  is a simple pole of  $f$  and  $\text{res}_a(f) = \frac{1}{b}$ .

**Proof:** Clearly, if  $l(a) \neq 0$ ,  $a$  is a zero of  $f' + bf^2$ . Hence, a zero  $a$  of  $h'l - hl' + bh^2$  that is not a zero of  $f' + bf^2$  is a pole of  $f$ . Now, when  $l(a) = 0$ , we have  $h(a) \neq 0$  hence  $l'(a) = bh(a) \neq 0$  and therefore  $a$  is a simple pole of  $f$  such that  $\frac{h(a)}{l'(a)} = \frac{1}{b}$ . But since  $a$  is a simple pole of  $f$ , we have  $\text{res}_a(f) = \frac{h(a)}{l'(a)}$  which ends the proof.

**Proof of Theorem 3:** Let  $f = \frac{P}{l}$  with  $P$  a polynomial,  $l \in \mathcal{A}(K)$  having no common zero with  $P$ . Then  $f' + bf^2 = \frac{P'l - l'P + bP^2}{l^2}$ . By hypothesis, this function has finitely many zeroes. Moreover, if  $a$  is a zero of  $P'l - l'P + bP^2$  but is not a zero of  $f' + bf^2$  then, by Lemma 2,  $a$  is a simple pole of  $f$  such that  $\text{res}_a(f) = \frac{1}{b}$ . Consequently,  $P'l - l'P + bP^2$  has finitely many zeroes and so we may write  $\frac{P'l - l'P + bP^2}{l^2} = \frac{Q}{l^2}$  with  $Q \in K[x]$ , hence  $P'l - l'P = -bP^2 + Q$ . But then, by Theorem 1,  $l$  is a polynomial, which ends the proof.

**Proof of Theorem 4:** Let  $g = \frac{f'}{f^2} + b$ . Suppose  $b = 0$ . Since all zeroes of  $f$  are simple zeroes except maybe finitely many,  $g$  has finitely many poles of order  $\geq 3$ , hence a primitive  $G$  of  $g$  has finitely many multiple poles. Consequently, by Theorem 2,  $g$  has infinitely many zeroes.

Now, suppose  $b \neq 0$ . Let  $\alpha$  be a zero of  $g$  and let  $f = \frac{h}{l}$  with  $h, l \in \mathcal{A}(K)$  having no common zero. Then  $\frac{f'}{f^2} + b = \frac{h'l - hl' + bh^2}{h^2}$ . Since  $\alpha$  is a zero of  $\frac{f'}{f^2} + b$ , it is not a zero of  $h$  and hence it is a zero of  $h'l - hl' + bh^2$ . Then by Lemma 2, if it is not a zero of  $f' + bf^2$ , it is a simple pole of  $f$  such that  $\text{res}_\alpha(f) = \frac{1}{b}$ , which ends the proof of Theorem 4.

## References.

- [1] **W. Bergweiler and X. C. Pang** *On the derivative of meromorphic functions with multiple zeros*, J. Math. Anal. Appl. 278 , pp. 285-292 (2003).
- [2] **K. Boussaf and J. Ojeda** *Value distribution of  $p$ -adic meromorphic functions*, to appear in the Bulletin of the Belgian Mathematical Society - Simon Stevin.
- [3] **K. Boussaf** *Picard values of  $p$ -adic meromorphic function*,  $p$ -adic Numbers, Ultrametric Analysis and Applications. Vol. 2, N. 4, pp. 285-292 (2010).
- [4] **A. Escassut** *Analytic Elements in  $p$ -adic Analysis*. World Scientific Publishing Co. Pte. Ltd. (Singapore, 1995).
- [5] **A. Escassut**  *$p$ -adic Value Distribution*. Some Topics on Value Distribution and Differentiability in Complex and  $P$ -adic Analysis, pp. 42-138. Mathematics Monograph, Series 11. Science Press (Beijing, 2008)
- [6] **A. Escassut and J. Ojeda** *Exceptional values of  $p$ -adic analytic functions and derivatives*. Complex Variables and Elliptic Equations. Vol 56, N. 1-4, pp. 263-269 (2011).
- [7] **W. Hayman** *Picard values of meromorphic functions and their derivatives*, Annals of Mathematics. Vol. 70, N. 1, pp. 9-42 (1959).
- [8] **P. C. Hu and C. C. Yang** *Meromorphic Functions over non-Archimedean Fields*. Kluwer Academic Publishers (2000).
- [9] **J. Ojeda** *Distribution de valeurs des fonctions méromorphes ultramétrique, applications de la Théorie de Nevanlinna*. Thèse (N. d'ordre D.U. 1865). Université Blaise Pascal, Clermont-Ferrand, France (2008).
- [10] **J. Ojeda** *Hayman's Conjecture over a  $p$ -adic field*. Taiwanese Journal of Mathematics. Vol. 12, N. 9, pp. 2295-2313 (2008).

Jacqueline Ojeda  
Departamento de Matemática  
Facultad de Ciencias Físicas y Matemáticas  
Universidad de Concepción  
Concepción  
Chile  
jacqojeda@udec.cl

Kamal Boussaf and Alain Escassut  
Laboratoire de Mathématiques  
Université Blaise Pascal  
24 Avenue des Landais  
63171 AUBIERE CEDEX, France

Kamal.Boussaf@math.univ-bpclermont.fr  
Alain.Escassut@math.univ-bpclermont.fr