

Electrical conductivity of hydrous silicate melts: Implications for the bottom-up hydration of Earth's upper mantle

D. Freitas, Geeth Manthilake

▶ To cite this version:

D. Freitas, Geeth Manthilake. Electrical conductivity of hydrous silicate melts: Implications for the bottom-up hydration of Earth's upper mantle. Earth and Planetary Science Letters, 2019, 523, pp.115712. 10.1016/j.epsl.2019.115712. hal-02455958

HAL Id: hal-02455958 https://uca.hal.science/hal-02455958

Submitted on 16 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Electrical conductivity of hydrous silicate melts: Implications for the bottom-up hydration of
2	Earth's upper mantle
3	D.Freitas ^{1*} , G. Manthilake ¹
4	¹ Université Clermont Auvergne, CNRS, IRD, OPGC, Laboratoire Magmas et Volcans,
5	F-63000 Clermont-Ferrand, France
6	(*corresponding author: damien.freitas@uca.fr)
7	

8 Abstract:

9 The upwelling of the hydrous mantle transition zone triggers dehydration-induced partial melting atop the 410-km discontinuity. Here we investigate the electrical conductivity 10 of hydrous silicate melts in the 200-400 km depth range and explore whether melting at the 11 12 410-km depths is responsible for the hydration of the upper mantle. Our experimental electrical conductivity data demonstrate that the mantle at 180-350 km depths is mostly melt 13 free, confirming the H₂O under-saturated conditions. However, the residual mantle from 14 15 partial melting atop the 410km discontinuity may contain various possible amounts of water according to the initial mantle transition zone and melt concentrations. This residual H₂O 16 could contribute to the hydration of the upper mantle either through diffusion or material 17 replacement by upwelling. Our calculations suggest that the diffusion may not be responsible 18 for the hydration of the upper mantle to present H₂O concentration of 50–200 ppm wt. 19 20 Melting of the upwelling mantle transition zone with less than 1500 ppm wt. H₂O produces residual peridotites with ~ 200 ppm H_2O at the 410-km discontinuity. Continuous upwelling 21 of such hydrous residues would gradually replace the dry upper mantle with depleted residual 22 23 hydrous peridotites in less than 260 Ma. In this study, we propose a bottom-up hydration mechanism for the Earth's upper mantle driven by dehydration-melting at the 410-km 24

discontinuity. The hydrous partial melting at the top of the asthenosphere appears to be a consequence of H_2O saturation in the upwelling residual peridotites.

27

Keywords: electrical conductivity, water storage capacity, hydrous melt, upper mantle,
mantle upwelling, mantle transition zone

30

31 **1. Introduction:**

The average H₂O content of the Earth's upper mantle is determined to be about 50-32 200 ppm wt. (Hirschmann, 2006; Michael, 1988; Saal et al., 2002; Simons et al., 2002). 33 34 These estimates are significantly lower than the H₂O storage capacity of peridotite mineral assemblages (Ardia et al., 2012; Demouchy and Bolfan-Casanova, 2016; Férot and Bolfan-35 Casanova, 2012), indicating water undersaturated conditions for the Earth's upper mantle. 36 37 The origin of H₂O in the upper mantle is still a subject of debate. However, two possible scenarios have been proposed. In the first hypothesis, the H₂O in the upper mantle has been 38 39 attributed to hydrous residues left behind following partial melting in the upper mantle. In the second scenario, H₂O in the upper mantle can be derived from the mixing of dry upper 40 mantle with a hydrous primitive source (Hirschmann, 2006). 41

At present, three distinct partial melting regimes, that could explain the origin of H_2O in the upper mantle have been proposed; dehydration melting at the 410-km discontinuity (Bercovici and Karato, 2003), melting in mantle wedges (Hirth and Kohlstedt, 1995) and partial melting of mantle plumes (Morgan and Morgan, 1999), all arguing that the residues left behind by the melting events could be responsible for H_2O contents in the upper mantle. While the depleted signature of mid oceanic ridge basalt (MORB)-source upper mantle strongly favors the partial melting hypotheses, the exact nature of the melting regime that led 49 to the hydration of the upper mantle, consistent with geochemical, geophysical and
50 geodynamic observations, remains poorly constrained (Hirschmann, 2006).

A recent study based on in situ sound velocity measurements confirmed the 51 dehydration induced melting in the upwelling mantle across the 410-km discontinuity 52 (Freitas et al., 2017). Geophysical observations of high electrical conductivity (up to 1 S/m) 53 (Toffelmier and Tyburczy, 2007) and reduced seismic wave velocities (\deltaVs 4-9%) 54 (Revenaugh and Sipkin, 1994; Tauzin et al., 2010) suggest a 50-60 km thick melt layer above 55 the Mantle Transition Zone (MTZ) (Revenaugh and Sipkin, 1994; Song et al., 2004; Tauzin 56 57 et al., 2013, 2010; Vinnik and Farra, 2007). The upwelling of hydrous MTZ with ~ 0.1-0.2 wt.% of H₂O (Freitas et al., 2017; Huang et al., 2005; Karato, 2011) and the significant 58 differences in H₂O storage capacities between the MTZ (Bolfan-Casanova et al., 2000; 59 Hirschmann et al., 2005; Kohlstedt et al., 1996) and the overlying mantle (Demouchy and 60 Bolfan-Casanova, 2016; Férot and Bolfan-Casanova, 2012) is the governing factor that 61 triggers melting in the upwelling mantle across the MTZ (Bercovici and Karato, 2003; Freitas 62 et al., 2017). The reported 3.6-9.4 % shear wave velocity drop is compatible with a 0.5-2.5 % 63 melt fraction (Freitas et al., 2017) in the partial melt layer. 64

The melt produced at the 410-km discontinuity is found to have high H₂O contents 65 (Freitas et al., 2017; Matsukage et al., 2005). The mineral-melt equilibrium following melting 66 at the top of the MTZ causes H₂O to diffuse from the hydrous melt into solid mineral phases 67 both above and below the melt layer. The H₂O diffusion may occur between melt and 68 wadsleyite, transferring H₂O back into the underlying MTZ and between melt-69 olivine/pyroxene/garnet, hydrating the overlying mantle. In addition to the diffusion process, 70 71 the mantle upwells passively away from subduction zones in response to intrusion of slab material into Earth's interior (Bercovici and Karato, 2003; Karato et al., 2006). The mantle 72 upwelling is significantly faster than H₂O diffusion into mantle minerals (Bercovici and 73

Karato, 2003), replacing the dry upper mantle with residual peridotites. The dehydration
melting in the upwelling mantle and subsequent H₂O transport mechanisms therefore play a
crucial role in water transfer between the upper mantle and the mantle transition zone
(Bercovici and Karato, 2003).

Laboratory measurements of electrical conductivity have been extremely useful in 78 characterizing the material circulation in the Earth's mantle (Dai and Karato, 2009, Yoshino, 79 80 2010). The laboratory measurements suggest electrical conductivity is a powerful tool for detecting melt or fluid and their occurrence within the solid mineral matrix at deep mantle 81 82 conditions (Freitas et al., 2019; Grotenhuis et al., 2005; Manthilake et al., 2016, 2015; Maumus et al., 2005). It is also sensitive to the chemical composition of melt, particularly to 83 volatiles (Gaillard et al., 2008; Karato, 1990; Manthilake et al., 2009; Yoshino, 2010) and 84 85 alkalis (Ni et al., 2011). In this study, we measure the electrical conductivity of hydrous peridotite at various pressures up to 12 GPa above their melting temperatures. The results 86 were compared with magnetotelluric profiles to gain insight into the extent of melting and 87 88 melt percolation if hydrous conditions are assumed for the Earth's upper mantle. Based on our results, we discuss different scenarios that could be responsible for the H₂O circulation in 89 the upper mantle. 90

91

92 2. Materials and Methods

93 2.1 Starting materials

Starting materials with a composition similar to KLB-1 peridotite (Wang and Takahashi, 2000) were prepared using reagent grade oxides, initially mixed in the absence of the required amount of Al(OH)₃ to yield the correct mineral composition (ranging from 1.63 to 4.86 mg to obtain 500 mg of peridotite powder with 400 to 1500 ppm H₂O, respectively). These powders were dried at 300 °C overnight to remove any adsorbed moisture. Na and K were added as NaCO₃ and K₂CO₃ and then de-carbonated. The decarbonation of the oxide mixtures was carried out by slowly increasing the temperature to 1000°C (1.6°C/min) and keeping it at 1000°C for about 10 hours to ensure complete decarbonation. The decarbonated powder mixture was then cooled to 200°C and stored in a high vacuum furnace at 120°C prior to the hot pressing runs. The required amount of Al(OH)₃ was then mixed with the decarbonated powder mixture to obtain the desired amount of H₂O in peridotite samples.

105 Samples with ~0, 500, 700 and 1200 ppm wt. H₂O as initial content in powders were prepared in order to have hydrous under saturated conditions at the different targeted 106 107 pressures. As water solubility increases with pressure in nominally hydrous phases (Ardia et al., 2012; Férot and Bolfan-Casanova, 2012), we always used initial water content under 108 solubility limit at each pressure. However, due to the absorbed moisture in the high pressure 109 110 synthesis assembly, the water content may be subjected to some possible water (H) diffusion into the samples even with sealed capsules. The X-Rays powder diffraction analyses using a 111 Philips PW 1830 (Cobalt wave-length) and micro-Raman analyses were carried out prior to 112 the electrical conductivity measurements to ensure the absence of additional hydrous phases 113 such as super hydrous Phase-B and partial melts in pre-sintered samples. 114

115

116 2.2 Experimental

The resulting powder mixtures were hot pressed to obtain solid sintered samples for electrical conductivity measurements. Hydrous and dry samples were synthesized at different pressures and temperature conditions in a first high pressure run with multi-anvil apparatus, these samples were recovered for *in situ* measurements that were performed in a second multi-anvil run. The details on samples conditions (pressure, temperature and time at temperature) are given in the table S1. Dry samples were synthesized at target or lower pressures to ensure low water solubility of the material (Ferot and Bolfan 2012, Kohlstedt et al. 1996). Hydrous samples were synthesized at target pressure (for 6 and 9 GPa) and 14 GPa.
The synthesis of hydrous samples at 14 GPa performing *in situ* experiments at 12 GPa allows
us to simulate the dehydration induced melting at the 410-km discontinuity (Freitas et al.
2017).

Dry samples and hydrous samples were synthetized for ~2 hours in Gold-Palladium (Au-Pd) capsules and temperatures between 1373 and 1573 K (table S1). These under saturated conditions are required during the synthesis process, in order to avoid hydrous silicate melts along grain boundaries, which could otherwise interfere with the electrical conductivity. Cylindrical core samples ~1.0 mm in length and ~1.2 mm in diameter were prepared from these pre-synthesized sample specimens.

The high-pressure and high-temperature experiments were performed using a 1500-134 ton Kawai-type multi-anvil apparatus at the Laboratoire Magmas et Volcans, Clermont-135 Ferrand, France. Both sample synthesis and the in situ measurements were performed in 136 multi-anvil apparatus using an octahedral pressure medium composed of MgO and Cr₂O₃ in a 137 14/8 multi-anvil configuration (octahedron edge length / anvil truncation edge length) for 138 experiments at 12 GPa and 18/11 for experiments at 6 and 9 GPa (Supplementary Figure S1). 139 The pre-synthesized cylindrical sample was inserted into a MgO sleeve. The sleeve also helps 140 insulate the sample electrically from the furnace. This furnace, composed of a 25 µm thick 141 cylindrical Re foil, has apertures for the electrode and the thermocouple wires. A zirconia 142 143 sleeve around the furnace was used as a thermal insulator. Thermal gradient was also taken in to account for the design of these experiments (Hernlund et al., 2006), and were estimated to 144 be negligible $<5^{\circ}$ with our sample dimensions (Supplementary Figure S2). We placed two 145 electrodes, made of Mo discs, at the top and bottom of the cylindrical sample. A tungsten-146 rhenium (W95Re5-W74Re26) thermocouple junction was placed at one end of the sample to 147 monitor the temperature. The opposite end was connected to a single W₉₅Re₅ wire. We 148

collected impedance spectra between the two $W_{95}Re_5$ wires. All ceramic parts of the cell assembly, including the pressure medium, were fired at 1373 K prior to their assemblage in order to remove any adsorbed moisture, and kept in vacuum furnaces (10⁻² Torr and 150°C) before assembling the experiment. Oxygen fugacity of the sample was not controlled during *in situ* measurements, but was expected to remain below the Mo-Mo₂ buffer.

EC measurements were performed using the ModuLab MTS Impedance/Gain-phase 154 analyzer in the frequency range of 10^{1} - 10^{6} Hz. Polyphasic samples are characterized by a 155 combination of resistor-capacitor/constant phase element (R-C/CPE) circuits and the 156 157 resistance can be obtained by fitting the impedance spectra to appropriate equivalent circuits (Supplementary Figure S3). Once the sample resistance has been determined, conductivity 158 can be calculated using the sample diameter and length. The experimental procedure for 159 160 electrical conductivity measurements is discussed elsewhere (Manthilake et al., 2016, 2015). Importance of a moisture removal procedure is crucial in these experiments and is detailed in 161 Supplementary Figure S3. 162

163

164 2.3 Chemical and micro-textural analyses

The chemical composition and micro-textures of pre-sintered samples and 165 166 experimental run products after electrical conductivity measurements were investigated using a Cameca SX100 electron probe micro analyzer and a Scanning Electron Microscope (SEM) 167 JEOL Jeol JSM-5910 LV, respectively, at the Laboratoire Magmas et Volcans of Clermont-168 Ferrand. Energy-dispersive X-ray spectroscopy (EDS) chemical mapping was used to 169 determine the mineral proportions and chemical compositions by selecting a large area of 170 each phase. The fine melt micro-textures were observed using a ZEISS supra 55VP field 171 emission gun (FEG) SEM at 2MATech, Aubière, France. Major element compositions of 172 solid phases are given in Supplementary Tables S2 and S3. 173

The presence of water (OH) in the crystalline phases of our peridotite samples was 174 qualitatively analyzed both before and after electrical conductivity measurements. These 175 analyzes were performed with micro-Raman spectroscopy (Freitas et al., 2017) 176 (Supplementary Figures S4-S6). Raman spectra were collected using an InVia confocal 177 Raman micro spectrometer, equipped with a 532 nm diode laser (output power of ~140 mW), 178 a Peltier-cooled CCD detector, a motorized XY stage and a Leica DM 2500 M optical 179 microscope, at the Laboratoire Magmas et Volcans, Clermont-Ferrand. The Raman 180 spectroscopy technique was preferred over FTIR analysis mainly due to its capacity to 181 182 analyze both OH and Si bonds in a non-destructive manner and its high spatial resolution allowing the analysis of small objects ($\sim 1 \mu m$). However, due to the unavailability of H₂O 183 calibration curves for pyroxene and garnet, we were not able to perform a complete estimate 184 185 of the bulk H₂O contents of our samples (qualitative analysis only).

186

187 **3. Results**

The melting of hydrous peridotite can be detected by a sudden or discontinuous increase in electrical conductivity at high temperature (Fig. 1). The electrical conductivity of partially molten peridotites varies from 0.1-1 S/m. The activation enthalpies of hydrous peridotites before melting indicate the conduction mechanism is characteristic of electron hopping (small polaron conduction) (Katsura et al., 2009).

The high resolution image analyses of samples after electrical conductivity measurements indicate the presence of melt in hydrous samples (Fig. 2). The experimental run products indicate well developed polygonal textures with average grain size of $<30 \mu m$. The melt distribution appears to be homogeneous throughout the samples and occurs as thin films along grains boundaries and as small pockets of a few hundred nanometers in size at triple junctions (Fig. 2 b, c, d). The melt forms an interconnected network showing a very low
dihedral angle with complete wetting of the grain boundaries (Freitas et al., 2017) (Fig 3).

Due to the extremely small size of the melt pockets (<1 μ m), we were not able to measure the chemical composition of the resulting hydrous silicate melt at 6 and 9 GPa. The melt fraction appears to be low (up to a few %). The chemical analyses of melt at 12 GPa indicate significant enrichment in incompatibles elements (Ca, Al, Na, K, Fe, Ti) (Freitas et al., 2017).

Samples synthesized prior to electrical conductivity measurements were melt-free 205 206 (confirmed by X-ray powder diffraction/micro-Raman) and their mineralogy was mainly composed of olivine, clinopyroxene and garnet (Fig. 2a). No evidence of hydrous phases such 207 as super-hydrous B were found in the starting materials. The proportions of the different 208209 crystalline phases vary between the samples depending on the degree of melting. The starting samples recovered after synthesis at 12 GPa have mineral proportions of about 60±2% 210 olivine, 20±2% garnet and 20±2% clinopyroxene. Upon partial melting at 12 GPa, the 211 proportion of olivine remained almost constant, while the proportion of Cpx decreased with 212 increasing melt fractions. The proportion of garnet slightly increased with melting, as would 213 be expected with mantle mineralogy (Kaminsky, 2012). For all experiments, the Mg number 214 of olivine is between 90 and 93. Nominally anhydrous sample have lower Mg numbers of 88-215 89, which is consistent with melt-absent conditions. Chemistry of solid phases for synthesis 216 217 and experimental runs are given in Supplementary Tables S2 and S3.

218

219 4. Discussion

220 4.1 Melting in the upper mantle

The magnetotelluric and geomagnetic depth sounding profiles of the upper mantle indicate significant regional variation in electrical conductivity (Grayver et al., 2017;

Kuvshinov et al., 2005; Toffelmier and Tyburczy, 2007). The majority of geophysical 223 profiles are compatible with electrical conductivity resulting from hydrous upper mantle 224 minerals (Novella et al., 2017b). The comparison of these electrical conductivities of hydrous 225 silicate melts with global electromagnetic and magnetotelluric profiles of the upper mantle 226 implies that the upper mantle in the depth range 180-350 km is mainly melt free (Fig. 4). 227 However, the estimated high electrical conductivity anomaly above the 410-km discontinuity 228 (Toffelmier and Tyburczy, 2007) appears to be compatible with the presence of melt in this 229 zone (Fig. 4). Thanks to our measurements, the melt fraction required to explain this 230 231 observed anomaly is estimated to be <10 vol. % (Fig. 4). This estimation is significantly higher than the melt fraction of 0.5-2.5 % obtained by sound velocity measurements (Freitas 232 et al., 2017) to explain the shear velocity reduction (\deltaVs) of 3.6-9.4 %. The possible over-233 234 estimation of melt fraction by laboratory electrical conductivity methods has been discussed in a recent study (Freitas et al., 2019) in addition to assumptions made on forward and 235 inversion models of electromagnetic data (Toffelmier and Tyburczy, 2007). In our discussion, 236 we use our electrical conductivity data to qualitatively determine the vertical extents of 237 melting in the upper mantle. 238

The absence of melt in the depth range 180-350 km confirms that the water content of 239 the upper mantle is below the saturation limits of peridotite (Ardia et al., 2012; Férot and 240 Bolfan-Casanova, 2012). This water undersaturated mid-upper mantle is sandwiched between 241 242 two partial melt layers. Melting at the top of the asthenosphere (above 180 km) has been discussed in terms of the reduced water solubility of aluminous orthopyroxene (Mierdel et al., 243 2007), and melting below 350 km has been linked to the reduced water solubility in 244 peridotites, in both cases melting is caused by the presence of excess H₂O. Here we 245 investigate whether the dehydration-induced water saturation and subsequent melting at the 246

top of the mantle transition zone has an influence on water circulation in the Earth's uppermantle.

249

$4.2 H_2O$ circulation in the upper mantle

The strong evidences supporting the presence of a melt layer atop the mantle 251 transition zone and the strict requirement of H₂O for deep mantle melting underline the 252 importance of understanding the mechanism that is responsible for H₂O circulation in the 253 upper mantle. Assuming that the mantle was initially dry and well mixed after the 254 255 establishment of the vigorously convecting global magma ocean, initiation of subduction in a plate tectonic context can be considered as having been the major process responsible for 256 mantle hydration. With average downward slab velocities of between 5 and 10 cm per year 257 258 (Gordon, 1995a) and assuming plates thicknesses of 7 km and a total ridge length of 70000 km, the annual amount of subducted material is estimated to be 24-49 km (Smyth and 259 Jacobsen, 2006). With a mean age of 80 Ma, the average subducted material is estimated to 260 have between 0.1 and 1.5% of H₂O (Dixon et al., 2002) leading to an H₂O flux of 0.1 to 1 261 km³ into the mantle per year. However, studies of ophiolites indicate that most hydrous 262 minerals break down at shallow mantle conditions and subducted rocks in eclogite facies 263 were shown to be significantly impoverished in volatiles, with values of up to 0.25% H₂O 264 (Katayama and Nakashima, 2003). Even in the case of extreme slab dehydration of up to 97% 265 266 (Dixon et al., 2002), H₂O contents of up to 1000 ppm wt. can be preserved in subducting slabs and transported down to the transition zone (Dixon et al., 2002). 267

Using slab downward speed of 5 cm per year and the MTZ thickness of 250 km, the time required for a portion of slab to cross the MTZ is estimated to be 5 Ma. While this time is relatively short on a planetary timescale, the slab deflection and stagnation in the MTZ observed in seismic tomography studies (Zhao, 2004) suggests that residence time of

subducted material would be much greater than the values assumed in our calculations. With a diffusion coefficient of 10^{-8} - 10^{-10} m²/s (Hae et al., 2006; Richard et al., 2006), a significant transfer of H₂O from the slab to the MTZ could be expected during that time.

The appearance of the melt layer above the MTZ can be inferred given the estimation 275 of H₂O slab fluxes in the mantle transition zone. If we assume 24.5 km³ slab material reaches 276 the MTZ per year (volume produced at the ridges for 5 cm/year spreading velocity along 277 70000 km of ridges with an average thickness of 7 km (Gordon, 1995b)), and that these rocks 278 contain 1000 ppm wt. H₂O, a mass of 9.4×10^{10} kg of H₂O is transferred to the MTZ every 279 year. Assuming the volume of the MTZ to be 6.5×10^7 km³, with a mass of 2.6×10^{20} kg (mean 280 density of 4000 kg/m³), and assuming complete slab dehydration into the MTZ, only 2.2-3.3 281 Ma after the beginning of modern subduction is required to reach 800-1200 ppm wt. in the 282 MTZ, which is roughly equal to the maximum H₂O solubility in nominally anhydrous 283 minerals (NAM's) forming the upper mantle (Ardia et al., 2012; Demouchy and Bolfan-284 Casanova, 2016; Férot and Bolfan-Casanova, 2012). Using the upper limit of the inferred 285 H₂O content of the mantle transition zone from geophysical methods of about 0.1-0.2 wt.% 286 H₂O (Freitas et al., 2017; Huang et al., 2005), the time required to transport H₂O from surface 287 to the MTZ is estimated to be 6 ± 0.6 Ma. This very simple estimation highlights the 288 efficiency of the hydration of the MTZ. It indicates clearly that melting at the 410-km 289 discontinuity would have been initiated early in the Earth's history. However, H₂O diffusion 290 291 into wadsleyite is a slow process (Hae et al., 2006). For example, the time required to diffuse 2000 ppm wt. H₂O to reach 500 ppm wt. at 500 km laterally along the MTZ is estimated to be 292 175 Ga. Given that the surface area of the MTZ is significantly large and the number of 293 294 active subduction zones is extremely limited, it is possible that the H₂O distribution in the MTZ remains heterogeneous, with regions of the MTZ maintaining dry conditions (Yoshino 295 et al., 2008). The absence of seismic anomalies in some regions (Tauzin et al., 2010) 296

indicates that melting is not pervasive above the MTZ. The upwelling of H_2O deprived regions (< 800-1200 ppm wt.) of the MTZ may not undergo partial melting upon entering the upper mantle.

The H₂O under-saturated MTZ, as demonstrated by recent experimental studies 300 (Freitas et al., 2017; Huang et al., 2005; Karato, 2011), can be a result of continuous removal 301 of H₂O from the MTZ (upward and/or downward) or partial dehydration of slabs at the MTZ. 302 The first scenario implies the presence of a global H₂O circulation cycle extracting H₂O from 303 the MTZ from an early stage. However, the quantity of H₂O stored in the melt layer above 304 305 the MTZ together with the moderately hydrous olivine present in equilibrium with melt over the 60 km thick layer, only represents a small portion of the total amount of H_2O in the MTZ 306 (<1%). This amount is a few orders of magnitude lower than the total amount transported by 307 308 subducted slabs per year. Similarly, the known deep volcanic sources of OIB show that source material may be only moderately hydrous with H₂O content estimated between 300 309 and 1000 ppm wt. (Rüpke et al., 2006; Simons et al., 2002). These calculations demonstrate 310 311 the possibility of large scale melting events occurring periodically in the Earth's history. Large scale deep mantle melting has occurred in the past, as demonstrated by the presence of 312 komatiites, kimberlites and numerous large igneous provinces (Vinnik and Farra, 2007). 313 However, at present, such large scale volcanism does not exist anymore and hot spot 314 volcanism and mantle plumes are alternative candidates for removing H₂O from the deep 315 316 Earth.

On the other hand, it also appears possible that slabs that are not fully dehydrated at the MTZ, and instead carry a substantial amount of H_2O on down to the lower mantle (Dixon et al., 2002). The high H_2O contents of up to 2 wt. % observed in some enriched mantle sources of oceanic island basalts (OIB) indicate that some H_2O may be incorporated into dense hydrous silicate phases (Nishi et al., 2014; Pamato et al., 2014) that penetrates down

into the lower mantle via subduction processes (Rüpke et al., 2006). These would imply a
slower hydration rate of the MTZ and more tardive apparition of the LVL.

324

325 *4.3 Bottom-up hydration of the upper mantle*

In this section, different possible mechanisms of bottom–up hydration of Earth's upper mantle and their consequences are explored.

Due to the H₂O under-saturated conditions of the upper mantle, the water contained in hydrous melts produced at the 410-km discontinuity may diffuse in to the overlying mantle. Based on the one dimensional diffusion equation (Crank, 1975), we calculate the diffusion time required to hydrate the upper mantle assuming that the upper mantle is initially dry (Supplementary Figure S7),

$$C(x,t) = C_0 + (C_1 - C_0) * erfc\left(\frac{x}{2\sqrt{D * t}}\right)$$

where C_0 is the initial concentration, C_1 is constant local concentration, x is the distance to the 334 border of concentration C_1 , t is the diffusion time and D the diffusion coefficient of hydrogen 335 in olivine. The value for D was estimated to be $7.82 \times 10^{-9} \pm 6.1 \times 10^{-8}$ m²/s at 1723 K for a 336 randomly oriented polycrystalline material (average of the three crystallographic direction) 337 (Novella et al., 2017b). Using average H₂O solubility for mantle (Demouchy and Bolfan-338 339 Casanova, 2016; Férot and Bolfan-Casanova, 2012), increasing from 300 ppm wt. at 200 km up to 1000 ppm wt. at 410 km, we calculate the time required for peridotite in equilibrium 340 with the hydrous melt layer to reach H₂O saturation. Our results indicate that the time 341 required to reach the H₂O saturation in the uppermost mantle is unrealistically high based on 342 diffusion alone, and cannot be achieved during the planetary life time (Supplementary Figure 343 S7, Fig. 5a). 344

On the other hand, the mantle away from subducting slabs is rising passively in response to the material injection by subduction. Estimated values using slab flux yield to upwelling velocities of 1 mm per year (Bercovici and Karato, 2003). The estimation based on convection models indicates convection rate as high as 1.5 cm/year (Becker et al., 1999). The upwelling rates from subduction fluxes indicates that only 260 Ma is sufficient to bring rocks from the top of the transition zone to the uppermost mantle (150 km depth) and even faster considering the highest convection rates. This estimation indicates that residual hydrous peridotite should be rising from the LVL to the uppermost mantle and saturates the upper mantle at various depths depending on the initial H₂O contents in peridotite (Fig. 5b).

Here we investigate the possible H₂O contents of peridotites in contact with the melt layer. The shear wave velocity reduction (δVs) at the melt layer is estimated to vary between 3.6-9.4 % (Revenaugh and Sipkin, 1994; Tauzin et al., 2010). Based on our previous study (Freitas et al., 2017), the resulting melt fraction corresponding to the δVs at the low velocity layer is estimated to be 0.5-2.5 vol.%

The average water content of the MTZ has been constrained by different methods 359 with variable results. Despite potential high water solubilities of MTZ minerals (Bolfan-360 Casaonova et al., 2018b), the actual content of the MTZ is believed to be lower than 361 saturation limit. Water contents inferred from inclusions in diamonds (Pearson et al., 2014) 362 with up to 2 % would suggest potentially high water content which also supported by 363 viscosity measurements (Fei et al. 2017). Particularly, the near-water saturated conditions 364 implied by hydrous ringwoodite inclusions found in natural diamond (Pearson et al., 2014) 365 may represent an episode of subduction-zone-related melting. However, most of geophysical 366 techniques such as elasticity, sound velocities (Inoue et al., 1998, Chang et al., 2015, Freitas 367 et al., 2017) and electrical conductivities (Huang et al., 2005, Yoshino et al., 2008, Yoshino 368 and Katsura 2012) indicating major part of MTZ contains < 0.2 wt. % H₂O. 369

Assuming a batch melting scenario and H₂O partitioning between peridotite and melt (Novella et al., 2014), readjusted to the mineral proportions expected at 12 GPa

(Supplementary Text S1), we could estimate the H₂O contents of peridotite in equilibrium 372 with hydrous melt for variable water contents in the mantle source for different melt fractions 373 (Fig. 6). Different scenarios could occur given the potential range of water content of the 374 residual peridotites (Fig. 6). Thus, if residual peridotite in the upwelling mantle contains 375 about 200 ppm wt. of H₂O, mantle saturation can be expected at range of depths in the upper 376 mantle (Fig. 6). If we assume the H₂O saturation curves of Férot and Bolfan-Casanova (Férot 377 378 and Bolfan-Casanova, 2012), Hirschmann et al. (Hirschmann et al., 2009), Tenner et al. (Tenner et al., 2012, 2009) and Ardia et al. (Ardia et al., 2012) (for garnet/olivine partitioning 379 380 of H₂O of 9.0), peridotite in the upper mantle reaches H₂O saturation limits above 200 km. In contrast, by using the extreme H₂O saturation scenario proposed by Ardia et al. (Ardia et al., 381 2012) with water partitioning between olivine and garnet of 0.9, the mantle saturation above 382 200 km depths can be expected for the water contents of 450 ppm. However, this value is 383 inconstant with the upper mantle water contents of 50-200 ppm. 384

The ascent of residual hydrous peridotites would initiate melting upon reaching their 385 storage capacities and could be responsible for the hydrous melting at the top of the 386 asthenosphere. Melting above a depth of 180 km can only be explained by moderate water 387 content in the MTZ (< 1500 ppm wt) for geologically relevant melt fractions at the 410-km 388 discontinuity (Fig. 6). The upwelling of the MTZ with water contents of less than 800 ppm 389 wt. may not undergo melting upon entering the upper mantle and may continue to hydrate the 390 391 upper mantle. However, such fertile peridotites are not able to explain the depleted signature of the upper mantle. Melting at the 410- km discontinuity is therefore a primary requirement 392 for a compatible hydration of the upper mantle with geochemical constrains. 393

The lithosphere-asthenosphere boundary (LAB) is defined as intersection of the conductive geotherm of mechanically strong crust with the fluid-saturated peridotite solidus in mechanically weak upper most part of the mantle (McKenzie and Bickle, 1988). The

location of the LAB is therefore corresponding to the depth at which the fluid saturation 397 occurs in upwelling peridotites. The geochemical constraints suggest that the LAB is not a 398 399 stable feature, which appear to move up or down over geological time scale (O'Reilly and Griffin, 2010). If other parameters remain constant, the lateral variation of the depth of the 400 LAB may indicate the variable water contents in residual peridotites, which corresponds to 401 the lateral variations of water contents in the MTZ. Further, the movement of the depth of the 402 403 LAB at the same location may suggest episodic variation of water contents in the MTZ over geological time scale. The upwelling of MTZ with a water content greater than 2000 ppm wt. 404 405 ppm could produce residual peridotites reaching the maximum water storage capacity (Fig. 6). The upwelling of such peridotites could continue to melt and form a gravitationally 406 unstable melt column in the upper mantle. The upwelling of the MTZ with high water 407 408 contents would trigger large-scale melting (Vinnik and Farra, 2007) and could be responsible for the formation of upper mantle plumes. 409

410

411 **5. Conclusions**

Based on our data, we propose a hydration mechanism for the Earth's upper mantle 412 that is consistent with geophysical observations (magnetotelluric and seismic profiles) and 413 geochemical constraints. In this bottom-up water transportation mechanism, the residual 414 hydrous peridotites, formed as a result of melting at the 410-km discontinuity, upwell across 415 416 the upper mantle, replacing initially dry mantle peridotites. The resulting water content in the residues is directly related to the water content of the mantle transition zone and the degree of 417 melting. The upper limit of 200 ppm observed for the upper mantle can be maintained in 418 419 residual peridotites if the water content of the MTZ is less than 1500 ppm . The slow H₂O diffusion rates in wadsleyite may enforce large gradient in H₂O distribution within the MTZ, 420 and as a result, regional variations of H₂O contents in the upper mantle would be expected. 421

The ascent of residual hydrous peridotites could be responsible for the hydrous melting at thetop of the asthenosphere.

424

425 Acknowledgements

- 426 We thank F. Schiavi for the assistance with Raman analyses, A. Mathieu for technical
- 427 assistance, J.M Hénot, J.L Devidal and 2MAtech staff for their help in SEM and EMP and

428 FEG-SEM analyses, respectively. We appreciate the discussion with J, Chantel, D, Andrault,

- 429 N. Bolfan-Casanova and M.A Bouhifd. G.M. acknowledges funding from the French PNP
- 430 program (INSU-CNRS). This research was financed by the French Government Laboratory
- 431 of Excellence initiative n°ANR-10-LABX-0006, the Région Auvergne and the European
- 432 Regional Development Fund. This is ClerVolc contribution number xx.
- 433

434 **References**

- Ardia, P., Hirschmann, M.M., Withers, A.C., Tenner, T.J., 2012. H2O storage capacity of
 olivine at 5-8GPa and consequences for dehydration partial melting of the upper mantle.
 Earth Planet. Sci. Lett. 345–348, 104–116. doi:10.1016/j.epsl.2012.05.038
- Becker, T.W., Kellogg, J.B., O' Connell, R.J., 1999. Thermal constraints on the survival of
 primitive blobs in the lower mantle. Earth Planet. Sci. Lett. 171, 351–365.
 doi:10.1016/S0012-821X(99)00160-0
- Bercovici, D., Karato, S.-I., 2003. Whole-mantle convection and the transition-zone water
 filter. Nature 425, 39–44. doi:10.1038/nature01918
- Bolfan-Casanova, N., Keppler, H., Rubie, D.C., 2000. Water partitioning between nominally
 anhydrous minerals in the MgO-SiO2-H2O system up to 24 GPa: Implications for the
 distribution of water in the Earth's mantle. Earth Planet. Sci. Lett. 182, 209–221.
 doi:10.1016/S0012-821X(00)00244-2
- 447 Bolfan-casanova, N., Schiavi, F., Novella, D., Bureau, H., Raepsaet, C., Khodja, H., Demouchy, S., Bolfan-casanova, N., Schiavi, F., Novella, D., Bureau, H., Raepsaet, C., 448 2018. Examination of Water Quantification and Incorporation in Transition Zone 449 Minerals: Wadsleyite, Ringwoodite and Phase D Using ERDA (Elastic Recoil 450 Detection Analysis) To cite this version : HAL Id : cea-01818386 Examination of Water 451 Quantification and Incorporation in Transition Zone Minerals: Wadsleyite 452 Ringwoodite and Phase D Using ERDA (Elastic Recoil Detection Analysis). 453 doi:10.3389/feart.2018.00075 454
- Chang, Y.-Y., Jacobsen, S. D., Bina, C. R., Thomas, S. M., Smyth, J. R., Frost, D. J., et al.
 (2015). Comparative compressibility of hydrous wadsleyite and ringwoodite: effect of H2O and implications for detecting water in the transition zone. J. Geophys. Res. Solid Earth 120, 259–8280.

- 459 Crank, J., 1975. The mathematics of diffusion. Clarendon Press, Oxford.
- 460 Dai, L., Karato, S., 2014. High and highly anisotropic electrical conductivity of the
 461 asthenosphere due to hydrogen diffusion in olivine. Earth Planet. Sci. Lett. 408, 79–86.
 462 doi:10.1016/j.epsl.2014.10.003
- Dai, L., Karato, S., 2009. Electrical conductivity of wadsleyite at high temperatures and high
 pressures. Earth Planet. Sci. Lett. 287, 277–283. doi:10.2138/am.2011.3796
- Demouchy, S., Bolfan-Casanova, N., 2016. Distribution and transport of hydrogen in the
 lithospheric mantle: A review. Lithos 240–243, 402–425.
 doi:10.1016/j.lithos.2015.11.012
- Dixon, J.E., Leist, L., Langmuir, C., Schilling, J., 2002. Recycled dehydrated lithosphere
 observed in plume-influenced mid-ocean-ridge basalt. Nature 420, 385–389.
- Fei, H., Yamazaki, D., Sakurai, M., Miyajima, N., Ohfuji, H., 2017. A nearly water-saturated
 mantle transition zone inferred from mineral viscosity. doi:10.1126/sciadv.1603024
- 472 Férot, A., Bolfan-Casanova, N., 2012. Water storage capacity in olivine and pyroxene to
 473 14GPa: Implications for the water content of the Earth's upper mantle and nature of
 474 seismic discontinuities. Earth Planet. Sci. Lett. 349–350, 218–230.
 475 doi:10.1016/j.epsl.2012.06.022
- Freitas, D., Manthilake, G., Chantel, J., Bouhfid, M., Andrault, D., 2019. Simultaneous
 measurements of electrical conductivity and seismic velocity of partially molten
 geological materials: Implications for melt fraction in the upper mantle. Phys. Chem.
 Miner. 0, 0. doi:10.1007/s00269-019-01021-5
- Freitas, D., Manthilake, G., Schiavi, F., Chantel, J., Bolfan-Casanova, N., Bouhifd, M.A.,
 Andrault, D., 2017. Experimental evidence supporting a global melt layer at the base of
 the Earth's upper mantle. Nat. Commun. 8, 2186. doi:10.1038/s41467-017-02275-9
- Gaillard, F., Malki, M., Iacono-Marziano, G., Pichavant, M., Scaillet, B., 2008. Carbonatite
 melts and electrical conductivity in the asthenosphere. Science (80-.). 322, 1363–1365.
 doi:10.1126/science.1164446
- Gordon, G., 1995a. Plate motions, crustal and lithospheric mobility, and paleomagnetism:
 Prospective viewpoint. J. Geophys. Res. 100, 24367–24392.
- 488 Gordon, G., 1995b. Global Earth Physics: A Handbook of Physical Constants, AGU Washin.
 489 ed.
- Grayver, A. V., Munch, F.D., Kuvshinov, A. V., Khan, A., Sabaka, T.J., Tøffner-Clausen, L.,
 2017. Joint inversion of satellite-detected tidal and magnetospheric signals constrains
 electrical conductivity and water content of the upper mantle and transition zone.
 Geophys. Res. Lett. 44, 6074–6081. doi:10.1002/2017GL073446
- Grotenhuis, S.M., Drury, M.R., Spiers, C.J., Peach, C.J., 2005. Melt distribution in olivine
 rocks based on electrical conductivity measurements 110, 1–11.
 doi:10.1029/2004JB003462
- Hae, R., Ohtani, E., Kubo, T., Koyama, T., Utada, H., 2006. Hydrogen diffusivity in
 wadsleyite and water distribution in the mantle transition zone. Earth Planet. Sci. Lett.
 243, 141–148. doi:10.1016/j.epsl.2005.12.035
- Hernlund, J., Einenweber, K., Locke, D., Tyburczy, J., 2006. A numerical model for steady state temperature distributions in solid-medium high-pressure cell assemblies 91, 295–
 305. doi:10.2138/am.2006.1938
- Hirschmann, M.M., 2006. Water, melting, and the deep Earth H2O cycle. Annu. Rev. Earth
 Planet. Sci. 34, 629–653. doi:10.1146/annurev.earth.34.031405.125211
- Hirschmann, M.M., Aubaud, C., Withers, A.C., 2005. Storage capacity of H2O in nominally
 anhydrous minerals in the upper mantle. Earth Planet. Sci. Lett. 236, 167–181.
 doi:10.1016/j.epsl.2005.04.022
- 508 Hirschmann, M.M., Tenner, T., Aubaud, C., Withers, A.C., 2009. Dehydration melting of

- nominally anhydrous mantle: The primacy of partitioning. Phys. Earth Planet. Inter. 176,
 54–68. doi:10.1016/j.pepi.2009.04.001
- Hirth, G., Kohlstedt, D.L., 1995. Experimental constraints on the dynamics of the partially
 molten upper mantle: Deformation in the diffusion creep regime. J. Geophys. Res. Solid
 Earth 100, 1981–2001. doi:10.1029/94JB02128
- Huang, X., Xu, Y., Karato, S., 2005. Water content in the transition zone from electrical
 conductivity of wadsleyite and ringwoodite. Nature 434, 746–749.
 doi:10.1038/nature03426
- Inoue, T.,Weidner, D. J., Northrup, P. A., and Parise, J. B. (1998). Elastic properties of
 hydrous ringwoodite (?-phase) in Mg2SiO4. Earth Planet. Sci. Lett. 160, 107–113. doi:
 10.1016/S0012-821X(98)00077-6
- Kaminsky, F., 2012. Earth-Science Reviews Mineralogy of the lower mantle : A review of '
 super-deep ' mineral inclusions in diamond. Earth Sci. Rev. 110, 127–147.
 doi:10.1016/j.earscirev.2011.10.005
- Karato, S., 1990. The role of hydrogen in the electrical conductivity of the upper mantle.
 Nature 347, 183–187. doi:10.1038/346183a0
- Karato, S., Bercovici, D., Leahy, G., Richard, G., Jing, Z., 2006. The Transition-Zone Water
 Filter Model for Global Material Circulation: Where Do We Stand? Earth's Deep
 Water Cycle 289–313. doi:10.1029/168GM22
- Karato, S.I., 2011. Water distribution across the mantle transition zone and its implications
 for global material circulation. Earth Planet. Sci. Lett. 301, 413–423.
 doi:10.1016/j.epsl.2010.11.038
- Katayama, I., Nakashima, S., 2003. Hydroxyl in clinopyroxene from the deep subducted
 crust : Evidence for H2O transport into the mantle. Am. Mineral. 88, 229–234.
- Katsura, T., Yoshino, T., Manthilake, G., Matsuzaki, T., 2009. Electrical conductivity of the
 major upper mantle minerals: a review. Russ. Geol. Geophys. 50, 1139–1145.
 doi:10.1016/j.rgg.2009.11.012
- 536 Kohlstedt, D.L., Keppler, H., Rubie, D.C., 1996. Solubility of water in the α , β and γ phases 537 of (Mg,Fe)2SiO4. Contrib. to Mineral. Petrol. 123, 345–357. 538 doi:10.1007/s004100050161
- Kuvshinov, A., Utada, H., Avdeev, D., Koyama, T., 2005. 3-D modelling and analysis of Dst
 C-responses in the North Pacific Ocean region, revisited. Geophys. J. Int. 160, 505–526.
 doi:10.1111/j.1365-246X.2005.02477.x
- Manthilake, G., Bolfan-Casanova, N., Novella, D., Mookherjee, M., Andrault, D., 2016.
 Dehydration of chlorite explains anomalously high electrical conductivity in the mantle
 wedges. Sci. Adv. 2.
- Manthilake, G., Mookherjee, M., Bolfan-Casanova, N., Andrault, D., 2015. Electrical
 conductivity of lawsonite and dehydrating fluids at high pressures and temperatures.
 Geophys. Res. Lett. 42, 7398–7405.
- Manthilake, M. a G.M., Matsuzaki, T., Yoshino, T., Yamashita, S., Ito, E., Katsura, T., 2009.
 Electrical conductivity of wadsleyite as a function of temperature and water content.
 Phys. Earth Planet. Inter. 174, 10–18. doi:10.1016/j.pepi.2008.06.001
- Matsukage, K.N., Jing, Z., Karato, S., 2005. Density of hydrous silicate melt at the conditions
 of Earth's deep upper mantle. Nature 438, 488–91. doi:10.1038/nature04241
- Maumus, J., Bagdassarov, N., Schmeling, H., 2005. Electrical conductivity and partial
 melting of mafic rocks under pressure. Geochim. Cosmochim. Acta 69, 4703–4718.
 doi:10.1016/j.gca.2005.05.010
- McKenzie D., Bickle M. J., 1988. The Volume and Composition of Melt Generated by
 Extension of the Lithosphere. J. Petrol. 29, 625–679.
- 558 Mibe, K., Fujii, T., Yasuda, A., 1999. Control of the location of the volcanic front in island

- arcs by aqueous fluid connectivity in the mantle wedge. Nature 401, 259–262.
 doi:10.1038/45762
- Mibe, K., Fujii, T., Yasuda, A., 1998. Connectivity of aqueous fluid in the Earth's upper
 mantle 25, 1233–1236.
- Michael, P.J., 1988. The concentration, behavior and storage of H20 in the suboceanic upper
 mantle: Implications for mantle metasomatism. Geochim. Cosmochim. Acta 52, 555–
 566.
- Mierdel, K., Keppler, H., Smyth, J.R., Langenhorst, F., 2007. Water Solubility in Aluminous
 Orthopyroxene and the Origin of Earth's Asthenosphere. Science (80-.). 315, 364–368.
 doi:10.1126/science.1135422
- Morgan, J.P., Morgan, W.J., 1999. Two-stage melting and the geochemical evolution of the
 mantle : a recipe for mantle plum-pudding. Earth Planet. Sci. Lett. 170, 215–239.
- Ni, H., Keppler, H., Manthilake, M.A.G.M., Katsura, T., 2011. Electrical conductivity of dry
 and hydrous NaAlSi3O8 glasses and liquids at high pressures. Contrib. to Mineral.
 Petrol. 162, 501–513. doi:10.1007/s00410-011-0608-5
- Nishi, M., Irifune, T., Tsuchiya, J., Tange, Y., Nishihara, Y., Fujino, K., Higo, Y., 2014.
 Stability of hydrous silicate at high pressures and water transport to the deep lower
 mantle 7. doi:10.1038/NGEO2074
- Novella, D., Dolejš, D., Myhill, R., Pamato, M.G., Manthilake, G., Frost, D.J., 2017a.
 Melting phase relations in the systems Mg2SiO4–H2O and MgSiO3–H2O and the
 formation of hydrous melts in the upper mantle. Geochim. Cosmochim. Acta 204, 68–
 82. doi:10.1016/j.gca.2016.12.042
- Novella, D., Frost, D.J., Hauri, E.H., Bureau, H., Raepsaet, C., Roberge, M., 2014. The
 distribution of H2O between silicate melt and nominally anhydrous peridotite and the
 onset of hydrous melting in the deep upper mantle. Earth Planet. Sci. Lett. 400, 1–13.
 doi:10.1016/j.epsl.2014.05.006
- Novella, D., Jacobsen, B., Weber, P.K., Tyburczy, J.A., Ryerson, F.J., Du Frane, W.L.,
 2017b. Hydrogen self-diffusion in single crystal olivine and electrical conductivity of
 the Earth's mantle. Sci. Rep. 7, 1–10. doi:10.1038/s41598-017-05113-6
- O'Reilly, S.Y., Griffin, W.L., 2010. The continental lithosphere-asthenosphere boundary:
 Can we sample it? Lithos 120, 1–13. doi:10.1016/j.lithos.2010.03.016
- Pamato, M.G., Myhill, R., Boffa Ballaran, T., Frost, D.J., Heidelbach, F., Miyajima, N.,
 2014. Lower-mantle water reservoir implied by the extreme stability of a hydrous
 aluminosilicate. Nat. Geosci. 8, 75–79. doi:10.1038/NGEO2306
- Pearson, D.G., Brenker, F.E., Nestola, F., McNeill, J., Nasdala, L., Hutchison, M.T.,
 Matveev, S., Mather, K., Silversmit, G., Schmitz, S., Vekemans, B., Vincze, L., 2014.
 Hydrous mantle transition zone indicated by ringwoodite included within diamond.
 Nature 507, 221–224.
- Revenaugh, J., Sipkin, S., 1994. Seismic evidance for silicate melt atop the 410-km mantle
 discontinuity. Nature 369, 474–476.
- Richard, G., Bercovici, D., Karato, S., 2006. Slab dehydration in the Earth 's mantle
 transition zone 251, 156–167. doi:10.1016/j.epsl.2006.09.006
- Rüpke, L., Morgan, J.P., Dixon, J.E., 2006. Implications of Subduction Rehydration for Earth
 's Deep Water Cycle, in: Jacobsen, S.D., Van Der Lee, S. (Eds.), Earth's Deep Water
 Cycle. American Geophysical Union, pp. 263–276.
- Saal, A.E., Hauri, E.H., Langmuir, C.H., Perfit, M.R., 2002. Vapour undersaturation in
 primitive mid-ocean-ridge basalt and the volatile content of Earth's upper mantle.
 Nature 419, 451–455. doi:10.1038/nature01073
- Simons, K., Dixon, J., Schilling, J.-G., Kingsley, R., Poreda, R., 2002. Volatiles in basaltic
 glasses from the Easter-Salas y Gomez Seamount Chain and Easter Microplate:

- Implications for geochemical cycling of volatile elements. Geochemistry, Geophys.
 Geosystems 3, 1–29. doi:10.1029/2001GC000173
- Smyth, J.R., Jacobsen, S.D., 2006. Nominally Anhydrous Minerals and Earth's Deep Water
 Cycle, in: Jacobsen, S.D., Van Der Lee, S. (Eds.), Earth's Deep Water Cycle,
 Geophysical Monograph Series. American Geophysical Union.
 doi:doi:10.1029/168GM02
- Song, T.-R.A., Helmberger, D. V, Grand, S.P., 2004. Low-velocity zone atop the 410-km
 seismic discontinuity in the northwestern United States. Nature 427, 530–533.
 doi:10.1038/nature02231
- Tauzin, B., Debayle, E., Wittlinger, G., 2010. Seismic evidence for a global low-velocity
 layer within the Earth's upper mantle. Nat. Geosci. 3, 718–721. doi:10.1038/ngeo969
- Tauzin, B., van der Hilst, R.D., Wittlinger, G., Ricard, Y., 2013. Multiple transition zone
 seismic discontinuities and low velocity layers below western United States. J. Geophys.
 Res. Solid Earth 118, 2307–2322. doi:10.1002/jgrb.50182
- Tenner, T.J., Hirschmann, M.M., Withers, A.C., Ardia, P., 2012. H2O storage capacity of 623 olivine and low-Ca pyroxene from 10 to 13 GPa: Consequences for dehydration melting 624 625 above the transition zone. Contrib. to Mineral. Petrol. 163, 297-316. doi:10.1007/s00410-011-0675-7 626
- Tenner, T.J., Hirschmann, M.M., Withers, A.C., Hervig, R.L., 2009. Hydrogen partitioning
 between nominally anhydrous upper mantle minerals and melt between 3 and 5 GPa and
 applications to hydrous peridotite partial melting. Chem. Geol. 262, 42–56.
 doi:10.1016/j.chemgeo.2008.12.006
- Toffelmier, D.A., Tyburczy, J.A., 2007. Electromagnetic detection of a 410-km-deep melt
 layer in the southwestern United States. Nature 447, 991–994. doi:10.1038/nature05922
- Vinnik, L., Farra, V., 2007. Low S velocity atop the 410-km discontinuity and mantle
 plumes. Earth Planet. Sci. Lett. 262, 398–412. doi:10.1016/j.epsl.2007.07.051
- Wang, W., Takahashi, E., 2000. Subsolidus and melting experiments of K-doped peridotite
 KLB-1 to 27 GPa: Its geophysical and geochemical implications. J. Geophys. Res. 105,
 2855–2868.
- Watson, E.B., Brenan, J.M., Baker, D.R., 1991. Continental Mantle. Oxford University Press.
- Yoshino, T., and Katsura, T. (2012). Re-evaluation of electrical conductivity of anhydrous
 and hydrous wadsleyite. Earth Planet. Sci. Lett. 337, 56–67. doi:
 10.1016/j.epsl.2012.05.023
- Yoshino, T., 2010. Laboratory Electrical Conductivity Measurement of Mantle Minerals.
 Surv. Geophys. 31, 163–206. doi:10.1007/s10712-009-9084-0
- Yoshino, T., Manthilake, G., Matsuzaki, T., Katsura, T., 2008. Dry mantle transition zone
 inferred from the conductivity of wadsleyite and ringwoodite 451, 1–4.
 doi:10.1038/nature06427
- Yoshino, T., Nishihara, Y., Karato, S. ichiro, 2007. Complete wetting of olivine grain
 boundaries by a hydrous melt near the mantle transition zone. Earth Planet. Sci. Lett.
 256, 466–472. doi:10.1016/j.epsl.2007.02.002
- Zhao, D., 2004. Global tomographic images of mantle plumes and subducting slabs : insight
 into deep Earth dynamics 146, 3–34. doi:10.1016/j.pepi.2003.07.032
- 652 653

654 **Figure captions**



Figure 1. Electrical conductivity of hydrous peridotites at 6, 9 and 12 GPa. The 656 peridotites before melting are indicated in squares and after partial melting are in circles. The 657 melt conductivities used for the discussion are shown with horizontal lines. Activation 658 enthalpies are given in eV, next to the individual fits. The error bars associated with the 659 electrical conductivity data measurements are less than the symbol size at high temperatures. 660 The uncertainties in the estimation of the electrical conductivity result from the estimations of 661 temperature, pressure, and sample dimensions as well as data fitting errors and are less than 662 5%. 663



Figure 2. High resolution microphotographs of the recovered samples before and after
the electrical conductivity measurements. (a) Hydrous peridotite sample showing mineral
distribution before melting at 12 GPa (M421). (b) The mineral and melt distribution of
partially molten sample after the electrical conductivity measurements at 6 GPa (M667), (c)
at 9 GPa (M668), (d) and at 12 GPa (M425).





Figure 3. Dihedral angles estimated in our partially molten samples as a function of pressure. The maximum temperature of each experiment is indicated next to its symbol. Dihedral angles were estimated using FEG SEM and SEM image analysis with Matlab code to measure the angles by the tangent method. Analytical error is estimated to be 5° (1 σ). Relevant literature values for similar or analogue systems at lower pressures have been represented for comparison (Mibe et al., 1999, 1998; Watson et al., 1991; Yoshino et al., 2007).



679

Figure 4. Electrical conductivity of hydrous silicate melts compared with geophysical 680 **profiles.** Electrical conductivities of peridotite prior to melting (squares) and partially molten 681 peridotites (circles) at 6, 9 and 12 GPa are displayed as a function of depth. Hashin Shtrikman 682 upper bound (HS+) conductivities for different melt fraction are displayed for 12 GPa 683 samples with dashed lines. The conductivity profiles are for oceanic mantle (pacific, KO5) 684 (Kuvshinov et al., 2005), continental (Tuscan SW USA, T07) (Toffelmier and Tyburczy, 685 2007) and global electrical conductivity model of Earth's mantle based on inverting satellite 686 magnetic field measurements (G17) (Grayver et al., 2017). Pink shaded area indicates the 687 possible distribution of hydrous melt at the top of the mantle transition zone. In this area the 688 high conductivity anomaly modeled from EM data has been represented (Toffelmier and 689 Tyburczy, 2007). Dry upper mantle electrical conductivity vs depth profile based on high 690 pressure electrical conductivity measurements are also displayed (Yoshino et al., 2008). 691 Electrical conductivities of oriented single crystal olivine from measurements (Dai and 692

Karato 2014) (4 GPa and extrapolated to mantle temperatures) and diffusion experiments (Novella et al., 2017b) (2 GPa, extrapolated to mantle temperatures and with 80 ppm of water) have been reported. Electrical conductivity anomaly of 350' cannot be reproduced by moderately anisotropic solid rock but rather require presence of melts. Our solid hydrous peridotites conductivities agree well with randomly oriented olivine values.



Figure 5. Comparison of water transport mechanisms of the Earth's upper mantle. (a) 700 Hydration of the Earth's upper mantle by diffusion. The estimations are based on the 701 diffusion coefficient of 7.89×10^{-9} m²/s (Novella et al., 2017a) at 1723K with initial H₂O 702 concentration of 800 ppm wt. and variable diffusion lengths (from 410-km to the saturation 703 depth at a given water content) (see also Supplementary Fig. S7). The water saturation curves 704 are for peridotite without considering garnet (F12) (Férot and Bolfan-Casanova, 2012) and 705 considering water partition between olivine and garnet (0.9) (A12) (Ardia et al., 2012). The 706 broken lines indicate the time required to saturate the upper mantle by diffusion at given 707 708 depths for a given bulk water content. The shaded in yellow indicates regions where dehydration-melting is expected. The shaded in pink indicate depth range for mantle H₂O 709 saturation which can't be explained by diffusion process. (b) Hydration by upwelling. The 710 dehydration melting at the 410-km discontinuity (Freitas et al., 2017) produces residual 711 peridotites with different H₂O concentration according melt fractions and compositions. The 712 continuous upwelling of hydrous peridotite replaces the initially dry peridotite in the upper 713 mantle, forming hydrous upper mantle. The ascending peridotite reaches water saturation 714 levels at different depths (Ardia et al., 2012; Férot and Bolfan-Casanova, 2012), and should 715 initiate hydrous melting. The broken lines indicate the time required to saturate the upper 716

mantle at given depths for a given bulk water content. The water saturation curves are for peridotite without considering garnet (F12) (Férot and Bolfan-Casanova, 2012) and with considering water partition between olivine and garnet (A12) (Ardia et al., 2012). The shaded in yellow indicates regions where dehydration-melting is expected. The shaded in pink indicate depth range for mantle H₂O saturation.



722

Figure 6. Bottom-up hydration of the upper mantle. The H₂O contents are calculated assuming batch melting scenario at the 410-km discontinuity. The H₂O contents in the source (MTZ) are marked next to individual lines. The Gray shaded area indicates the melt fraction expected at the partially molten layer above the MTZ. The area shaded in Yellow indicates H2O content of the upper mantle. Vertical lines (F12/D16) (Demouchy and Bolfan-Casanova, 2016; Férot and Bolfan-Casanova, 2012) and (A12) (Ardia et al., 2012) indicate water saturation limits at different depths based on water solubilities of peridotites.

Supplementary materials for:

Electrical conductivity of hydrous silicate melts: Implications for the bottom-up hydration of

Earth's upper mantle

D.Freitas^{1*}, G. Manthilake¹

¹Université Clermont Auvergne, CNRS, IRD, OPGC, Laboratoire Magmas et Volcans,

F-63000 Clermont-Ferrand, France

(*corresponding author: damien.freitas@uca.fr)

Supplementary materials

Here we give supplementary information concerning:

- The estimation of the H₂O partition coefficient for peridotite/melt at 12 GPa (Text S1)
- Experimental conditions and parameters of syntheses and *in situ* measurements (table S1)
- Average chemical composition of solid mineral phases at 6 and 9 GPa both before and after EC measurements (Table S2)
- Average chemical composition of solid mineral phases at 12 GPa. (Table S3)
- Schematic cross section of the high pressure assemblies (Figure S1)
- Modeling of thermal gradient in our assembly (Figure S2)
- Impedance spectra of the sample at different stages of heating (Figure S3)
- Representative Raman spectra from starting materials and recovered experimental runs at 6 GPa (Figure S4)
- Representative Raman spectra from starting materials and recovered experimental runs at 9 GPa (Figure S5)
- Representative Raman spectra from recovered experimental runs at 12 GPa. (Figure S6)

• Water diffusion calculations for the hydration of Earth's upper mantle. (Figure S7)

Supplementary Text S1: The estimation of the H₂O partition coefficient for peridotite/melt at 12 GPa.

The H_2O partition coefficient between peridotite and melt is available at 6 GPa (Novella et al., 2014). Due to the significant variation of mineral proportions at 12 GPa, we have estimate the H_2O partition coefficient between peridotite and melt based on mantle mineralogy at 12 GPa. The bulk partition coefficient can be calculated by:

$$D_{H20}^{pdt/melt} = \sum X_i K d_i$$

Where X is the fraction of each mineral (i) and Kd its partition coefficient for water with melt. The Kd values are taken from Novella et al. 2014 (Novella et al., 2014) and mineral proportions where estimated using the mineral proportions estimated at 12 GPa in our study. Mineral proportion at 12 GPa are observed to be 60 % of olivine, 28 % of garnet (Gt) and 12 % of clinopyroxene (Cpx).

	Kd (Novella	STD
	et al., 2014)	510
Melt/olivine	0.004	0.0006
Melt/Opx	0.0064	0.0004
Melt/Cpx	0.0115	0.0016
Melt/garnet	0.0032	0.0008
D _{Melt/pdt} 6 GPa	0.006288	0.000343
D _{Melt/pdt} 12 GPa	0.004676	0.000424

Table S1. Conditions of experimental runs (syntheses and *in situ* measurements). Each run conditions are given in the first section: pressure, temperature and duration at maximum temperature. *In situ* experiments were only kept for short times above melting temperature (<1h) to avoid any melt escape or reaction. Water content are then given within the 3 next columns: initial starting water content in powders, water content in solid samples after syntheses and water content after *in situ* electrical conductivity measurements. Water content on solid samples/samples after *in situ* measurements was estimated qualitatively (or with large errors) by Raman spectroscopy (see methods section and Freitas et al. 2017). Partially molten samples characteristics are given, with melt fractions and estimated melting temperatures based on electrical conductivity. Dihedral angles measurements are given with average and median angles for each partially molten sample. In the last column (remarks), the type of each experiment is given; synthesis run number used for electrical conductivity measurements is given in parenthesis. Errors given in parenthesis are 1 standard deviation.

¹ Water content of the bulk sample estimated with quenched glass quantification and partition coefficient (Novella et al. 2014) with mineral proportions inferred from BSE image analyses.

² Melt fractions estimated by mass balance, partition coefficients and image surface analyses.

³ Melt fractions estimated by image surface analysis only.

	Experimer	ntal conditions		Water contents (ppm)			Partial melting		Dihedral angle Θ			
Run #	Pressure (GPa)	Tmax (K)	Time	initial powder	Solid sample	After <i>in situ</i> EC ¹	%melt ²	Tmelting (K) ⁴	Number of measurements	Mean	Median	Remarks
M405	5	1523 (20)	3h45	0	-	-	-	-	-	-	-	Synthesis
M407	5	1473 (20)	1h	0	-	-	-	-	-	-	-	Synthesis
M408	14	1373 (20)	7h	≈500	>3000	-	-	-	-	-	-	Synthesis
M410	14	1373 (20)	2h10	≈500	3267 (650)	-	-	-	-	-	-	Synthesis
M412	12	1738 (5)	<1h	0	-	3098 (959)	7.00 (0.27)	1143 (5)	77.00	12.20 (11.68)	10.51	EC (M405)
M415	5	1450 (20)	2h30	0	-	-	-	-	-	-	-	Synthesis
M421	12	1385 (5)	<1h	0	-	993 (260)	-	-	-	-	-	EC (M415)
M425	12	1738. (5)	<1h	≈500	≈3000	3030 (627)	2.00 (0.45)	1493 (5)	139.00	7.88 (8.79)	6.23	EC (M408)
M455	14	1373 (20)	2h	≈700	≈3600	-	-	-	-	-	-	Synthesis
M456	12	1503 (5)	<1h	≈700	≈3600	-	3.50 (2.44)	1303 (5)	59.00	10.12 (10.26)	8.50	EC (M455)
M481	2	1473 (20)	2h	0	-	-	-	-	-	-	-	Synthesis
M482	14	1323 (20)	2h10	-	500-700	-	-	-	-	-	-	Synthesis (M481)
M490	12	1780 (5)	<1h	0	500-700	800-1650	25.65 (2.50)	1561 (5)	36.00	16.60 (9.52)	17.09	Freitas et al. 2017 (M481)
M492	12	1363 (5)	<1h	0	500-700	490 (125)	-	-	-	-	-	Freitas et al. 2017 (M482)
M665	6	1273 (20)	4h15	≈700	1579 (413)	-	-	-	-	-	-	Synthesis
M666	9	1323 (20)	3h30	≈700	1740 (523)	-	-	-	-	-	-	Synthesis
M667	6	1378 (5)	<1h	≈700	1579 (413)	-	3.68 (1.86) ³	1163 (5)	104.00	28.46 (13.19)	27.26	EC (M665)
M668	9	1480.20 (5)	<1h	≈700	1105 (422)	-	6.33 (1.01) ³	1181 (5)	109.00	20.32 (9.53)	19.49	EC (M666)

Table S2. Average chemical composition of solid mineral phases at 6 and 9 GPa both before and after EC measurements. Each value is an average of 3 to 10 analyses; the standard deviation is indicated in parenthesis. Chemical analyses of quenched melt were not possible due to the extremely small size of the melt pockets.

	Starti	ng material at	6 GPa	After EC exp. at 6 GPa			Starting material at 6 GPa			After EC exp. at 6 GPa		
	Ol	Срх	Gt	Ol	Срх	Gt	Ol	Срх	Gt	Ol	Срх	Gt
SiO ₂	40.00 (0.90)	54.82 (0.22)	46.13 (0.92)	40.65 (0.87)	54.42 (0.73)	42.13 (2.84)	39.30 (1.00)	53.58 (0.63)	39.81 (2.45)	39.28 (1.39)	51.14 (0.96)	40.68 (0.22)
TiO ₂	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Al ₂ O ₃	0.29 (0.37)	2.34 (0.64)	14.66 (0.49)	0.44 (0.43)	2.59 (0.36)	15.41 (6.69)	1.03 (0.03)	1.48 (0.15)	18.50 (4.68)	0.20 (0.20)	1.79 (0.22)	20.53 (0.54)
Cr_2O_3	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
FeO	10.71 (1.86)	3.93 (1.35)	8.80 (0.77)	10.79 (2.54)	2.77 (0.09)	9.01 (1.00)	8.81 (0.22)	4.22 (3.06)	10.84 (1.85)	9.18 (1.34)	3.06 (0.30)	9.17 (0.12)
MnO	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
MgO	40.00 (0.90)	19.12 (0.46)	22.75 (0.74)	47.17 (2.57)	19.70 (0.68)	42.13 (2.84)	50.00 (1.25)	19.30 (0.94)	23.29 (5.35)	48.25 (2.45)	20.91 (2.57)	20.47 (2.93)
CaO	0.09 (0.05)	18.38 (2.84)	6.79 (1.53)	0.16 (0.11)	18.97 (0.60)	5.04 (2.87)	0.17 (0.01)	18.88 (4.42)	4.81 (1.22)	0.14 (0.10)	18.65 (1.82)	5.87 (0.71)
Na ₂ O	0.22 (0.28)	1.91 (0.68)	0.72 (0.11)	0.31 (0.37)	1.79 (0.27)	0.60 (0.19)	0.58 (0.02)	1.78 (0.90)	0.49 (0.28)	0.27 (0.24)	1.42 (0.19)	0.17 (0.21)
NiO	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
K ₂ O	-	-	-	-	0.01 (0.02)	0.01 (0.02)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	-	0.01 (0.01)	0.05 (0.02)
Total	99.91 (1.15)	100.52 (0.57)	99.84 (0.86)	99.55 (0.94)	100.19 (0.33)	99.89 (1.36)	99.9 (2.50)	99.24 (0.47)	97.74 (1.11)	97.36 (0.28)	96.97 (0.58)	96.95 (1.53)

Ol: olivine, Cpx: clino-pyroxenes, Gt: garnet

Table S3. Average chemical composition of solid mineral phases at 12 GPa. Each value is an average of 5 to 20 analyses.	The standard
deviation is given on the average value in parenthesis.	

	М	412-7 (0.27)% M		M421-0% Melt			M425-2 (0.45)% Melt			M456-3.5 (2.44)% Melt		
	Ol	Срх	Gt	Ol	Срх	Gt	Ol	Срх	Gt	Ol	Срх	Gt
SiO ₂	41.60 (0.38)	56.48 (1.01)	47.23 (0.89)	40.90 (0.87)	54.62 (0.65)	40.69 (1.38)	41.36 (0.26)	54.78 (0.67)	43.95 (1.46)	41.99 (0.48)	55.73 (0.36)	46.51 (4.01)
TiO ₂	0.00 (0.00)	0.03 (0.01)	0.78 (0.38)	0.04 (0.03)	0.10 (0.04)	0.57 (0.39)	0.04 (0.03)	0.06 (0.03)	0.43 (0.16)	0.05 (0.04)	0.10 (0.05)	0.32 (0.26)
Al ₂ O ₃	0.12 (0.02)	2.66 (0.29)	8.87 (2.21)	0.04 (0.08)	1.66 (0.46)	22.19 (3.04)	0.44 (0.60)	1.74 (0.57)	17.13 (5.31)	0.15 (0.04)	1.83 (0.40)	13.17 (10.42)
Cr ₂ O ₃	0.12 (0.02)	0.01 (0.02)	0.01 (0.01)	0.01 (0.02)	0.01 (0.02)	0.03 (0.02)	0.01 (0.01)	0.02 (0.01)	0.03 (0.02)	0.00 (0.00)	0.00 (0.02)	0.00 (0.00)
FeO	7.15 (0.24)	4.83 (0.48)	9.77 (1.77)	8.99 (1.12)	2.97 (0.71)	7.73 (0.58)	7.01 (0.78)	2.90 (0.35)	6.34 (0.61)	5.55 (1.96)	2.96 (0.15)	5.17 (0.99)
MnO	0.02 (0.02)	0.02 (0.02)	0.05 (0.02)	0.02 (0.02)	0.01 (0.01)	0.06 (0.02)	0.02 (0.02)	0.02 (0.02)	0.06 (0.03)	0.01 (0.01)	0.02 (0.01)	0.05 (4.59)
MgO	50.78 (0.58)	33.61 (0.73)	13.64 (8.72)	48.80 (0.68)	18.74 (5.83)	18.51 (4.17)	50.10 (1.20)	19.09 (1.18)	24.97 (7.53)	52.67 (2.02)	17.83 (0.62)	24.57 (4.59)
CaO	0.18 (0.03)	1.70 (0.23)	15.99 (4.72)	0.11 (0.12)	20.04 (4.33)	8.77 (7.09)	0.36 (0.30)	19.79 (1.07)	6.94 (1.97)	0.23 (0.14)	20.96 (0.50)	10.58 (2.07)
Na ₂ O	0.02 (0.02)	0.21 (0.13)	1.46 (0.42)	0.02 (0.04)	1.31 (0.29)	0.06 (0.02)	0.03 (0.02)	0.97 (0.18)	0.11 (0.07)	0.03 (0.02)	1.23 (0.12)	0.31 (0.25)
NiO	0.01 (0.01)	0.07 (0.02)	0.00 (0.00)	-	-	-	-	-	-	0.01 (0.01)	0.00 (0.00)	0.02 (0.01)
K ₂ O	0.01 (0.01)	0.01 (0.02)	0.06 (0.06)	0.01 (0.01)	0.03 (0.02)	0.01 (0.02)	0.00 (0.00)	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.02 (0.01)	0.02 (0.01)
Total	99.88 (0.93)	100.16 (1.14)	97.86 (5.21)	98.94 (0.67)	99.50 (1.40)	98.62 (1.05)	99.38 (0.45)	99.39 (0.57)	99.95 (1.70)	100.70 (0.29)	100.68 (0.61)	100.71 (1.03)

Ol:olivine, Cpx: clino-pyroxenes, Gt: garnet



Figure S1. Schematic cross section of the high pressure assemblies. Assembly geometries on the left were used for the synthesis of the peridotite samples and on the right were used for electrical conductivity measurements at 6, 9 and 12 GPa. Top assemblies are 14/8 geometries whereas bottom ones are 18/11.



Figure S2. Modeling of thermal gradient in our assembly. Results of simulations using the software codes of Hernlund et al. 2006 (Hernlund et al., 2006) . The maximum temperature gradient of the assembly at 1500 K was estimated to be less than 20 K.



Figure S3. Impedance spectra of the sample at different stages of heating. (a) during assembly dehydration at 673 K (b) sample annealing at 1173 K. (c) Impedance spectra recorded after assembly dehydration and sample annealing steps. Moisture tails were not observed anymore during this cycle. (d) Impedance spectra recorded after the onset of melting, these spectra are characterized by the disappearance of the semicircular shape for induction patterns.



Figure S4. Representative Raman spectra from starting materials and recovered experimental runs at 6 GPa. Low-wavenumber (200-1200 cm⁻¹) bands corresponding to vibrations of the silicate network (a, c, e) and high-wavenumber bands (3100-3800 cm⁻¹) corresponding to O-H vibrations (b, d, f) of olivine, pyroxene and garnet, respectively. The presence of H₂O is visible in all samples. H₂O content is significantly reduced in the experiments that undergoes partial melting. Black and gray spectra were taken in starting material after the synthesis run, green spectra were acquired on samples recovered after the *in situ* measurements.



Figure S5. Representative Raman spectra from starting materials and recovered experimental runs at 9 GPa. Low-wavenumber (200-1200 cm⁻¹) bands corresponding to vibrations of the silicate network (a, c, e) and high-wavenumber bands (3100-3800 cm⁻¹) corresponding to O-H vibrations (b, d, f) of olivine, pyroxene and garnet, respectively. The presence of H₂O is visible in all samples. H₂O content is significantly reduced in the experiments that undergoes partial melting. Black and gray spectra were taken in starting material after the synthesis run, green spectra were acquired on samples recovered after the *in situ* measurements.



Figure S6. Representative Raman spectra from recovered experimental runs at 12 GPa. Red spectra were measured on samples with 600 ppm wt. bulk H₂O contents. Black spectra were acquired on the samples that undergone partial melting. Spectra on the left column (a, c, e) were made on silicate frequency region (200-1200 cm⁻¹) whereas spectra displayed on the right column (b, d, f) were performed on O-H vibration frequency region (3100-3800 cm⁻¹) of olivine, pyroxene and garnet, respectively.



Figure S7. Water diffusion calculations for the hydration of Earth's upper mantle. (a). Water concentration profiles as a function of time at depth of 160 km (250 km of diffusion length). The diffusion profile is calculated for an initially dry upper mantle ($C_0 = 0$ ppm) and considering an olivine peridotite in contact with hydrous melt at 410-km discontinuity with a constant, sustained water concentration C1 = 800 ppm wt. The calculations are based on the diffusion coefficient of 7.82×10^{-9} m²/s (Novella et al., 2017) at 1723K (b) Water concentration profiles in the upper mantle for different times of diffusion (0.01 to 4 Ga) for an initially dry upper mantle and 800 ppm wt. olivine peridotite in equilibrium with hydrous melt at the top of the MTZ. The water saturation curves are for peridotite without considering garnet (F12+D16) (Demouchy and Bolfan-Casanova, 2016; Férot and Bolfan-Casanova, 2012) and with considering water partition between olivine and garnet (0.9) (A12) (Ardia et al., 2012). The diffusion profiles even for Earth's time scale can't reach the saturation curve. Diffusion is able to keep large gradients of water concentration in Earth's upper mantle over geological times.

References:

- Ardia, P., Hirschmann, M.M., Withers, A.C., Tenner, T.J., 2012. H2O storage capacity of olivine at 5-8GPa and consequences for dehydration partial melting of the upper mantle.
 Earth Planet. Sci. Lett. 345–348, 104–116. doi:10.1016/j.epsl.2012.05.038
- Demouchy, S., Bolfan-Casanova, N., 2016. Distribution and transport of hydrogen in the lithospheric mantle: A review. Lithos 240–243, 402–425. doi:10.1016/j.lithos.2015.11.012
- Férot, A., Bolfan-Casanova, N., 2012. Water storage capacity in olivine and pyroxene to 14GPa: Implications for the water content of the Earth's upper mantle and nature of seismic discontinuities. Earth Planet. Sci. Lett. 349–350, 218–230. doi:10.1016/j.epsl.2012.06.022
- Freitas, D., Manthilake, G., Schiavi, F., Chantel, J., Bolfan-Casanova, N., Bouhifd, M.A., Andrault, D., 2017. Experimental evidence supporting a global melt layer at the base of the Earth's upper mantle. Nat. Commun. 8, 2186. doi:10.1038/s41467-017-02275-9
- Hernlund, J., Einenweber, K., Locke, D., Tyburczy, J., 2006. A numerical model for steadystate temperature distributions in solid-medium high-pressure cell assemblies 91, 295– 305. doi:10.2138/am.2006.1938
- Novella, D., Frost, D.J., Hauri, E.H., Bureau, H., Raepsaet, C., Roberge, M., 2014. The distribution of H2O between silicate melt and nominally anhydrous peridotite and the onset of hydrous melting in the deep upper mantle. Earth Planet. Sci. Lett. 400, 1–13. doi:10.1016/j.epsl.2014.05.006
- Novella, D., Jacobsen, B., Weber, P.K., Tyburczy, J.A., Ryerson, F.J., Du Frane, W.L., 2017. Hydrogen self-diffusion in single crystal olivine and electrical conductivity of the Earth's mantle. Sci. Rep. 7, 1–10. doi:10.1038/s41598-017-05113-6