



HAL
open science

Lu-Hf systematics of 4.0 – 2.3 Ga old zircons from the Turee Creek Group (Pilbara Craton, W. Australia): Implications on the rise of atmospheric oxygen and global glaciation during the Paleoproterozoic

Tom Caquineau, Jean-Louis Paquette, Abdelmouhcine Gannoun, Pascal Philippot

► To cite this version:

Tom Caquineau, Jean-Louis Paquette, Abdelmouhcine Gannoun, Pascal Philippot. Lu-Hf systematics of 4.0 – 2.3 Ga old zircons from the Turee Creek Group (Pilbara Craton, W. Australia): Implications on the rise of atmospheric oxygen and global glaciation during the Paleoproterozoic. *Precambrian Research*, 2020, 348, pp.105859. 10.1016/j.precamres.2020.105859 . hal-02906321

HAL Id: hal-02906321

<https://uca.hal.science/hal-02906321>

Submitted on 24 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Lu-Hf systematics of 4.0 – 2.3 Ga old zircons from the Turee Creek**
2 **Group (Pilbara Craton, W. Australia): Implications on the rise of**
3 **atmospheric oxygen and global glaciation during the Paleoproterozoic**

4
5 Tom Caquineau^a, Jean-Louis Paquette^{b*}, Abdelmouhcine Gannoun^b, Pascal Philippot^{c,d,a}

6
7 ^a Institut de Physique du Globe de Paris, Sorbonne-Paris Cité, Université Paris Diderot,
8 CNRS, 1 rue Jussieu 75238 Paris Cedex 05, France.

9 ^b Université Clermont Auvergne, CNRS, IRD, OPGC, Laboratoire Magmas et Volcans, F-
10 63000 Clermont-Ferrand, France

11 ^c Géosciences Montpellier, CNRS-UMR 5243, Université de Montpellier, Montpellier
12 Cedex5, France

13 ^d IAG, Universidade de São Paulo, Rua do Matão, 1226 – Cidade Universitária, Butantã,
14 05508-090 São Paulo – SP, Brazil

15

16 *Corresponding author e-mail: J.L.paquette@opgc.univ-bpclermont.fr

17

18 **Abstract**

19 We investigated the Hf isotopic systematics of detrital zircons in a succession of
20 siliciclastic sediments and glacial diamictites from the early Paleoproterozoic sequence of the
21 Boolgeeda Iron Formation (Hamersley Group) and overlying Turee Creek Group of the
22 Pilbara Craton, Western Australia. About 400 detrital zircons yielding > 95% concordant U-
23 Pb ages were analyzed for Hf isotopes to constrain their magmatic sources. 70% of the
24 analyzed zircons display super-chondritic initial Hf isotopic compositions, demonstrating
25 crystallization in mantle-derived magmas. Most of the data are comprised between model age

26 lines at ~2.5 and 3.2 Ga, which suggests a sub-continuous crust generation by extraction from
27 the depleted mantle during this time period. A single grain yields a 4.0 Ga age, which
28 represents the first Hadean age for a zircon from the Pilbara Craton. Our results are
29 significantly different from zircon Hf isotope data of the Glenburgh Terrane, in the southern
30 border of the Turee Creek Group, or older successions of the Pilbara, Kaapvaal and Superior
31 cratons, but show overlap with some of the Yilgarn Craton. This together with the occurrence
32 of a Hadean zircon crystal preserved in the Boolgeeda glacial diamictite with similar Hf
33 isotopic signature than the Jack Hills zircons makes the Yilgarn Craton a possible source
34 material for the Boolgeeda glacial horizon. Alternatively, the majority of the zircons analyzed
35 show ages which are consistent with those of the underlying 2.45 – 2.78 Ga Hamersley and
36 Fortescue groups, formed by sedimentary successions interleaved with thick subaerial
37 volcanic sequences associated with the emplacement of Large Igneous Provinces. Such
38 subaerial volcanic rocks could account for the relatively juvenile character of the zircon
39 analyzed. A local provenance of the siliciclastic material delivered to the Turee Creek Basin
40 would support the role of large subaerial magmatic provinces as triggers of the rise of
41 atmospheric oxygen and the onset of glaciations at the beginning of the Proterozoic.

42

43 *Keywords*

44 Zircon; Hafnium isotopes; Turee Creek Group; Great Oxidation Event; Paleoproterozoic
45 glaciations; Hadean.

46

47 **1. Introduction**

48 The Archean-Proterozoic transition was marked by important redox and climatic
49 changes, specifically the rise of atmospheric oxygen, the so-called Great Oxidation Event
50 (GOE) (Holland, 2002), and several Paleoproterozoic glacial events (e.g., Rasmussen *et al.*,

51 2013; Caquineau *et al.*, 2018). The causal relationships between these events remain largely
52 debated (e.g., Kopp *et al.*, 2005; Teitler *et al.*, 2014). One model proposed by Kump and
53 Barley (2007) explained the rise of atmospheric oxygen by a change in the eruptive dynamic
54 of the Earth at ca. 2.5 Ga that shifted the main style of volcanism from marine to subaerial.
55 This increase in the subaerial proportion of Large Igneous Provinces (LIPs) would have
56 buffered the greenhouse gases content of the atmosphere due to carbon dioxide consumption
57 through enhanced silicate weathering, thus enabling atmospheric oxygen to build up and
58 surface temperature to cool down, leading to global glaciation (Cox *et al.*, 2016). In addition,
59 subaerial eruptions are known to produce more oxidized lavas and gases (Holland *et al.*, 2002;
60 Gaillard *et al.*, 2011), which could also have contributed further to reduce oxygen sinks.

61 Zircon is a very resistant mineral, of which Hf is a major constituent (~ 1wt%), that
62 represents the only witness of eroded and unknown source protoliths (Iizuka *et al.*, 2017). The
63 incorporation of significant amounts of mantle-derived volcanic material within the upper
64 continental crust can generate a large increase of juvenile Hf isotopic signatures in the zircons
65 from the sediments. Glacial diamictites are considered to record the average chemical
66 composition of the upper continental crust from which they formed by erosion (Gaschnig *et*
67 *al.*, 2014). Consequently, U-Pb-Hf systematics in zircons sampled in these sediments appears
68 as a valuable strategy to test the possible occurrence of materials derived from mantle sources.
69 Here, we performed a Lu-Hf isotopic study of detrital zircons previously analyzed for U-Pb
70 geochronology (Caquineau *et al.*, 2018) from a suite of glaciogenic and non-glaciogenic
71 sediments of Paleoproterozoic age that spans the GOE. The rocks investigated are from the
72 Boolgeeda Iron Formation, which represents the uppermost succession of the Hamersley
73 Group, and the Turee Creek Group (TCG) in the Hamersley Basin at the southern margin of
74 the Pilbara Craton, Western Australia (Fig. 1). The main objective of this study is to (i)
75 identify the major sources and further constrain the formation's model of this sedimentary

76 sequence and (ii) to test the hypothesis of massive subaerial LIPs emplacement in the Pilbara
77 Craton and its implications on the GOE and the Paleoproterozoic glaciations.

78

79 **2. Geological setting**

80 The Turee Creek Group (TCG) is a ca. 4 km-thick continuous sedimentary succession
81 conformably overlying the Boolgeeda Iron Formation of the Hamersley Group and
82 unconformably overlain by the Beasley River Quartzite and the Cheela Springs Basalt of the
83 lower Wyloo Group (Figs. 1, 2; Martin, 1999; Van Kranendonk *et al.*, 2015). From bottom to
84 top, the TCG consists of siliciclastic sedimentary rocks, minor stromatolitic carbonates and
85 glaciogenic strata of the Kungarra Formation, quartz-rich sandstones of the Koolbye
86 Formation, and shales and stromatolitic marine carbonates of the Kazput Formation (Fig. 2b;
87 Trendall, 1981; Martin, 1999; Van Kranendonk *et al.*, 2015). The TCG sedimentary sequence
88 has been described as a broad shallowing upward profile from deep-water iron formation of
89 the Boolgeeda Iron Formation, through fine-grained siliciclastic deposits of the Kungarra
90 Formation, to fluvial and shallow marine strata of the Koolbye and Kazput formations
91 (Martin, 1999; Mazumder *et al.*, 2015; Van Kranendonk *et al.*, 2015). The Boolgeeda Iron
92 Formation overlies the Woongarra Rhyolite and Weeli Wooli Formation, which have been
93 interpreted as a subaerial bimodal mafic and felsic volcanic sequence emplaced at 2450 Ma
94 (Barley *et al.*, 1997). This intense volcanic event would have produced large volumes of lavas
95 ($>30\,000\text{ km}^3$, Barley *et al.*, 1997) covering the continental surface at the time of the glacial
96 event recorded in the Boolgeeda Iron Formation, and possibly during the subsequent
97 deposition of the overlying TCG. The unconformably overlying Beasley River Quartzite
98 represents a terrestrial deposit composed of a basal fluvial conglomerate and overlain by an
99 unit of fine-grained quartzite (Mazumder and Van Kranendonk, 2013). Metamorphism of the
100 group did not exceed sub-greenschist facies conditions (Rasmussen *et al.*, 2005).

101 Several sections of the TCG have been described in the southwestern margin of the
102 Hamersley Basin (Martin, 1999; Van Kranendonk *et al.*, 2015; Krapež *et al.*, 2017). These
103 include from East to West: the Turee Creek, Hardey and Duck Creek synclines, as well as the
104 Yeera Bluff area (Fig. 1b). Glaciogenic horizons were used as reference markers to establish
105 stratigraphic correlations among these different sections. The main glacial body is the ~400m
106 thick Meteorite Bore Member (MBM) that was originally described in the Hardey Syncline
107 section (Trendall, 1981; Figs. 1, 2b and e). More recently, Van Kranendonk and Mazumder
108 (2015) reported a second level of massive diamictites located above the reference Meteorite
109 Bore Member in the Hardey Syncline; in the following, we will refer to this horizon as the
110 ‘second Kungarra diamictite’. A third diamictite horizon associated with sandstones and
111 siltstones was described in the Duck Creek Syncline at the Boundary Ridge locality and at
112 Yeera Bluff located 50 and 150 km north westward from the Hardey Syncline, respectively
113 (Fig. 1b, Martin, 1999; Van Kranendonk *et al.*, 2015). These diamictites, associated with deep
114 marine sediment of the Boolgeeda Iron Formation, were interpreted as distal equivalents of
115 the Meteorite Bore Member formed as sediment gravity flows and rain-out from rare icebergs
116 (Martin, 1999; Van Kranendonk *et al.*, 2015). However, Caquineau *et al.* (2018) and Philippot
117 *et al.* (2018) recently reported the occurrence of a 2m-thick diamictite horizon within the
118 underlying Boolgeeda Iron Formation in the Hardey Syncline (drill core TCDP1, see below
119 and Fig. 2d). Because this diamictite layer is stratigraphically located ~1500 meters below the
120 Meteorite Bore Member, it thus represents a different and older glacial event. Based on new
121 geochronological (see below) and stratigraphic constraints, Caquineau *et al.* (2018) and
122 Philippot *et al.* (2018) showed that this diamictite layer can be correlated with the diamictites
123 horizons reported in the Boolgeeda Iron Formation at Yeera Bluff and the Boundary Ridge
124 locality. This glacial horizon is named hereafter the ‘Boolgeeda diamictite’.

125 Time constraints on the deposition of the sedimentary sequence of interest include: (i)
126 a U-Pb zircon age of 2450 ± 3 Ma for the underlying Woongarra Rhyolite (Figs. 1, 2a and b;
127 Trendall *et al.*, 2004), (ii) a U-Pb detrital zircon maximum age of 2460 ± 9 Ma for the
128 Boolgeeda diamictite (Caquineau *et al.*, 2018), (iii) a Re-Os age of 2312 ± 6 Ma obtained on
129 diagenetic pyrite from the Meteorite Bore Member diamictite (Philipipot *et al.*, 2018) in
130 agreement with a U-Pb maximum age on detrital zircon of 2340 ± 22 Ma (Caquineau *et al.*,
131 2018) for this glacial formation, and (iv) a U-Pb zircon age of 2209 ± 15 Ma for the overlying
132 Cheela Springs Basalt (Martin *et al.*, 1998). A baddeleyite U-Pb age of 2208 ± 10 Ma was
133 obtained on dolerite sills that intrude the TCG and are truncated by the unconformably
134 overlying Beasley River Quartzite (Müller *et al.*, 2005). Thus, the significance of the age of
135 the Cheela Springs Basalt has been questioned (Müller *et al.*, 2005). Accordingly, the age of
136 the Beasley River Quartzite remains loosely bracketed between ~ 2210 Ma and 2031 ± 6 Ma
137 (Fig. 2a, Martin *et al.*, 1998; Müller *et al.*, 2005).

138 The sedimentation of the TCG is considered to have occurred in the McGrath Trough,
139 an asymmetric basin structurally linked with the migration to the northeast of a geanticline or
140 a thrust-fold belt system (e.g., Martin *et al.*, 2000; Van Kranendonk *et al.*, 2015). This basin
141 has been interpreted either as a retroarc foreland basin (Martin *et al.*, 2000; Müller *et al.*,
142 2005; Krapež *et al.*, 2017) or as an intracontinental basin (Van Kranendonk *et al.*, 2015). It is
143 generally accepted that the Turee Creek Group formed in a long-lived basin between ~ 2.45
144 and 2.2 Ga (Martin *et al.*, 2000; Van Kranendonk *et al.*, 2015). An alternative view (Krapež *et*
145 *al.*, 2017), which proposes that the TCG formed in a short-lived foreland basin that closed by
146 ~ 2.43 Ga or shortly after, can be ruled out by recent geochronological constraints on the
147 MBM diamictites mentioned above (Caquineau *et al.*, 2018; Philipipot *et al.*, 2018). The
148 provenance of the sediments forming the TCG is subject of debate, mainly because it relies on
149 controversial paleocurrent data, which established that the detritus was principally derived

150 from the south and the southwest (Martin *et al.*, 2000; Van Kranendonk *et al.*, 2015; Krapež
151 *et al.*, 2017). The major source terranes include the Pilbara Craton, the Hamersley Province to
152 the north (Martin *et al.*, 2000; Van Kranendonk *et al.*, 2015), and the Glenburgh Terrane to
153 the south (Krapež *et al.*, 2017), a continental exotic fragment that is thought to have collided
154 with the southwestern margin of the Pilbara Craton ca. 2.2 Ga ago (Johnson *et al.*, 2011;
155 2013). The occurrence of three discrete glacial diamictite horizons, and new age constraints of
156 the TCG led Caquineau *et al.* (2018) to propose that the ca. 2.45 Ga-old Boolgeeda diamictite
157 can be correlated with (1) the pre-2.43 Ga Makganyene Formation of the South African
158 Transvaal Supergroup (Gumsley *et al.*, 2017), (2) the basal glaciogenic horizon of the Ramsay
159 Lake Formation of the North American Huronian Supergroup (Rasmussen *et al.*, 2013), and
160 the ca. 2.43 Ga old Fennoscandian glacial diamictites (Melezhik *et al.*, 2013). This early
161 glacial event at about 2.45 Ga recognized in five different cratons could therefore have been
162 global in extent (i.e., Snowball Earth). Caquineau *et al.* (2018) further proposed that the two
163 diamictites of the Kungarra Formation (the MBM and the second Kungarra diamictite) can be
164 correlated with the pre-2.31 Ga Gowganda Formation (North America) and upper Deutschland
165 (South Africa) glacial diamictites (Rasmussen *et al.*, 2013), and the ca. 2.26 Ga old upper
166 Timeball Hill (North America) and Ditojana (South Africa) diamictites (Rasmussen *et al.*,
167 2013), as well as the Enchantment Lake Formation (North America).

168

169 **3. Samples**

170 The samples analyzed in this study have been described in detail in Caquineau *et al.*
171 (2018). These include drill core samples collected in the course of the Turee Creek Drilling
172 Project (TCDP1 and TCDP2) obtained at the base and intermediate level of the Turee Creek
173 Group, respectively, as well as surface samples collected in lithostratigraphic levels not
174 intercepted by the drill cores (Fig. 2). Six samples were selected from different stratigraphic

175 levels in drill cores TCDP 1 and TCDP 2 and eight complementary samples were collected in
176 the field (Fig. 2; drill core samples are noted T1-X or T2-X, with X referring to the depth;
177 surface samples are labelled with Pi- followed by the sampling year and the sample number).
178 These include from bottom to top: Three glaciogenic dropstone-bearing sandstones of the
179 Boolgeeda glacial diamictite (T1-152 and Pi-09.07 and Pi-09.08 of the Boundary Ridge
180 Locality), a sandstone layer at the base of TCDP 2 (T2-332) and four samples of the main
181 glacial diamictite deposit of the Meteorite Bore Member (T2-272, T2-205, T2-169, T2-130),
182 three samples of the second Kungarra glacial diamictites (Pi-15.15, Pi-15.16, Pi-15.17), two
183 samples from meter-scale massive sandstone bars of the upper part of the Kazput Formation
184 (Pi 15.41, Pi-15.42), and one sample of the Beasley River Quartzite (Pi-15.02).

185 The Hf isotopic composition was measured in 382 detrital zircon grains from
186 diamictites, sandstones and quartzites that previously gave > 95% concordant U-Pb dates
187 (Fig. 2 and Table 2). These include: (1) 82 zircons from the Boolgeeda diamictite both from
188 the Hardey (drill core sample T1-152) and Duck Creek (surface samples Pi 09.07 and Pi
189 09.08) synclines (Fig. 2c and d), (2) 108 zircons from the Meteorite Bore Member diamictite
190 (drillcore samples T2-130, T2-169, T2-205 and T2-272) and an underlying sandstone bar (T2-
191 332) (Fig. 2e), (3) 101 zircons from the second Kungarra diamictite (surface samples Pi
192 15.15, Pi15.16, Pi 15.17), (4) 59 zircons from two sandstone layers of the upper part of the
193 Kazput formation (surface samples Pi 15.41, Pi 15.42) (Fig. 2f), and (5) 32 zircons from the
194 Beasley River Quartzite (surface sample Pi 15.02).

195

196 **4. Methods**

197 The cathodoluminescence image (CL) of the Hadean mounted zircon grain was
198 generated using a Zeiss Sigma FEG-SEM equipped with a detector operating at 15kV current.
199 Mineral separation involved crushing and sieving at the threshold of 300 μm , Wilfley table,

200 Frantz magnetic separator and heavy liquids. U-Th-Pb isotopic data on zircons were obtained
201 by laser ablation inductively coupled plasma mass spectrometry (LA-ICPMS) at the
202 Laboratoire Magmas & Volcans (LMV, Clermont-Ferrand, France). Detailed analytical
203 techniques are reported in Hurai *et al.* (2010), Paquette *et al.* (2014) and Caquineau *et al.*
204 (2018).

205 Hafnium isotope analyses were performed with a Thermo Scientific Neptune Plus
206 Multicollector (MC) ICPMS coupled to a Resonetics M50E 193nm Excimer laser system at
207 the LMV (Clermont-Ferrand). The operating conditions and instrument settings are described
208 in Moyen *et al.* (2017) and Paquette *et al.* (2017) and detailed in S1. Repeated analyses of
209 both natural (91500, GJ-1 and Mud-Tank) and synthetic Hf and REE-doped (Fischer *et al.*,
210 2011) zircon reference materials are reported in Table S1. More than 50 replicates of 91500,
211 GJ-1 and Mud Tank zircon reference materials yielded $^{176}\text{Hf}/^{177}\text{Hf}$ of 0.282307 ± 16 ,
212 0.282011 ± 15 and 0.282520 ± 12 , respectively, which well agree with reference values (eg.
213 Hu *et al.*, 2012). Mean and two standard deviation values obtained on MUNZirc 0-2a
214 ($^{176}\text{Hf}/^{177}\text{Hf} = 0.28213 \pm 3$), MUNZirc 1-2b (0.28214 ± 2); MUNZirc 3-2b (0.28215 ± 2) and
215 MUNZirc 4-2b (0.28216 ± 3) are fully consistent with reference values (Fisher *et al.*, 2011).
216 The ϵ_{Hf} values are calculated as parts per 10,000 deviation of the $^{176}\text{Hf}/^{177}\text{Hf}$ of the zircons at
217 the time of their formation relative to the Chondritic Uniform Reservoir (CHUR) defined by
218 the ^{176}Lu decay constant of $1.867 \times 10^{-11}\text{yr}^{-1}$ (Söderlund *et al.*, 2004) and the CHUR
219 parameters ($^{176}\text{Hf}/^{177}\text{Hf} = 0.282793$; $^{176}\text{Lu}/^{177}\text{Hf} = 0.0338$; Iizuka *et al.*, 2015). A $^{176}\text{Lu}/^{177}\text{Hf}$
220 ratio of 0.015 typical of bulk continental crust (Griffin *et al.*, 2004) has been used for
221 representing model age lines.

222

223 **5. Zircon Hf isotopic compositions**

224 Hf isotopic data are presented in Table 1. The zircons from the Boolgeeda diamictite
225 yielded a range of $^{207}\text{Pb}/^{206}\text{Pb}$ ages between 2436 and 3216 Ma, as well as a single Hadean
226 grain (discussed below), and initial $\epsilon_{\text{Hf}}(t)$ values between -12.4 to +6.2 (Fig. 3a). About half
227 of the analyzed zircons display a supra-chondritic Hf isotopic composition (i.e. > 0) reflecting
228 the addition of material derived from partial melting of the depleted mantle at an unknown
229 time. In contrast, the remaining zircons analyzed shows a sub-chondritic Hf initial isotopic
230 composition (i.e. < 0) indicating that zircons parental melts recycled an evolved crust.

231 Zircons from the MBM diamictite yielded a range of $^{207}\text{Pb}/^{206}\text{Pb}$ ages between 2359
232 and 3000 Ma and initial $\epsilon_{\text{Hf}}(t)$ values between -5.2 and +7.6 (Fig. 3b). The mantle-derived
233 component is proportionally more important in the MBM diamictite than in the Boolgeeda
234 diamictite and the time interval appears shifted toward slightly younger ages. Zircons from
235 the second Kungarra diamictite yielded a range of $^{207}\text{Pb}/^{206}\text{Pb}$ ages between 2430 and 3350
236 Ma and initial $\epsilon_{\text{Hf}}(t)$ values between -11.8 and +6.7 (Fig. 3 c). Again the addition of a mantle-
237 derived component appears predominant as recorded by chondritic to supra-chondritic Hf
238 initial isotopic compositions. This pattern appears very similar to that of the Booldeega
239 diamictite. Zircons from the sandstones of the upper part of the Kazput Formation yielded a
240 range of $^{207}\text{Pb}/^{206}\text{Pb}$ ages between 2430 and 3230 Ma and initial $\epsilon_{\text{Hf}}(t)$ values between -7.4
241 and +6.4 (Fig. 3d). Most of the data display supra-chondritic Hf initial isotopic composition,
242 which demonstrates a prevailing input of mantle-derived component. These Hf isotope data
243 share similarities with those of the MBM diamictite (Fig. 4b).

244 Zircons from the Beasley River Quartzite yielded a range of $^{207}\text{Pb}/^{206}\text{Pb}$ ages between
245 2450 and 3230 Ma and initial $\epsilon_{\text{Hf}}(t)$ values between -10.4 and +5.9 (Fig. 3e). Almost equal
246 proportions of sub- and supra-chondritic zircon Hf isotopic composition were recorded. The
247 results present similar ages and Hf initial isotopic signatures than the other formations.

248 When all the analyses are plotted together (Fig. 3f), two major points are evidenced:
249 (1) the majority of the U-Pb ages range from 2.45 Ga to 3.20 Ga, and (2) more than 70% of
250 the analyses plot in the supra-chondritic field, which demonstrates the prevailing occurrence
251 of mantle-derived materials. Considering a $^{176}\text{Lu}/^{177}\text{Hf}$ ratio of 0.015 typical of bulk
252 continental crust (Griffin et al., 2004), the data can be confined between two model age lines
253 at *c.a.* 2.54 Ga and 3.20 Ga. As discussed by Vervoort and Kemp (2016), these ages are
254 dependent of a number of factors such as mantle source models and Lu/Hf values and
255 therefore should not be taken at face values and only considered as indicative of the time
256 when the source of the zircon's protoliths separated from the mantle.

257

258 **6. The first record of a Hadean zircon in the Pilbara Craton**

259 Four analyses of a single zircon crystal (120 x 60 μm in size) from sample Pi 09.07 of
260 the 2.45 Ga Boolgeeda diamictite at the Boundary Ridge locality yielded an upper intercept at
261 4013 ± 69 Ma. One of the analyses is concordant at 3998 ± 15 Ma (Fig. 4a and Table 2). The
262 Th/U ratios (0.2 to 0.8) favour a magmatic origin for the crystal. This zircon crystal represents
263 the first record of a Hadean relict within the Pilbara Craton. The zircon shows a negative
264 initial $\epsilon_{\text{Hf}}(t)$ value of -3.3 (Fig. 4b), which is consistent with the filtered Hf isotope dataset
265 for Jack Hills zircons in the Yilgarn Craton after Whitehouse et al. (2017). These authors
266 interpreted these data to represent zircon growth events during the Hadean, although the
267 nature of these events remains unconstrained. As discussed by Kemp et al. (2010), it is
268 therefore tempting, although speculative, to propose the crystallization of this Hadean zircon
269 in a magma source derived from the Depleted Mantle and carrying an evolved crustal
270 component earlier than 4.0 Ga. Depending on the nature of the crust (i.e. felsic or mafic) from
271 which the zircon crystallized, the Hf model ages are 4.27 or 4.49 Ga (Fig. 5b), which are in

272 agreement with Nd model ages obtained on samples from the 3.5 Ga Dresser Formation in the
273 Pilbara Craton (Tessalina *et al.*, 2010).

274

275 **7. Discussion**

276 **7.1 Summary of the zircon Hf isotopic signatures**

277 Zircon Hf isotopic data appear relatively consistent throughout the sedimentary
278 sequence of the Boolgeeda Iron Formation and the Turee Creek Group. About 70 % of the
279 analyzed zircons are characterized by chondritic to supra-chondritic Hf isotopic compositions,
280 with 30% of the analyses showing $\epsilon_{\text{Hf}}(t)$ values higher than +3. Most of the data plot between
281 model age lines at ~2.5 and 3.2 Ga (Fig. 3f). Although these ages are not representative of the
282 time of extraction of juvenile magmas from the depleted mantle (Vervoort and Kemp, 2016),
283 the lack of evidence for a crustal contribution older than 3.2 Ga indicates that the majority of
284 the data is consistent with a sub-continuous or multi-episodic crust generation through
285 depleted mantle differentiation after about 3.2 Ga. Eo- and Paleo-Archean crustal rocks did
286 not contribute significantly to the sedimentary material of the Boolgeeda diamictite and TCG
287 sediments, in agreement with zircon age peaks at 2.54, 2.68, 2.82, 2.95 and 3.15 Ga
288 (Caquineau *et al.*, 2018). The recognition that the zircon Hf isotopic compositions of the
289 Beasley River Quartzite (unconformably overlying the Turee Creek Group) are
290 undistinguishable from the Boolgeeda and Turee Creek zircon Hf compositions further
291 support the notion of a relatively homogeneous source over a substantial long period of time
292 between ~2.45 and 2.0 Ga.

293

294 **7.2 Paleogeographic significance of the zircon Hf isotopic data**

295 To evaluate the main sources of the Boolgeeda diamictite and the TCG sediments we
296 have considered our isotope data in the broader context of published radiogenic isotope data

297 for igneous rocks from several representative cratons in Australia (Glenburgh Terrane and
298 Pilbara and Yilgarn cratons), South Africa (Kaapvaal Craton) and North America (Superior
299 Craton).

300

301 *7.2.1 Glenburgh Terrane*

302 The Glenburgh Terrane has been defined as an exotic continental fragment that
303 collided with the southwestern margin of the Pilbara Craton during the ~2.22 – 2.15 Ga
304 Ophthalmian orogeny (Blake & Barley, 1992; Martin *et al.*, 2000; Johnson *et al.*, 2011).
305 Zircons from the 2430–2555 Ma old Halfway Gneiss, which represents the oldest component
306 of the Glenburgh Terrane, have a variety of Paleoproterozoic and Archean ages and a
307 relatively narrow range of sub-chondritic Hf isotopic compositions interpreted to reflect
308 reworking of an older 2600–2730 Ma crust (Johnson *et al.*, 2011). Some of the ages and Hf
309 isotopic data overlap those of the Boolgeeda diamictite and TCG detrital zircons (Fig. 5a), but
310 only a few of the zircons from the Glenburgh Terrane match the supra-chondritic values
311 recorded in our samples. Hence, while the Glenburgh Terrane may have contributed to Turee
312 Creek basin (Krapež *et al.*, 2017), it cannot be considered as a major source material of the
313 Boolgeeda diamictite and TCG sediments.

314

315 *7.2.2 Pilbara Craton.*

316 Magmatic zircons from the Pilbara Craton mainly crystallized at about 3.2–3.3 Ga or
317 before (Fig. 5b; Gardiner *et al.*, 2017; 2019; Petersson *et al.*, 2019 and references therein).
318 These TTG suites are interpreted as derived from partial melting of basaltic crust with the
319 contribution from older felsic components (e.g. xenoliths of the Strelley Pool Formation up to
320 3.7 Ga; Gardiner *et al.*, 2019). After ca. 3.3 Ga, the trend toward more negative Hf values is
321 interpreted by these authors as recording more extensive crust reworking. In the eastern

322 Pilbara Craton, multiple pulses of magmatism produced voluminous granitic complexes from
323 3.5 Ga to 2.95-2.92 Ga and 2.85-2.82 Ga. This age range is much older than the one recorded
324 for the Boolgeeda and TCG detrital zircons. As shown in figure 5b, the Hf isotope data from
325 both central and eastern Pilbara do not overlap with those of the TCG, ruling out any
326 significant contribution of the Pilbara Craton to this sedimentary succession.

327

328 *7.2.3 Northern Yilgarn Craton*

329 The Yilgarn Craton located south of the Pilbara Craton and Glenburgh Terrane
330 consists of several distinct tectonic entities (the Narryer, Youanmi and South West terranes
331 and the Eastern Goldfields Superterrane) that have been accreted to the amalgamated Pilbara
332 Craton–Glenburgh Terrane during the ~1.96 – 2.0 Ga Glenburgh orogeny (Occhipinti *et al.*
333 2004; Johnson *et al.*, 2011, 2013), about 200 Ma after deposition of the TCG (i.e. 2210 Ma,
334 Müller *et al.*, 2005). Available Hf isotope data of magmatic zircons from the Yilgarn Craton
335 (Griffin *et al.*, 2004) show similarities with our results. Both display supra-chondritic values
336 during Neoproterozoic times, a quite similar range of ages and only a few zircon grains older than
337 3.2 Ga (Fig. 5c). However, the Boolgeeda sample also contains younger grains with supra-
338 chondritic Hf isotopic compositions down to 2.4 Ga compared to the Yilgarn zircons (2.6 Ga).
339 Nevertheless, the strong overlap between our Hf isotope data and those of the Yilgarn Craton
340 together with the occurrence of a Hadean zircon crystal in the Boolgeeda diamictite, similar in
341 age and Hf isotopic signature to the Jack Hills zircons (Whitehouse *et al.*, 2017 and references
342 therein) makes the Yilgarn Craton a possible source for the Boolgeeda glacial horizon. This
343 conclusion is in apparent contradiction with the cratonisation history and paleogeographic
344 reconstruction of the Yilgarn and Pilbara cratons, which are considered to be part of two
345 different supercratons during the early Paleoproterozoic (Cheney, 1996; de Kock *et al.*, 2009;
346 Smirnov *et al.*, 2013), and to have accreted during the ~1.96 – 2.0 Ga Glenburgh orogeny

347 (Johnson *et al.*, 2013), that is about 200 Ma later than the upper age limit of the TCG (Müller
348 *et al.*, 2005; see chapter 7.4 below for further discussion).

349

350 *7.2.4 Kaapvaal Craton*

351 Magmatic zircons from the Kaapvaal Craton mainly crystallized from material that is
352 much older than the one recorded for the Boolgeeda and TCG detrital zircons (Fig. 5e, Zeh *et*
353 *al.*, 2009). Assuming the ‘Vaalbara’ hypothesis (Cheney, 1996; deKock *et al.*, 2009; see
354 Eriksson *et al.*, 2011 for an alternative hypothesis), the post-2080 Ma Magaliesberg Quartzite,
355 which is stratigraphically located at the top of the Transvaal Supergroup, could represent an
356 analog of the TCG sediments. Two populations of zircons can be distinguished on the basis of
357 U-Pb ages and initial $\epsilon_{\text{Hf}}(t)$ values (Fig. 5d, Zeh *et al.*, 2016). One is characterized by Archean
358 age and both supra- and sub-chondritic Hf isotopic signatures; the second population shows
359 Paleoproterozoic age and sub-chondritic Hf isotopic compositions (Fig. 5d). Our data from
360 the Boolgeeda diamictite and TCG sediments mostly plot between these two zircon
361 populations, which argue for distinctly different sources. The zircons of Archean age
362 (population 1) of the Magaliesberg Quartzite have been attributed to the erosion of the
363 Pietersburg Block of the Kaapvaal Craton. The absence of major overlap with the Boolgeeda
364 and TCG zircons indicates that this terrane was not a major source of the Boolgeeda and TCG
365 sediments. The source of the Paleoproterozoic zircons (population 2) is unclear. The dominant
366 crustal recycling signature indicated by the strongly negative $\epsilon_{\text{Hf}}(t)$ values could be linked to
367 the break-up of the Vaalbara Supercontinent between ~2.45 and 2.2 Ga. Accordingly, U-Pb
368 geochronology, Hf isotopes and paleogeographic reconstructions all argue against the
369 Kaapvaal Craton as a possible source for the Boolgeeda diamictite and TCG sediments.

370

371 *7.2.5 Superior Craton*

372 The Paleoproterozoic glacial events also affected the Superior Craton during the
373 deposition of the Huronian Supergroup (Young, 1970). Available zircon Hf isotopic data from
374 the Superior Craton are mostly supra-chondritic (Corfu and Noble, 1992; Corfu and Stott,
375 1993 and 1996; Davis *et al.*, 2005) and characterize samples older than 2.7 Ga. As shown in
376 figure 5f, data from the Boolgeeda diamictite, the TCG and the Superior Craton are mostly
377 different, likely due to different cratonization histories.

378

379 **7.3 Implications for a prevailing mantle-derived magma source**

380 The comparison of our detrital zircon $\epsilon_{\text{Hf}}(t)$ data with those of the literature shows that
381 the Glenburgh Terrane and the Pilbara, Kaapvaal, Zimbabwe and Superior cratons did not
382 contribute significantly to the Boolgeeda and TCG sediments. Over 80% of the analyzed
383 zircons yielded ages spanning those of the Hamersley and Fortescue groups between 2.45 and
384 2.78 Ga (Arndt *et al.*, 1991; Trendall *et al.*, 2004). This supports the interpretation that the
385 Boolgeeda Iron Formation and the TCG successions mainly derived from the erosion of the
386 underlying Hamersley and Fortescue groups. Martin *et al.* (2000) and Van Kranendonk *et al.*,
387 (2015), originally proposed this interpretation based on geological and stratigraphic
388 reconstructions. As shown by their supra-chondritic Hf isotopic compositions, detrital zircons
389 from the Boolgeeda diamictite and the TCG successions preferentially incorporated mantle-
390 derived material. This is not typical of terrigenous sedimentation in cratonic settings, where
391 erosion products generally record diverse lithologies of various ages reflecting the erosion of
392 older crustal basement characterized by sub-chondritic Hf isotopic compositions. The marked
393 differences in zircon Hf compositions between the Boolgeeda diamictite and TCG sediments,
394 and the Pilbara Craton (Fig. 5b) and the Glenburgh Terrane (Fig. 5a) is particularly striking,
395 as these two continental blocks are the most likely sources in addition to the Fortescue and
396 Hamersley groups.

397 A possible explanation for the minor proportion of sub-chondritic Hf isotopic
398 compositions in the Boolgeeda diamictite and TCG zircons can be a lack of emerged crustal
399 blocks carrying a sub-chondritic Hf isotope record at the time of deposition. This is
400 particularly critical for the Pilbara Craton to the north and the Glenburgh Terrane to the south,
401 which have been considered as the main potential sources of the TCG successions (e.g.,
402 Krapez *et al.*, 2017). The scarcity or lack of surface exposure of Pilbara Craton lithologies
403 seems unlikely as it comprises a wide range of rocks of different ages between 2.7 and 3.5 Ga
404 that crop out on thousands of km² (Hickman and Van Kranendonk, 2012). Additionally, the
405 Mount Bruce Supergroup, which contains the TCG and the underlying Hamersley and
406 Fortescue groups, was unconformably deposited on the basement of the Pilbara Craton (Arndt
407 *et al.*, 1991). The lack of detrital zircon much older than 3.2 Ga within the TCG may also be
408 related to the absence of efficient transport and/or erosion of Archean material. However,
409 sandstones of the Koolbye Formation and of the Beasley River Quartzite (see Fig. 2) have
410 been interpreted as fluvial to shallow marine, and fluvial to aeolian deposits, respectively
411 (Mazumder and Van Kranendonk, 2013; Mazumder *et al.*, 2015), witnessing the activities of
412 rivers and wind during the deposition of the TCG.

413 About 75% of the zircons analyzed here come from glaciogenic deposits
414 (i.e. Boolgeeda, MBM and second Kungarra diamictites). Glaciogenic rocks typically derive
415 from widespread physical erosion of the upper continental crust by icesheets and are therefore
416 geographically dispersed (e.g., Gaschnig *et al.*, 2014). The supra-chondritic Hf isotopic
417 composition recorded in the major part of the zircon populations from the Boolgeeda and
418 Turee Creek sediments is at odd with a sub-chondritic Hf isotopic composition generally
419 expected for the recycled upper continental crust (e.g., Condie, 2014). The Welli Wooli
420 Formation of the Hamersley Group, comprising basalts, dolerites as well as the Woongarra
421 Rhyolite, has been interpreted as a subaerial bimodal Large Igneous Province (LIP) emplaced

422 at 2450 Ma (Barley *et al.*, 1997). This intense volcanic event would have produced large
423 volumes of lavas ($>30\,000\text{ km}^3$, Barley *et al.*, 1997) covering the continental surface at the
424 time of the glacial event recorded in the Boolgeeda Iron Formation, and possibly during the
425 subsequent deposition of the overlying TCG. Thus, it is suggested that the erosion of these
426 volcanic materials by ice sheets and their incorporation as terrigenous detritus within the
427 glacial diamictites of the Boolgeeda and Turee Creek sedimentary successions could account
428 for the supra-chondritic Hf isotopic compositions of many analyzed zircons (Fig. 3).
429 Similarly, zircons corresponding to age peaks at 2540 and 2680 Ma could have been produced
430 by previous volcanic eruptions attending deposition of the Fortescue and Hamersley groups
431 and preserved at the continental surface. It is noteworthy that the Hf isotopic compositions of
432 zircons from non-glacial sediments, i.e the sandstones of the Kazput Formation and the
433 Beasley River Quartzite, do not exhibit different $\epsilon_{\text{Hf}}(t)$ vs. time patterns than the zircons from
434 the glacial diamictites (Fig. 3d and 3e). This suggests that these sandstones and quartzites may
435 represent the erosional products of the same kind of LIP-derived volcanic material involved in
436 the formation of the glacial diamictites.

437 The hypothesis that the Earth's surface was covered by juvenile lavas during the
438 Archean-Proterozoic transition, at least on the Pilbara Craton, may also have important
439 implications on the concomitant rise of atmospheric oxygen (the GOE) and drastic cooling of
440 the Earth's surface. As shown by Kump and Barley (2007), a rapid increase of the subaerial
441 proportion of LIPs at ca. 2.5 Ga would have abruptly and permanently diminished the
442 predominant sink for oxygen (submarine volcanism) and increased the sink for carbon dioxide
443 (silicate weathering). This in turn would have enabled oxygen to build up into the atmosphere
444 and triggered a glacial period (Cox *et al.*, 2016).

445

446 **7.4 Geodynamic implications of a Hadean zircon in the Pilbara Craton**

447 Several studies have suggested that the Pilbara Craton developed on a Hadean
448 continental substrate (Smithies *et al.*, 2003, 2007; Champion, 2013; Tessalina *et al.*, 2010).
449 Such ancient crust has, however, never been identified in outcrop. The oldest rock reported so
450 far is the metamorphosed Mount Webber Gabbro located in the western part of the Shaw
451 Granitic Complex, which formed at 3.59–3.58 Ga (Petersson *et al.*, 2019), and therefore
452 predates the oldest known volcanic activity of the 3.53–3.23 Ga Pilbara Supergroup within
453 the East Pilbara Terrane. Many Hadean zircons identified in Western Australia are from the
454 Jack Hills meta-conglomerate of the Narryer Formation in the neighboring Yilgarn Craton
455 (Wilde *et al.*, 2001, see above). The significant overlap between our Hf isotope data and those
456 of the Yilgarn Craton, together with the occurrence of an Hadean zircon crystal preserved in
457 the Boolgeeda diamictite with similar age and Hf isotopic signature than the Jack Hills
458 zircons (Whitehouse *et al.*, 2017 and references therein) makes the Yilgarn Craton a possible
459 source material for the Boolgeeda glacial horizon. Available paleogeographic reconstructions
460 based on paleomagnetic data and cratonization history suggest that during the
461 Paleoproterozoic, the Yilgarn and Pilbara cratons were part of two different supercratons, the
462 Zimgarn comprising the Zimbabwe and Yilgarn cratons (Smirnov *et al.*, 2013), and the
463 Vaalbara composed of the Kaapvaal and Pilbara cratons (Cheney, 1996; de Kock *et al.*, 2009).
464 This argues against Yilgarn and Zimbabwe cratons as a source for the Turee Creek and
465 Boolgeeda zircons (Fig. 5c and d). However, Gumsley *et al.* (2017) recently proposed that
466 despite their dissimilar cratonization history, the Zimbabwe and Yilgarn cratons may have
467 been part of a much larger supercraton, the Supervaalbara, which comprised Superior,
468 Wyoming, Hearne, Kola, Karelia, Kaapvaal, and Pilbara. Considering that the 2.45 Ga old
469 Boolgeeda glacial diamictite has been correlated with several glacial horizons in South Africa
470 (pre-2.43 Ga Makganyene Formation of the Griqualand West region), North America (2.45
471 Ga Ramsay Lake Formation of the Superior Province) and Arctic Europe (2.43 Ga

472 Urkkavaara and Polisarka Formations of Fennoscandian Craton), Caquineau et al., (2018)
473 proposed that it could represent a global glacial event. Accordingly, although the Yilgarn
474 craton may have been located at a latitude very different from that of the Turee Creek Basin
475 during the Paleoproterozoic, it cannot be excluded that the Hadean zircon preserved in the
476 Boolgeeda diamictite has been transported by icesheets over significant distances.

477

478 **8. Conclusions**

479 Our zircon U-Pb-Hf isotopic study indicates that Neoproterozoic (to Mesoproterozoic)
480 mantle-derived material was the major detrital component delivered to the Hamersley basin
481 during deposition of the glacial and non-glacial sediments of the Boolgeeda Iron Formation
482 and the Turee Creek Group, as well as the overlying Beasley River Quartzite. In this study,
483 we discovered the first occurrence of a Hadean zircon grain in the Pilbara Craton. Rare
484 materials older than 3.2 Ga were recycled within the Turee Creek basin. Our data highlight a
485 much larger proportion of mantle-derived source materials than in comparable cratons and
486 basins, for which zircon Hf isotopic data are available. The predominantly supra-chondritic
487 Hf isotopic signatures can be explained by a relatively large proportion of subaerial volcanic
488 rocks during deposition of the underlying 2.45 – 2.78 Ga Hamersley and Fortescue groups. A
489 local origin of the material delivered to the Turee Creek basin is in agreement with an
490 intracontinental depositional setting. The massive emplacement of subaerial LIPs in the
491 Hamersley and Fortescue groups during the Neoproterozoic followed by subsequent erosion and
492 alteration attending the formation of the Turee Creek basin may have triggered the rise of
493 atmospheric oxygen and associated Paleoproterozoic glacial events.

494

495 ***Acknowledgements***

496 This work was supported by grants from the UnivEarths Labex Program at Sorbonne
497 ParisCité (ANR-10-LABX-0023 and ANR-11-IDEX-0005-02), the Programme National de
498 Planétologie and the São Paulo Research Foundation (FAPESP, grant 2015/16235-2). This
499 article has benefited from the constructive comments of Dr. Fernando Corfu and an
500 anonymous reviewer as well as associate editor Pr. Wilson Teixeira.

501

502 **References**

503 1 Arndt, N.T., Nelson, D.R., Compston, W., Trendall, A.F., and Thorne, A.M. (1991) The age
504 of the Fortescue Group, Hamersley Basin, Western Australia, from ion microprobe zircon
505 U-Pb results. *Australian Journal of Earth Sciences*, **38**, 261–281.

506 2 Barley, M.E., Pickard, A.L., and Sylvester, P.J. (1997) Emplacement of a large igneous
507 province as a possible cause of banded iron formation 2.45 billion years ago. *Nature*, **385**,
508 55–58.

509 3 Blichert-Toft, J. and Albarède, F. (2008) Hafnium isotopes in Jack Hills zircons and the
510 formation of the Hadean crust. *Earth and Planetary Science Letters*, **265**, 686–702.

511 4 Blichert-Toft, J. and Puchtel, I.S. (2010) Depleted mantle sources through time: evidence
512 from Lu–Hf and Sm–Nd isotope systematics of Archean komatiites. *Earth and Planetary
513 Science Letters*, **297**, 598–606.

514 5 Caquineau, T., Paquette, J.-L., and Philippot, P. (2018) U-Pb detrital zircon geochronology
515 of the Turee Creek Group, Hamersley Basin, Western Australia: Timing and correlation of
516 the Paleoproterozoic glaciations. *Precambrian Research*, **307**, 34–50.

517 6 Cheney, E.S. (1996) Sequence stratigraphy and plate tectonic significance of the Transvaal
518 succession of southern Africa and its equivalent in Western Australia. *Precambrian
519 Research*, **79**, 3–24.

520 7 Condie, K.C. (2014) Growth of continental crust: a balance between preservation and

- 521 recycling. *Mineralogical Magazine*, **78**, 623–638.
- 522 8 Corfu, F. and Noble, S.R. (1992) Genesis of the southern Abitibi greenstone belt, Superior
523 Province, Canada: evidence from zircon Hf isotope analyses using a single filament
524 technique. *Geochimica et Cosmochimica Acta*, **56**: 2081-2097.
- 525 9 Corfu, F. and Stott, G.M. (1993) Age and petrogenesis of two late Archean magmatic suites,
526 northwestern Superior Province, Canada: zircon U-Pb and Lu-Hf isotopic relations.
527 *Journal of Petrology*, **34**: 817-838.
- 528 10 Corfu, F. and Stott, G.M. (1996) Hf isotopic composition and age constraints on the
529 evolution of the Archean central Uchi Subprovince, Ontario, Canada. *Precambrian*
530 *Research*, **78**: 53-63.
- 531 11 Cox, G.M., Halverson, G.P., Stevenson, R.K., Vokaty, M., Poirier, A., Kunzmann, M., Li,
532 Z.-X., Denyszyn, S.W., Strauss, J.V., and Macdonald, F.A. (2016) Continental flood
533 basalt weathering as a trigger for Neoproterozoic Snowball Earth. *Earth and Planetary*
534 *Science Letters*, **446**, 89–99.
- 535 12 Davis, D.W., Amelin, Y., Nowell, G.M., and Parrish, R.R. (2005) Hf isotopes in zircon
536 from the western Superior province, Canada: Implications for Archean crustal
537 development and evolution of the depleted mantle reservoir. *Precambrian Research*, **140**,
538 132–156.
- 539 13 Eriksson, P.G., Lenhardt, N., Wright, D.T., Mazumder, R., and Bumby, A.J. (2011) Late
540 Neoproterozoic-Palaeoproterozoic supracrustal basin-fills of the Kaapvaal craton: Relevance
541 of the supercontinent cycle, the “Great Oxidation Event” and “Snowball Earth”? *Marine*
542 *and Petroleum Geology*, **28**, 1385–1401.

- 543 14 Fisher, C.M., Hanchar, J.M., Samson, S.D., Dhuime, B., Blichert-Toft, J., Vervoort, J.,
544 Lam, R. (2011) Synthetic zircon doped with hafnium and rare earth elements: a reference
545 material for in situ hafnium isotope analysis. *Chemical Geology*, **286**, 32-47.
- 546 15 Gaillard, F., Scaillet, B. and Arndt, N.T. (2011) Atmospheric oxygenation caused by a
547 change in volcanic degassing pressure. *Nature*, **478**, 229-232.
- 548 16 Gardiner, N.J., Hickman, A.H., Kirkland, C.L., Lu, Y., Johnson, T., and Zhao, J.-X. (2017)
549 Processes of crust formation in the early Earth imaged through Hf isotopes from the East
550 Pilbara Terrane. *Precambrian Research*, **297**, 56–76.
- 551 17 Gardiner, N.J., Wacey, D., Kirkland, C.L., Johnson, T.E., Jeon, H. (2019) Zircon U-Pb,
552 Lu-Hf and O isotopes from the 3414 Ma Strelley Pool Formation, East Pilbara Terrane,
553 and the Paleoarchean emergence of a cryptic cratonic core. *Precambrian Research*, **321**,
554 64-84.
- 555 18 Gaschnig, R.M., Rudnick, R.L., McDonough, W.F., Kaufman, A.J., Hu, Z., and Gao, S.
556 (2014) Onset of oxidative weathering of continents recorded in the geochemistry of
557 ancient glacial diamictites. *Earth and Planetary Science Letters*, **408**, 87–99.
- 558 19 Griffin, W.L., Belousova, E.A., Shee, S.R., Pearson, N.J., and O'Reilly, S.Y. (2004)
559 Archean crustal evolution in the northern Yilgarn Craton: U–Pb and Hf-isotope evidence
560 from detrital zircons. *Precambrian Research*, **131**, 231–282.
- 561 20 Gumsley, A.P., Chamberlain, K.R., Bleeker, W., Söderlund, U., Kock, M.O. de, Larsson,
562 E.R., and Bekker, A. (2017) Timing and tempo of the Great Oxidation Event. *Proceedings*
563 *of the National Academy of Sciences*, **114**, 1811–1816.
- 564 21 Hawkesworth, C.J., Dhuime, B., Pietranik, A.B., Cawood, P.A., Kemp, A.I.S., and Storey,
565 C.D. (2010) The generation and evolution of the continental crust. *Journal of the*
566 *Geological Society*, **167**, 229–248.
- 567 22 Hickman, A.H. and Van Kranendonk, M.J.V. (2012) Early Earth Evolution: Evidence

- 568 from the 3.5-1.8 Ga Geological History of the Pilbara Region of Western Australia.
569 *Episodes*, **35**, 283–297.
- 570 23 Holland, H.D. (2002) Volcanic gases, black smokers, and the great oxidation event.
571 *Geochimica et Cosmochimica Acta*, **66**, 3811–3826.
- 572 24 Hu, Z., Liu, Y., Gao, S., Liu, W., Zhang, W., Tong, X., Lin, L., Zong, K., Li, M., Chen, H.,
573 Zhou, L., Yang L. (2012) Improved in situ Hf isotope ratio analysis of zircon using newly
574 designed X skimmer cone and jet sample cone in combination with the addition of nitrogen
575 by laser ablation multiple collector ICP-MS. *Journal of analytical Atomic Spectrometry*, **27**,
576 1391-1399.
- 577 25 Hurai, V., Paquette, J.L, Huraiová, M., Konečný, P. (2010) Age of deep crustal magmatic
578 chambers in the intra-Carpathian back-arc basin inferred from LA-ICPMS U-Th-Pb dating
579 of zircon and monazite from igneous xenoliths in alkali basalts. *Journal of Volcanology
580 and Geothermal Research*, **198**, 275-287.
- 581 26 Iizuka, T., Yamaguchi, T., Hibiya, Y., and Amelin, Y. (2015) Meteorite zircon constraints
582 on the bulk Lu–Hf isotope composition and early differentiation of the Earth. *Proceedings
583 of the National Academy of Sciences*, **112**, 5331–5336.
- 584 27 Iizuka, T., Yamaguchi, T., Itano, K., Hibiya, Y., and Suzuki, K. (2017) What Hf isotopes
585 in zircon tell us about crust–mantle evolution. *Lithos*, **274–275**, 304–327.
- 586 28 Johnson, S.P., Sheppard, S., Rasmussen, B., Wingate, M.T.D., Kirkland, C.L., Muhling,
587 J.R., Fletcher, I.R., and Belousova, E.A. (2011) Two collisions, two sutures: Punctuated
588 pre-1950 Ma assembly of the West Australian Craton during the Ophthalmian and
589 Glenburgh Orogenies. *Precambrian Research*, **189**, 239–262.
- 590 29 Johnson, S.P., Thorne, A.M., Tyler, I.M., Korsch, R.J., Kennett, B.L.N., Cutten, H.N.,
591 Goodwin, J., Blay, O., Blewett, R.S., Joly, A., and others. (2013) Crustal architecture of
592 the Capricorn Orogen, Western Australia and associated metallogeny. *Australian Journal*

- 593 *of Earth Sciences*, **60**, 681–705.
- 594 30 Kemp, A.I.S., Wilde, S.A., Hawkesworth, C.J., Coath, C.D., Nemchin, A., Pidgeon, R.T.,
595 Vervoort, J.D., DuFrane, S.A. (2010) Hadean crustal evolution revisited: New constraints
596 from Pb–Hf isotope systematics of the Jack Hills zircons. *Earth and Planetary Science*
597 *Letters*, **296**, 45–56.
- 598 31 Kemp, A.I.S., Hickman, A.H., Kirkland, C.L., and Vervoort, J.D. (2015) Hf isotopes in
599 detrital and inherited zircons of the Pilbara Craton provide no evidence for Hadean
600 continents. *Precambrian Research*, **261**, 112–126.
- 601 32 de Kock, M.O., Evans, D.A.D., and Beukes, N.J. (2009) Validating the existence of
602 Vaalbara in the Neoproterozoic. *Precambrian Research*, **174**, 145–154.
- 603 33 Kopp, R.E., Kirschvink, J.L., Hilburn, I.A., and Nash, C.Z. (2005) The Paleoproterozoic
604 snowball Earth: A climate disaster triggered by the evolution of oxygenic photosynthesis.
605 *Proceedings of the National Academy of Sciences of the United States of America*, **102**,
606 11131–11136.
- 607 34 Krapež, B., Müller, S.G., Fletcher, I.R., and Rasmussen, B. (2017) A tale of two basins?
608 Stratigraphy and detrital zircon provenance of the Palaeoproterozoic Turee Creek and
609 Horseshoe basins of Western Australia. *Precambrian Research*, **294**, 67–90.
- 610 35 Kump, L.R. and Barley, M.E. (2007) Increased subaerial volcanism and the rise of
611 atmospheric oxygen 2.5 billion years ago. *Nature*, **448**, 1033–1036.
- 612 36 Martin, D.M. (1999) Depositional setting and implications of Paleoproterozoic
613 glaciomarine sedimentation in the Hamersley Province, Western Australia. *Geological*
614 *Society of America Bulletin*, **111**, 189–203.
- 615 37 Martin, D.M., Li, Z.X., Nemchin, A.A., and Powell, C.M. (1998) A pre-2.2 Ga age for
616 giant hematite ores of the Hamersley Province, Australia? *Economic Geology*, **93**, 1084–
617 1090.

- 618 38 Martin, D.M., Powell, C.M., and George, A.D. (2000) Stratigraphic architecture and
619 evolution of the early Paleoproterozoic McGrath Trough, Western Australia. *Precambrian*
620 *Research*, **99**, 33–64.
- 621 39 Mazumder, R. and Van Kranendonk, M.J. (2013) Palaeoproterozoic terrestrial
622 sedimentation in the Beasley River Quartzite, lower Wyloo Group, Western Australia.
623 *Precambrian Research*, **231**, 98–105.
- 624 40 Mazumder, R., Van Kranendonk, M.J., and Altermann, W. (2015) A marine to fluvial
625 transition in the Paleoproterozoic Koolbye Formation, Turee Creek Group, Western
626 Australia. *Precambrian Research*, **258**, 161–170.
- 627 41 Melezhik, V.A., Young, G.M., Eriksson, P.G., Altermann, W., Kump, L.R., and Lepland,
628 A. (2013) 7.2 Huronian-Age Glaciation. Pp. 1059–1109 in: *Reading the Archive of*
629 *Earth's Oxygenation* (V.A. Melezhik, A.R. Prave, E.J. Hanski, A.E. Fallick, A. Lepland,
630 L.R. Kump, and H. Strauss, editors). *Frontiers in Earth Sciences*, Springer Berlin
631 Heidelberg.
- 632 42 Moyen, J.-F., Paquette, J.-L., Ionov, D.A., Gannoun, A., Korsakov, A.V., Golovin, A.V.,
633 and Moine, B.N. (2017) Paleoproterozoic rejuvenation and replacement of Archaean
634 lithosphere: Evidence from zircon U–Pb dating and Hf isotopes in crustal xenoliths at
635 Udachnaya, Siberian craton. *Earth and Planetary Science Letters*, **457**, 149–159.
- 636 43 Müller, S.G., Krapež, B., Barley, M.E., and Fletcher, I.R. (2005) Giant iron-ore deposits of
637 the Hamersley province related to the breakup of Paleoproterozoic Australia: New
638 insights from in situ SHRIMP dating of baddeleyite from mafic intrusions. *Geology*, **33**,
639 577–580.
- 640 44 Nebel-Jacobsen, Y., Nebel, O., Wille, M., Cawood, P.A. (2018) A non-zircon Hf isotope
641 record in Archean black shales from the Pilbara craton confirms changing crustal
642 dynamics ca. 3 Ga ago. *Nature Scientific Reports*, **8**, 1-7.

- 643 45 Paquette, J.L., Ionov, D.A., Agashev, A.M., Gannoun, A., and Nikolenko, E.I. (2017) Age,
644 provenance and Precambrian evolution of the Anabar shield from U-Pb and Lu-Hf isotope
645 data on detrital zircons, and the history of the northern and central Siberian craton.
646 *Precambrian Research*, **301**, 134–144.
- 647 46 Petersson, A., Kemp, A.I.S., Hickman, A.H., Whitehouse, M.J., Martin, L., Gray, C.M.
648 (2019) A new 3.59 Ga magmatic suite and a chondritic source to the east Pilbara Craton.
649 *Chemical Geology*, **511**, 51-70.
- 650 47 Philippot, P., Avila, J., Killingsworth, B., Tessalina, S., Baton, F., Caquineau, T., Muller,
651 E., Pecoits, E., Cartigny, P., Lalonde, S., Ireland, T., Thomazo, C., Van Kranendonk, M.,
652 Busigny, V. (2018) Globally asynchronous sulphur isotope require re-definition of the
653 Great Oxidation Event. *Nature Communications*, **9**, 1-10.
- 654 48 Rasmussen, B., Fletcher, I.R., and Sheppard, S. (2005) Isotopic dating of the migration of
655 a low-grade metamorphic front during orogenesis. *Geology*, **33**, 773–776.
- 656 49 Rasmussen, B., Bekker, A., and Fletcher, I.R. (2013) Correlation of Paleoproterozoic
657 glaciations based on U–Pb zircon ages for tuff beds in the Transvaal and Huronian
658 Supergroups. *Earth and Planetary Science Letters*, **382**, 173–180.
- 659 50 Scherer, E., Münker, C., and Mezger, K. (2001) Calibration of the Lutetium-Hafnium
660 Clock. *Science*, **293**, 683–687.
- 661 51 Smirnov, A.V., Evans, D.A., Ernst, R.E., Söderlund, U., and Li, Z.-X. (2013) Trading
662 partners: tectonic ancestry of southern Africa and western Australia, in Archean
663 supercratons Vaalbara and Zimgarn. *Precambrian Research*, **224**, 11–22.
- 664 52 Teitler, Y., Le Hir, G., Fluteau, F., Philippot, P., and Donnadiou, Y. (2014) Investigating
665 the Paleoproterozoic glaciations with 3-D climate modeling. *Earth and Planetary Science
666 Letters*, **395**, 71–80.
- 667 53 Tessalina, S.G., Bourdon, B., Van Kranendonk, M., Birck, J.-L., and Philippot, P. (2010)

- 668 Influence of Hadean crust evident in basalts and cherts from the Pilbara Craton. *Nature*
669 *Geoscience*, **3**, 214–217.
- 670 54 Trendall, A.F. (1981) The Lower Proterozoic Meteorite Bore Member, Hamersley Basin,
671 Western Australia. *Earth's Pre-Pleistocene glacial record: Cambridge University Press*,
672 *Cambridge*, 555–557.
- 673 55 Trendall, A.F., Compston, W., Nelson, D.R., De Laeter, J.R., and Bennett, V.C. (2004)
674 SHRIMP zircon ages constraining the depositional chronology of the Hamersley Group,
675 Western Australia. *Australian Journal of Earth Sciences*, **51**, 621–644.
- 676 56 Van Kranendonk, M.J., Mazumder, R., Yamaguchi, K.E., Yamada, K., and Ikehara, M.
677 (2015) Sedimentology of the Paleoproterozoic Kungarra Formation, Turee Creek Group,
678 Western Australia: A conformable record of the transition from early to modern Earth.
679 *Precambrian Research*, **256**, 314–343.
- 680 57 Van Kranendonk, M.J.V. and Mazumder, R. (2015) Two Paleoproterozoic glacio-eustatic
681 cycles in the Turee Creek Group, Western Australia. *Geological Society of America*
682 *Bulletin*, **127**, 596–607.
- 683 58 Whitehouse, M.J., Nemchin, A.A., Pidgeon, R.T. (2017) What can Hadean detrital zircon
684 really tell us? A critical evaluation of their geochronology with implications for the
685 interpretation of oxygen and hafnium isotopes. *Gondwana Research*, **51**, 78-91.
- 686 59 Wilde, S.A., Valley, J.W., Peck, W.H., and Graham, C.M. (2001) Evidence from detrital
687 zircons for the existence of continental crust and oceans on the Earth 4.4 Gyr ago. *Nature*,
688 **409**, 175–178.
- 689 60 Young, G.M. (1970) An extensive early proterozoic glaciation in North America?
690 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **7**, 85–101.
- 691 61 Zeh, A., Gerdes, A., and Barton, J.M. (2009) Archean Accretion and Crustal Evolution of
692 the Kalahari Craton—the Zircon Age and Hf Isotope Record of Granitic Rocks from

693 Barberton/Swaziland to the Francistown Arc. *Journal of Petrology*, **50**, 933–966.

694 62 Zeh, A., Wilson, A.H., and Ovtcharova, M. (2016) Source and age of upper Transvaal
695 Supergroup, South Africa: Age-Hf isotope record of zircons in Magaliesberg quartzite and
696 Dullstroom lava, and implications for Paleoproterozoic (2.5–2.0Ga) continent
697 reconstruction. *Precambrian Research*, **278**, 1–21.

698

699 **Figure captions**

700 Figure 1: Localization of the Pilbara Craton in Western Australia (a). Geological maps of (b)
701 the Pilbara Craton and Hamersley Basin in Western Australia with Turee Creek Group
702 outcrops in yellow and (c) the Hardey Syncline area with locations of the TCDP drill cores
703 and surface samples (showed with numbers from 1 to 5) analyzed in this study (Modified
704 after Martin et al. (2000) and Van Kranendonk et al. (2015).

705

706 Figure 2: Stratigraphic logs of (a) the Mount Bruce Supergroup, which comprises the
707 Fortescue, Hamersley, and Turee Creek groups, and the overlying lower Wyloo Group with
708 geochronological constraints, (b) the complete section of Turee Creek Group in the Hardey
709 Syncline, (c) the Turee Creek Group section at the Boundary Ridge locality, (d) drill core
710 TCDP1, (e) drill core TCDP2, and (f) the upper Kazput cross section (see Fig. 1 b). (a)
711 Modified after Trendall *et al.* (2004) and Philippot *et al.* (2018), (b, d, e) after Philippot *et al.*
712 (2018) and (c, f) after Caquineau *et al.* (2018). Samples analyzed for their Hf isotope
713 compositions include from bottom to top: (1) Boolgeeda diamictite (T1-152, Pi-09.07, Pi-
714 09.08), (2) a sandstone layer of the Kungarra Formation (T2-332) and diamictites of the
715 Meteorite Bore Member (T2-272, T2-205, T2-169 and T2-130), (3) second Kungarra
716 diamictite MBM2 (Pi-15.41, Pi-15.42) (4) two massive sandstones of the upper Kazput
717 section (Pi-15.15, Pi-15.16, Pi-15.17) and (5) Beasley River Quartzite (Pi-15.02).

718 Abbreviations: WoD, Woolly Dolomite, CSB, Cheela Springs Basalt, BRQ, Beasley River
719 Quartzite, Ka, Kazput Formation, Kb, Koolbye Formation, MBM, Meteorite Bore Member,
720 Ku, Kungarra Formation, TCG, Turee Creek Group, Bo, Boolgeeda Iron Formation, WR,
721 Woongarra Rhyolite, WW, Weeli Wolli Formation, B IF, Brockman Iron Formation, MSW,
722 Mount Mac Rae Shale – Mt. Sylvia Formation – Wittenoon Formation, M IF, Marra Mamba
723 Iron Formation, Jee, Jeerinah Formation, Mad/Bu, Maddina Basalt/Bunjinah Formation,
724 Tu/Py, Tumbiana Formation/Pyradie Formation, Ky/Bo, Kylena Formation/Boongal
725 Formation, MRB, Mount Roe Basalt, Be, Bellary Formation. References for the
726 geochronological constraints are: (1) Müller *et al.* (2005), (2) Martin *et al.*(1998), (3)
727 Philippot *et al.* (2018), (4) Caquineau *et al.* (2018), (5) Trendall *et al.* (2004), (6) Arndt *et al.*
728 (1991).

729

730 Figure 3: $\epsilon_{Hf}(t)$ vs age diagrams for each sample: (a) Boolgeeda diamictite, (b) MBM
731 diamictite, (c) second Kungarra diamictite, (d) upper Kazput sandstones, (e) Beasley River
732 Quartzite, (f) all data. CHUR, Chondritic Uniform Reservoir, DM, Depleted Mantle. The DM
733 evolution line corresponds to a $^{176}\text{Lu}/^{177}\text{Hf}$ ratio of 0.03933 (Blichert-Toft and Puchtel, 2010)
734 evolving from the chondritic initial composition starting at 4567 Ma (Iizuka *et al.*, 2015).The
735 data can be confined between two model age lines at *c.a.* 2.54 Ga and 3.20 Ga. Hf model age
736 lines are represented for a $^{176}\text{Lu}/^{177}\text{Hf}$ ratio of 0.015 typical of bulk continental crust (Griffin
737 *et al.*, 2004).

738

739 Figure 4: A Hadean zircon in the Pilbara Craton. (a) U-Pb Concordia plot of zircon analyses
740 showing an upper intercept at 4013 ± 69 Ma and a Concordia date of 3998 ± 15 Ma. Inset:
741 cathodoluminescence image of the grain with laser spots indicated. (b) $\epsilon_{Hf}(t)$ vs. age
742 diagrams of the Hadean zircon with model age lines for two different natures of crust, a felsic

743 crust with a $^{176}\text{Lu}/^{177}\text{Hf}$ ratio of 0.015 (Griffin *et al.*, 2004), and a mafic crust with a
744 $^{176}\text{Lu}/^{177}\text{Hf}$ of 0.026 (Blichert-Toft and Albarède, 2008). The filtered domain of Jack Hills
745 zircon analyses after Whitehouse *et al.* (2017) is represented by the yellow box.

746

747 Figure 5: Comparison of $\varepsilon_{\text{Hf}}(t)$ data obtained in this study with literature data. The Boolgeeda
748 diamictite and Turee Creek Group data are represented as grey squares in all diagrams. (a)
749 Halfway Gneiss of the Glenburgh Terrane, (Johnson *et al.*, 2011), (b) Pilbara Craton, (Kemp
750 *et al.*, 2015; Gardiner *et al.*, 2017 and 2019; Petersson *et al.*, 2019), (c) The northern part of
751 the Yilgarn Craton, (Griffin *et al.*, 2004). (d,) South Africa: Magaliesberg Quartzite, (Zeh *et*
752 *al.*, 2016), Kalahari Craton (Zeh *et al.*, 2009). (e) Superior craton (Corfu and Noble, 1992;
753 Corfu and Stott, 1993 and 1996; Davis *et al.*, 2005).

754

755 **Tables Content**

756 **Table1.** Zircon Lu-Hf isotope data obtained by in *situ* Laser Ablation MC-ICP-MS.

757

758 **Table 2.** Zircon U-Th-Pb data from the Hadean zircon crystal obtained by in situ Laser
759 Ablation ICP-MS.

760

761 **Supplementary Table S1.** Detail of analytical conditions for Lu-Hf isotope analysis by *in*
762 *situ* Laser Ablation MC-ICP-MS. Lu-Hf isotope data on natural GJ-1, 91500 and Mud Tank
763 zircon reference materials and synthetic zircons doped with hafnium and rare earth elements
764 (Fischer *et al.*, 2011b) are included.