



**HAL**  
open science

# Effect of a 10-month residential multidisciplinary weight loss intervention on food reward in adolescents with obesity

Maud Miguet, Kristine Beaulieu, Alicia Fillon, Marwa Khammassi, Julie Masurier, Celine Lambert, Martine M. Duclos, Yves Boirie, Graham Finlayson, David Thivel

## ► To cite this version:

Maud Miguet, Kristine Beaulieu, Alicia Fillon, Marwa Khammassi, Julie Masurier, et al.. Effect of a 10-month residential multidisciplinary weight loss intervention on food reward in adolescents with obesity. *Physiology & behavior*, 2020, 223 (9), pp.112996. 10.1016/j.physbeh.2020.112996 . hal-03013128

**HAL Id: hal-03013128**

**<https://uca.hal.science/hal-03013128>**

Submitted on 25 Jun 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



UNIVERSITY OF LEEDS

This is a repository copy of *Effect of a 10-month residential multidisciplinary weight loss intervention on food reward in adolescents with obesity*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/168420/>

Version: Accepted Version

---

**Article:**

Miguet, M, Beaulieu, K [orcid.org/0000-0001-8926-6953](https://orcid.org/0000-0001-8926-6953), Fillon, A et al. (7 more authors) (2020) Effect of a 10-month residential multidisciplinary weight loss intervention on food reward in adolescents with obesity. *Physiology & Behavior*, 223. 112996. ISSN 0031-9384

<https://doi.org/10.1016/j.physbeh.2020.112996>

---

© 2020, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

**Weight loss induced by a 10-month residential multidisciplinary intervention is associated with a decrease in food reward in adolescents with obesity**

Journal:	<i>British Journal of Nutrition</i>
Manuscript ID	Draft
Manuscript Type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Miguet, Maud; Clemmont Auvergne University, AME2 Beaulieu, Kristine; University of Leeds, School of Psychology Fillon, Alicia; Clermont Auvergne University - Cezeaux Campus, ame2p Khammassi, Marwa; Clermont-Ferrand I University, ame2p Julie, MASURIER; UGECAM, AUVERGNE RHONE ALPES Céline, LAMBERT; DRCI, Auvergnés Rhone Alpes Duclos, Martine; CHU, Sports medicine Borie, Yves; INRA UMR1019: Unit of Human Nutrition , CRNH Auvergne; Centre Hospitalier Universitaire de Clermont-Ferrand, Service de Nutrition Clinique Finlayson, Graham; University of Leeds, School of Psychology Thivel, David; University of Clermont, AME2P
Keywords:	Weight Loss, Multidisciplinary Intervention, Energy Intake, Food Reward
Subject Category:	Behaviour, Appetite and Obesity

SCHOLARONE™  
Manuscripts

## Title page

---

**Title:**

Weight loss induced by a 10-month residential multidisciplinary intervention is associated with a decrease in food reward in adolescents with obesity

**Authors:**

Maud Miguet<sup>1</sup>, Kristine Beaulieu<sup>2</sup>, Alicia Fillon<sup>1</sup>, Marwa Khammassi<sup>1</sup>, Julie Masurier<sup>3</sup>, Céline Lambert<sup>4</sup>, Martine Duclos<sup>5</sup>, Yves Boirie<sup>6</sup>, Graham Finlayson<sup>2</sup>, David Thivel<sup>1</sup>

**MIGUET****Affiliations:**

1: Clermont Auvergne University, EA 3533, Laboratory of the Metabolic Adaptations to Exercise under Physiological and Pathological Conditions (AME2P), Clermont-Ferrand, France.

2: School of Psychology, Faculty of Medicine and Health, University of Leeds, Leeds, LS2 9JT, UK

3: UGECAM Nutrition Obesity Ambulatory Hospital, Clermont-Ferrand, France

4: Clermont-Ferrand University hospital, Biostatistics unit (DRCI), Clermont-Ferrand, France

5: Department of Sport Medicine and Functional Explorations, Clermont-Ferrand University Hospital, G. Montpied Hospital, Clermont-Ferrand, France.

6: Department of Human Nutrition, Clermont-Ferrand University Hospital, G. Montpied Hospital, Clermont-Ferrand, France

**Corresponding author:**

Maud Miguet

[maud.miguet@gmail.com](mailto:maud.miguet@gmail.com)

Laboratory of Metabolic Adaptations to Exercise under Physiological and Pathological conditions, EA3533, Blaise Pascal University, Clermont University, France

Phone. 0473407679 / Fax. 0473407679

**Running head:**

Food reward and weight loss in adolescents

**Keywords:**

Weight Loss, Multidisciplinary Intervention, Energy Intake, Food Reward

**Title:**

2 **Weight loss induced by a 10-month residential multidisciplinary intervention is**  
3 **associated with a decrease in food reward in adolescents with obesity**

4

**Authors:**

6 Maud Miguet<sup>1</sup>, Kristine Beaulieu<sup>2</sup>, Alicia Fillon<sup>1</sup>, Marwa Khammassi<sup>1</sup>, Julie Masurier<sup>3</sup>,  
7 Céline Lambert<sup>4</sup>, Martine Duclos<sup>5</sup>, Yves Boirie<sup>6</sup>, Graham Finlayson<sup>2</sup>, David Thivel<sup>1</sup>

8

**Affiliations:**

10 1: Clermont Auvergne University, EA 3533, Laboratory of the Metabolic Adaptations to Exercise  
under Physiological and Pathological Conditions (AME2P), Clermont-Ferrand, France.

12 2: School of Psychology, Faculty of Medicine and Health, University of Leeds, Leeds, LS2 9JT, UK

3: UGECAM Nutrition Obesity Ambulatory Hostipal, Clermont-Ferrand, France

14 4: Clermont-Ferrand University hospital, Biostatistics unit (DRCI), Clermont-Ferrand, France

5: Department of Sport Medicine and Functional Explorations, Clermont-Ferrand University Hospital,  
16 G. Montpied Hospital, Clermont-Ferrand, France.

6: Department of Human Nutrition, Clermont-Ferrand University Hospital, G. Montpied Hospital,  
18 Clermont-Ferrand, France

**Corresponding author:**

Maud Miguet

22 [maud.miguet@gmail.com](mailto:maud.miguet@gmail.com)

Laboratory of Metabolic Adaptations to Exercise under Physiological and Pathological  
24 conditions, EA3533, Blaise Pascal University, Clermont University, France

Phone. 0473407679 / Fax. 0473407679

26

**Running head:**

28 Food reward and weight loss in adolescents

**Keywords:**

Weight Loss, Multidisciplinary Intervention, Energy Intake, Food Reward

32

**Conflicts of interest statement:** The authors declare no conflicts of interest.

## 34 1. Abstract

36 **Background.** While multidisciplinary weight loss programs - including physical activity - have  
38 been suggested to improve eating disorders, emotional eating, and the sensitivity of the appetite  
control system; this study examined the effect of a specific multidisciplinary intervention on  
appetite control and food reward in adolescents with obesity.

40 **Methods.** Thirty adolescents with obesity (11-15 years) took part in a 10-month inpatient  
multidisciplinary weight loss program. Body composition, energy intake, appetite, food reward  
42 and eating behavior traits were measured at baseline, 5 months and the end of the 10-month  
intervention.

44 **Results.** *Ad libitum* intake decreased after the 10-month intervention ( $p < 0.001$ ). Fasting hunger  
( $p = 0.02$ ) and desire to eat ( $p = 0.01$ ), daily hunger ( $p = 0.001$ ) and pre-meal liking for high-fat  
46 savoury foods (HFSA;  $p = 0.03$ ), low-fat savoury foods (LFSA;  $p = 0.04$ ), high-fat sweet foods  
(HFSW;  $p = 0.009$ ), low-fat sweet foods (LFSW;  $p = 0.005$ ) increased after 5 months. Fasting and  
48 daily hunger ( $p = 0.02$  and  $p = 0.008$ , respectively), desire to eat ( $p < 0.001$  and  $p = 0.005$ ,  
respectively), prospective food consumption (PFC;  $p = 0.048$  and  $p = 0.03$ , respectively); pre-  
50 meal liking for HFSA ( $p = 0.007$ ), LFSA ( $p < 0.001$ ), HFSW ( $p = 0.02$ ), LFSW ( $p < 0.001$ );  
emotional eating ( $p < 0.001$ ) and uncontrolled eating ( $p = 0.009$ ), decreased between months 5  
52 and 10. Post-meal liking for HFSA ( $p < 0.001$ ), LFSA ( $p = 0.002$ ), HFSW ( $p = 0.02$ ) and LFSW  
( $p < 0.001$ ) decreased between baseline and month 5 and remained unchanged between months  
54 5 and 10.

**Conclusion.** These results indicate possible improvements in the reward response to food in  
56 adolescents with obesity and may contribute to the beneficial effect of a multicomponent weight  
loss intervention in this population. These findings suggest that adaptive mechanisms to weight  
58 loss occurring in the short-to-medium term are attenuated in the longer term with the persistence  
of weight loss.

60

**Key words.** Weight Loss, Multidisciplinary Intervention, Energy Intake, Food Reward

62

## 2. Introduction

64

In the current obesogenic environment, the interaction between the homeostatic and hedonic  
66 appetite systems in the control of energy intake is of considerable interest (1). The increasing  
availability of highly palatable food is likely to favor overeating, independently of dietary status  
68 (2).

Adolescents are particularly vulnerable to increased consumption of unhealthy foods and rising  
70 obesity rates (3), with one out of five children having obesity in Europe (4). During adolescence,  
consumption of 'junk food' or other high-fat foods is related to pleasure, social influence from  
72 peers and independence from parental control (5). In that regard, the relationship between food  
reward and obesity has been previously demonstrated, with higher relative reinforcing value of  
74 food predicting greater weight gain (6), while lower food reinforcement is associated with better  
weight loss (7). Since childhood is an important period to enhance food acceptance and induce  
76 a wider dietary range, research is needed to identify strategies to make children's food  
preferences and food choices healthier during this critical time. This is especially important  
78 because 80% of adolescents with obesity are at high risk of becoming adults with obesity (8).

Recently, physical activity has been described as a good strategy to decrease eating disorders,  
80 emotional eating and energy intake in adults and adolescents with obesity (9–12) as well as  
improving the sensitivity of the appetite control system in adults (13). Moreover, a recent  
82 systematic review revealed that liking and wanting for high-energy dense food decreased  
following weight management interventions in adults (14), contrary to the notion that adaptive  
84 mechanisms may increase appetite and food reward to counteract weight loss. Such results  
suggest that the effect of physical activity and weight loss interventions on overall health, body  
86 weight and body composition might be in part mediated by modifications in eating behaviors.  
Physiological pathways have been identified to explain some of these nutritional responses to  
88 exercise (15); however, the neurocognitive factors remain less explored. Specifically, food  
reward appears to be an important factor likely to be implicated in the success of weight loss  
90 programs, but this remains to be investigated.

Currently, weight management interventions tend to rest on multidisciplinary approaches  
92 involving nutritional counselling, psychological support and exercise (16). Such interventions  
have been shown effective (at least in the short term), leading to significant improvements in  
94 body weight and body composition (17), physical fitness (17), cardio-metabolic profile (18),

and health-related quality of life (19), among others. However, their effects on eating behaviors are mixed. Although a recent meta-analysis found reduced daily energy intake in response to multicomponent weight loss programs in adolescents with obesity (10), it also noted that most studies assessed energy intake as a secondary outcome using self-reported food diaries, which have been shown to favor underestimation of food consumption in this population (20). The authors therefore expressly identified the need for additional studies using alternative dietary assessment strategies with less potential bias (10). According to some recent results from our research group, adolescents with obesity significantly increased their weighed daily *ad libitum* energy intake in response to multidisciplinary weight loss interventions (21,22). Importantly, our work also highlighted the need to consider the adolescents' individual eating behavior traits, given that the most cognitively restrained eaters appeared to be the most vulnerable to increased energy intake in tempting situations (23), such as *ad libitum* buffet meals (21,22). The potential underlying mechanisms remain to be elucidated, particularly the potential effect of such weight loss intervention on the hedonic and cognitive control of food consumption.

The primary aim of this study was to investigate the homeostatic (energy and macronutrient intake and appetite sensations) and food reward (explicit liking and implicit wanting) responses to a 10-month multidisciplinary weight loss intervention among youths with obesity. The secondary aims were to examine the effects of the intervention on cognitive control over eating (eating behaviour traits).

### 114 3. Methods

#### 3.1. Participants

116 A total of 30 adolescents (mean age:  $13.1 \pm 1.0$  years) with obesity as defined by Cole et al. (24) (mean BMI z-score:  $2.3 \pm 0.2$ ; BMI percentile:  $98.7 \pm 0.5$ ) were enrolled in this study (7 boys and 23 girls) through local pediatric consultations. Convenience sampling was used in this study. To be included, the adolescents had to: i) be aged between 11 and 15 years; ii) present a BMI equal or above the 95th percentile for their gender and age (24); iii) not take any medication that could interact with the protocol (e.g., thyroid medication, stimulant medication, medication for diabetes); iv) present no contraindication to physical activity; and v) take part in less than 2 hours of physical activity per week (according to the International Physical Activity Questionnaire – IPAQ). All the adolescents and their legal representative received information sheets and signed up consent forms as requested by the ethical authorities.



### 126 3.2. Protocol design

127 After a medical screening conducted by a pediatrician to ensure their ability to complete the  
128 study, the adolescents were enrolled in a 10-month residential multidisciplinary weight loss  
129 program in a local Pediatric Obesity Center (Tza Nou, La Bourboule, France). Maturation status  
130 was assessed by the physician using the Tanner stages during the first medical visit.  
131 Anthropometric measurements, body composition (dual-energy X-ray absorptiometry), daily  
132 energy intake (*ad libitum* buffet meals), food reward (Leeds Food Questionnaire Preference),  
133 eating behavior traits (Three-Factor Eating Questionnaire) and appetite sensations (visual  
134 analogue scales) were assessed before (T0), after 5 months (T1), and at the end of the 10-month  
135 program (T2). This study was conducted in accordance with the Helsinki declaration and  
136 received ethical approval (CPP Sud Est VI: 2015-33; Clinical Trial NCT02626273).

### 3.3. Measurements

138 The following measurements were performed at baseline (T0), 5 months (T1) and the end of  
the 10-month program (T2).

#### 140 **Anthropometric characteristics and body composition**

141 Body weight and height were recorded to the nearest 0.1 kg and 0.5 cm, respectively, while  
142 wearing light clothes and bare-footed, using a digital scale and a standard wall-mounted  
stadiometer, respectively. BMI was calculated as weight (kg) divided by height squared (m<sup>2</sup>).  
144 Afterwards, BMI was reported in the sex and age dependent French reference curves to get the  
BMI percentile. Fat mass (FM) and fat-free mass (FFM) were assessed, under fasted state, by  
146 dual-energy X-ray absorptiometry (DXA) following standardized procedures (QDR4500A  
scanner, Hologic, Waltham, MA, USA).

#### 148 **Eating behavior traits**

149 The Three-Factor Eating Questionnaire (TFEQ R-21; (25)) was used to assess eating behavior  
150 traits. Three domains were evaluated: uncontrolled eating (UE: tendency to overeating in  
response to loss of control over intake); emotional eating (EE: overeating in response to  
152 negative moods) and cognitive restraint (CR: individuals' efforts to limit their food intake to  
control body weight or to promote weight loss). Participants completed the 21 items in a four-  
154 point Likert scale for items 1–20 and on an eight-point numerical rating scale for item 21. Their  
answers were coded following the instructions given by Cappelleri et al. (25). Items 1 to 16  
156 were reverse coded and item 21 was recoded as follows: 1–2 scores as 1; 3–4 as 2; 5–6 as 3;

7–8 as 4. Means for each domain were then calculated (domain score also range from 1 to 4),  
158 with higher score indicate more UE, EE and CR.

### ***Ad libitum* energy and macronutrient intake**

160 Daily *ad libitum* energy intake was assessed during three single test days performed at T0, T1  
and T2. At 08:00, after an overnight fast, the adolescents consumed a standardized calibrated  
162 breakfast respecting the recommendations for their age ( $\approx 500$  kcal, (26). Lunch (12 p.m.) and  
dinner (6:30 p.m.) meals were *ad libitum* served using a buffet-type meal. The content of the  
164 buffets was determined based on the adolescent's food preferences and eating habits. Top rated  
items as well as disliked ones and items liked but not usually consumed were excluded to avoid  
166 over-, under- and occasional/opportunistic consumption. At lunch, the menu was composed of  
beef steaks, pasta, mustard, cheese, yogurt, apple sauce, fruits and bread. Dinner menu was  
168 composed of ham/turkey, beans, mashed potatoes, cheese, yogurt, apple sauce, fruits and bread.  
Food items were presented in abundance and accompanied with water (1-L carafe) only.  
170 Adolescents made their choices and composed their trays individually before joining their  
habitual table (5 adolescents per table). Adolescents were told to eat until feeling comfortably  
172 satiated and had access to extra food if wanted. Food items were weighed by the experimenters  
before and after consumption, and the macronutrient distribution (proportion of fat,  
174 carbohydrate and protein) and total energy intake in kcal were calculated using the software  
Bilnut 4.0 (4.0 SCDA, Nutrisoft, Software, France). Total daily energy intake was calculated  
176 by summing breakfast, lunch and dinner meals. This methodology has been previously  
validated and published (27). Between the test meals, the adolescents did not have access to  
178 any food but were free to drink water *ad libitum*, and were requested not to engage in any  
moderate to vigorous physical activity and mainly performed sedentary activities such as  
180 reading, homework, or board games.

### **Subjective appetite sensations**

182 Appetite sensations were collected throughout the test day using visual analogue scales (150-  
mm scales). Adolescents reported hunger, fullness, desire to eat and prospective food  
184 consumption at six regulated times: before and immediately after the breakfast, lunch, and  
dinner. The questions were: i) "How hungry do you feel?", ii) "How full do you feel?", iii)  
186 "Would you like to eat something?", iv) "How much do you think you can eat?" (adolescents  
were asked to respond on a scale from "not at all" to "a lot"). This method has been previously  
188 validated (28). Area under the curve (AUC) was calculated based on the trapezoid method for  
daily appetite sensations.

### 190 **Food liking and wanting**

192 The reward value of food was assessed pre-lunch (hungry state) and post-lunch (fed state) during the test days. The Leeds Food Preference Questionnaire (described in greater methodological detail by Dalton and Finlayson (29) provided measures of food preference and food reward. Participants were presented with an array of pictures of individual food items common in the diet. Foods in the array were chosen by the local research team from a validated database to be either predominantly high (>50% energy) or low (<20% energy) in fat but similar in familiarity, protein content, palatability and suitable for the study population. Explicit liking was measured by participants rating the extent to which they like each food (“How pleasant would it be to taste this food now?”). The food images were presented individually, in a randomized order and participants make their ratings using a 100-mm VAS. Implicit wanting was assessed using a forced choice methodology in which the food images were paired so that every image from each of the four food types was compared to every other type over 96 trials (food pairs). Participants were instructed to respond as quickly and accurately as they could to indicate the food they want to eat the most at that time (“Which food do you most want to eat now?”). To measure implicit wanting, reaction times for all responses were covertly recorded and used to compute mean response times for each food type after adjusting for frequency of selection. Responses on the LFPQ were used to compute mean scores for high fat, low fat, sweet or savoury food types (and different fat-taste combinations).

#### 3.4. *Multidisciplinary weight loss program*

210 The 10-month residential multidisciplinary weight loss program combined physical activity, nutritional education and psychological support. The physical activity intervention was composed of four 60-minute physical activity sessions per week including aerobic training, strength training, aquatic activities and leisure-time activities (e.g. soccer). Concomitantly, the adolescents attended 2 hours of physical education per week at school. The adolescents also attended nutritional education classes twice a month led by a dietician and received psychological support through individualized consultations with a professional once a month. During the intervention, the adolescents were prescribed a balanced diet, based on their age and sex recommendations for energy requirements in accordance with the national nutrition guidelines (30). Adolescents were not subjected to energy restriction. The focus of the nutritional intervention was on the regularity of the meal, the variety and the quality of the food (31).

### 222 3.5. Statistical analyses

223 Statistical analysis was performed using Stata software (version 13; StataCorp, College Station,  
224 Texas, USA). All tests were two-sided, with a Type I error set at 0.05. As discussed by Feise in  
2002, the adjustment of Type I error were not proposed systematically but according to clinical  
226 and not only statistical considerations (32). Continuous data were expressed as mean and  
standard deviation. To measure the effect of the 10-month intervention over time (T0, T1, and  
228 T2) on weight, BMI, FM, FFM, EI, macronutrients, appetite sensations, eating behaviors  
scores, statistical analyses were conducted with linear mixed models to take into account the  
230 repeated measurements per subject (time as fixed effect and subject as random effect). Paired  
t-tests were used to compare changes during phase 1 and phase 2. To measure the meal effect  
232 (before/after meal) on food reward, linear mixed models were also considered at each time,  
with meal (before/after meal) as fixed effect and subject as random effect. Interaction between  
234 time (T0, T1, and T2) and meal (before/after meal) was tested before performing subgroup  
analyses.

## 236 4. Results

### 238 Participant characteristics

Among the initial 30 children included, 24 completed the 10-month intervention (80%  
240 retention). Of the six participants who left the trial, two were excluded for disciplinary reasons,  
three discontinued the intervention for family reasons and one for school-related reasons. None  
242 of the dropouts was related to the study protocol.

Anthropometric and body composition changes are presented in Table 1. As expected, body  
244 weight, body mass index and percent fat mass decreased significantly throughout the 10-month  
weight loss intervention ( $p < 0.001$ ). There was also a slight but significant reduction in fat-free  
246 mass after the intervention.

Weight and fat mass loss were greater during the first phase. Between baseline and mid-  
248 intervention adolescents lost on average  $5.7 \pm 4.3$ g body weight and  $4.3 \pm 1.9$  % of fat mass  
*versus*  $4.4 \pm 3.3$ g and  $2.1 \pm 2.2$  % during the second phase of the intervention ( $p < 0.0001$ ).

250

**Energy intake, dietary profile and appetite sensations**

252 As shown in Table 2, there was a significant time effect for lunch *ad libitum* energy intake  
( $p < 0.001$ ) with a decrease of 198 kcal between T0 and T2 ( $p = 0.006$ ) and of 282 kcal between  
254 T1 and T2 ( $p < 0.001$ ). There was a non-significant increase in dinner (58 kcal) and decrease in  
total (139 kcal) *ad libitum* energy intake from T0 to T2.

256 The proportion of fat consumed decreased over time ( $p < 0.001$ , with significant decreases  
between T0 and T2 and between T1 and T2  $p < 0.001$ ), whereas energy intake derived from  
258 carbohydrate increased ( $p < 0.001$ , with significant increases between T0 and T2 and between  
T1 and T2  $p < 0.001$ ).

260 Regarding eating behaviors traits, uncontrolled and emotional eating decreased over time  
( $p < 0.001$  between T0 and T2), whereas restrained eating remained stable.

262 As shown in Figure 1, fasting hunger and fasting desire to eat both increased between T0 and  
T1 ( $p = 0.02$  and  $p = 0.01$ , respectively) and decreased between T1 and T2 ( $p = 0.02$  and  $p < 0.001$ ,  
264 respectively). Fasting prospective food consumption decreased between T1 and T2 ( $p = 0.048$ ).  
There was a time effect for daily hunger and desire to eat ( $p = 0.002$  and  $p = 0.02$ , respectively)  
266 with daily hunger increasing between T0 and T1 ( $p = 0.001$ ) and decreasing between T1 and T2  
( $p = 0.008$ ), and daily desire to eat decreasing between T1 and T2 ( $p = 0.005$ ). Daily prospective  
268 food consumption decreased between T1 and T2 ( $p = 0.03$ ).

**Food reward**

270 Regarding food reward (Table 3), main changes were found for explicit liking. We observed a  
272 time effect in the hungry state (pre-meal) for explicit liking for all food categories, which all  
increased between T0 and T1, then decreased between T1 and T2 (see Table 3 for details). In  
274 the fed state (post-meal), values in explicit liking for HFSA, LFSA and LFSW followed a  
different trend; they decreased from T0 to T1 then remained stable at T2. The interactions  
276 between time and meal revealed a different effect of the meal on explicit liking over time.  
Mainly, explicit liking decreased significantly from hungry to fed for each food category at T1  
278 and T2 whereas at T0 there was no meal effect. In terms of implicit wanting, there were no  
meal, time or interaction effects at any time point.

280 Figure 2 illustrates the variability in food reward changes (hungry state) between the first phase  
(T0 to T1) and the second phase (T1 to T2) of the program. As highlighted, there was a clear

282 trend for an increase in explicit liking during the first phase and a decrease in explicit liking  
during the second phase for all food categories. On the other hand, the variability in the change  
284 in implicit wanting was large and appeared to be evenly distributed across those that increased  
and those that decreased, resulting in little change in the overall mean.

286

## 5. Discussion

288

The present study suggests non-linear temporal changes in appetite and food reward during the  
290 10-month multidisciplinary weight loss intervention in adolescents with obesity. Interestingly,  
our results suggest an increase in appetite and liking for all food categories mid-intervention,  
292 followed by a decrease at the end. Thus, the homeostatic and hedonic appetite adaptations to a  
multidisciplinary program might depend on the degree and timing of the weight loss, with  
294 opposite responses being observed between the first 5 months of intervention (phase 1) and the  
last 5 months (phase 2). Finally, after 5 months and until the end of the intervention, food  
296 consumption (*ad libitum* lunch meal) induced a decrease in explicit liking for all food  
categories, suggesting a reconnection between the hedonic and homeostatic systems of appetite  
298 control. Overall, this reduction in food reward was accompanied by a reduction in *ad libitum*  
energy intake at lunch and a slight reduction in daily energy intake.

300 In developed countries, the obesogenic environment – promoting a high-energy intake and a  
sedentary lifestyle – favors an hedonic rather than homeostatic control of food intake (1). This  
302 shift toward an hedonic over an homeostatic control of eating behavior has been linked with  
reduced sensitivity to internal cues for satiety (33), leading to overeating. In the long term, the  
304 resulting positive energy balance is responsible for weight gain and then the development of  
overweight and obesity. In that context, obesity management has been subject to much research,  
306 aiming at improving both its prevention and treatment. Previous studies have shown that  
multifaceted approaches are more successful, leading to body composition, cardiovascular,  
308 metabolic and well-being improvements, compared with nutritional interventions alone (34).  
Moreover, physical activity should be considered as an important component in weight  
310 management programs, especially since it is now recognized to affect not only energy  
expenditure, but also appetite control and energy intake (35). While the first study examining  
312 the relationship between exercise and food intake in children was conducted 70 years ago (36),

many others have been conducted since then demonstrating the beneficial effect of exercise on  
314 the control of food intake (for reviews see (15,37). While the homeostatic mechanisms linking  
exercise and food intake have been largely investigated lately, the neurocognitive pathways  
316 remain less explored, especially among youth. Identifying interventions that modulate the  
hedonic aspects of food intake in adolescents is of major interest given the current tendencies  
318 of the rates of obesity.

In that context, this study investigated for the first time the effect of a multidisciplinary weight  
320 loss intervention on food reward in adolescents with obesity. According to our results, distinct  
changes with the intervention occurred over the two phases. During the first 5 months of the  
322 intervention, we found an increase in explicit liking in the hungry state for all food categories.  
In a similar manner, subjective appetite sensations of hunger and desire to eat both increased  
324 during the first 5 months of the intervention. Although it did not reach significance, lunch  
energy intake and daily energy intake also slightly increased during this phase. Together these  
326 results suggest, in line with previous theories, the presence compensatory adaptations in  
appetite in the first phase of weight loss (38,39). The compensatory mechanisms observed  
328 during the first phase of the intervention in the current study are consistent with recently  
published results showing an increase in the subjective sensation of hunger (40) and energy  
330 intake (21,22) after a multicomponent weight loss intervention in adolescents with obesity. A  
recent meta-analysis from Doucet et al. (39) synthesized the effect of weight loss on energy  
332 balance. According to them, there are both homeostatic and hedonic adaptations leading to an  
increase in appetite and drive to eat after weight loss. An attenuation of the anorexigenic  
334 peripheral signals (leptin, PYY, GLP-1) in response to weight loss may explain compensatory  
increases in food intake through weakened satiety (41). On the other hand, weight loss via  
336 energy restriction has been previously associated with increases in food reward in adults with  
obesity (42). However, if very few studies take mid-point measures, they may miss important  
338 temporal information occurring throughout an intervention. Indeed, we observed a different  
trend during the second phase of the program, leading to a decrease in explicit liking for all  
340 food categories, hunger, desire to eat, prospective food consumption and *ad libitum* lunch  
energy intake. These observations are in line with previous studies highlighting a reduced  
342 difficulty to maintain a stable body weight after weight reduction (43). Klem et al. reported that  
the effort and the attention allocated to weight maintenance decreased as the duration of weight  
344 maintenance increased. In other words, the “cost” related to weight maintenance after weight  
loss appears to decrease over time. Importantly, during the second phase, weight and fat mass



346 loss were lower, compared with the first phase of the intervention, suggesting that the degree  
of weight loss could be implicated in the compensatory mechanisms observed, with more rapid  
348 weight loss being counterproductive. It appears indeed that compensatory mechanisms are  
decreased in response to the slower weight loss period. These assumption need to be further  
350 investigated.

Moreover, while there were no changes in food reward in response to the meal at baseline, from  
352 the fifth month, the meal induced a significant decrease in food reward. As the reduction in  
food reward is typically observed from the hungry to fed states (44,45), this lack of effect at  
354 baseline suggests that the reward system wasn't sensitive to homeostatic satiety signals at this  
time. This assumption is in line with a previous study from Stice at al. demonstrating that  
356 adolescents with obesity experienced less satisfaction from eating food compared with  
adolescents who are lean, due to a reduced response in the brain reward circuit (46). Advanced  
358 theories support both a hyper-responsiveness of reward circuitry in the fasted state, and a hypo-  
responsiveness in the fed state in people with obesity, both increasing the risk for overeating  
360 (47). Consequently, identifying interventions that can modulate these reward "abnormalities"  
is of high relevance to treat and prevent obesity. In a recent systematic review, Oustric at al.  
362 report a decrease in food reward after weight loss in adults with obesity (all interventions  
combined: dietary, pharmacological, behavioral and cognitive), and hypothesized that short  
364 term food deprivation may enhance food reward, whereas longer term deprivation might  
attenuate it (Oustric et al., 2018), which is in line with the present results.

366 The reduction of lunch *ad libitum* energy intake observed at 10-month is also in line with the  
systematic review by Schwartz et al. (10) who found that physical activity interventions in  
368 adolescents with obesity induced a decrease in daily energy intake. The reduction in daily  
energy intake in the current study was smaller in magnitude. Furthermore, we recorded an  
370 overall shift in macronutrient intake towards less fat and more carbohydrate, which couldn't be  
analyzed in the review from Schwartz et al. due to the lack of evidence regarding the relative  
372 contribution of each macronutrient (in percentage) to total energy intake. Further specific  
studies are therefore needed. One of the strength of the present study is the robust methodology  
374 we used to asses both energy intake (considering the relative contribution of fat, carbohydrate  
and protein) and food reward (considering liking and wanting). Indeed, as highlighted by  
376 Schwartz et al., the homeostatic appetite adaptations to weight loss are often considered as  
secondary outcomes and food intake is commonly assessed using dietary recall which is a  
378 highly subjective method, especially in this specific population. Instead of using a self-reported



380 methods, we measured food intake via *ad libitum* buffet meals, which is most likely to reflect  
objective changes in food intake. Moreover, we used a psychometric methodology to assess  
382 food reward (the LFPQ), which provides a better reflection of the behavioral hedonic responses,  
compared with other methodologies using for example fMRI or BOLD signal. Finally, since it  
384 appears essential to better understand the control of energy intake throughout weight loss, we  
considered eating behavior traits of the participants as a variable of interest. Regarding these  
behavioural traits, we found coherent and encouraging results, demonstrating a significant  
386 decrease in emotional and uncontrolled eating, while restrained eating remained stable across  
time. As aforementioned, in the present environment favoring disinhibited eating and loss of  
388 control over eating, it is a priority to target the control of food intake and to reduce eating  
behaviors related with weight gain (e.g. emotional, external, uncontrolled and restrained eating)  
390 during weight loss interventions.

Interestingly, there were no overall changes in implicit wanting in response to the intervention  
392 and only mean liking was altered. A similar effect of changes in liking but not wanting in  
response to diet-induced weight loss was observed by Oustric et al. (48). In their study, adult  
394 women were subjected to continuous or intermittent energy restriction to 5% weight loss within  
12 weeks. In contrast, Beaulieu et al. report a reduction in wanting but not liking after a 12-  
396 week exercise training intervention in adults with overweight and obesity (49). Together these  
results suggest potential differences in reward-related mechanisms affected by diet- and  
398 exercise-induced weight loss. This warrants further investigations, especially since some  
evidence suggests that wanting may be more important than liking in weight management (50).  
400 Yet, there appeared to be more inter-individual variability in implicit wanting than in explicit  
liking. Almost as many adolescents had increases in implicit wanting as adolescents who had  
402 decreases, resulting in no overall mean differences. Exploratory analyses revealed no  
differences in outcomes between those that increased or decreased wanting (data not shown).  
404 Understanding what leads individuals to increase or decrease implicit wanting during weight  
loss, and the implications for weight management, is warranted.

406 The present results must be considered and interpreted in light of some limitations. First, the  
relatively small sample size did not allow us to further examine the inter-individual variability  
408 in the homeostatic adaptations to weight loss, which seems interesting in view of our results.  
Moreover, it would have been relevant to conduct a follow-up assessment to analyze whether  
410 the observed changes were maintained over time after the intervention. Finally, it would have  
been valuable to measure food reward using a French version of the LFPQ; however, at the

412 time of our study, the French version had not yet been validated (the validation process is  
currently ongoing).

## 414 **6. Conclusion**

416 The present work is the first to evaluate the impact of weight loss on food reward in adolescents  
with obesity and provides new insights regarding the effect of a 10-month multidisciplinary  
418 intervention on both the control of food intake and appetite in this population. We show that  
mean liking in the hungry state increased during the first phase of the multicomponent weight  
420 loss program and returned to baseline after the completion of the intervention whereas liking in  
the fed state decreased during the first phase and remained unchanged in the second phase. No  
422 changes were observed for wanting. Concurrently, appetite sensations and *ad libitum* energy  
intake increased during the first 5 months, followed by a reduction, suggesting that adaptive  
424 mechanisms to weight loss occurring in the short-to-medium term can be attenuated with  
persistence of weight loss in the longer term. The implications are that clinicians and patients  
426 need to be careful with potential compensatory adaptations occurring in the earlier phases of a  
weight loss intervention as this is a risky time for attrition and loss of motivation. Finally, while  
428 these results seem to reinforce the beneficial effect of a multicomponent weight loss  
intervention including exercise on the hedonic appetite system controlling food intake in  
430 adolescents with obesity, further work is required to understand the individual variability in the  
adaptations in appetite in response to weight loss.

432

## **7. Acknowledgements**

434 The authors are grateful to all of the adolescents that participated in the program, and to the  
Nutrition Obesity Ambulatory Hospital (UGECAM) that provided their generous support. We  
436 also want to thank the Auvergne Regional Council for its help through its 2016 New Researcher  
Award.

## 438 **8. Funding**

This research did not receive any specific grant from funding agencies in the public,  
440 commercial, or not-for-profit sectors.

## 9. Declarations of interest

442 The authors have not conflict of interest to disclose.

## 10. References

- 444 1. Berthoud H-R. Metabolic and hedonic drives in the neural control of appetite: who is the boss? *Curr Opin Neurobiol.* déc 2011;21(6):888-96.
- 446 2. Davis C. From Passive Overeating to “Food Addiction”: A Spectrum of Compulsion and Severity. *ISRN Obes* [Internet]. 15 mai 2013 [cité 31 juill 2019];2013. Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3901973/>
- 448
3. Han JC, Lawlor DA, Kimm SY. Childhood obesity. *The Lancet.* 15 mai 2010;375(9727):1737-48.
- 450
4. WHO European Childhood Obesity Surveillance Initiative. 9th Meeting of the WHO European Childhood Obesity Surveillance Initiative (COSI). Meeting Report (2016) [Internet]. 2017 [cité 4 avr 2018]. Disponible sur: <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/publications/meeting-reports/technical-meetings-and-workshops/9th-meeting-of-the-who-european-childhood-obesity-surveillance-initiative-cosi.-meeting-report-2016>
- 452
- 454
- 456
5. Chapman G, Maclean H. “Junk food” and “healthy food”: meanings of food in adolescent women’s culture. *Journal of Nutrition Education.* 1 mai 1993;25(3):108-13.
- 458
6. Epstein LH, Yokum S, Feda DM, Stice E. Food reinforcement and parental obesity predict future weight gain in non-obese adolescents. *Appetite.* nov 2014;82:138-42.
- 460
7. Buscemi J, Murphy JG, Berlin KS, Raynor HA. A behavioral economic analysis of changes in food-related and food-free reinforcement during weight loss treatment. *J Consult Clin Psychol.* août 2014;82(4):659-69.
- 462
8. Guo N, Chumlea N. Tracking of body mass index in children in relation to overweight in adulthood. *Am J Clin Nutr.* juill 1999;70(1 Part 2):145S-148S.
- 464
9. Blanchet C, Mathieu M-È, St-Laurent A, Fecteau S, St-Amour N, Drapeau V. A Systematic Review of Physical Activity Interventions in Individuals with Binge Eating Disorders. *Curr Obes Rep.* mars 2018;7(1):76-88.
- 466
- 468
10. Schwartz C, King NA, Perreira B, Blundell JE, Thivel D. A systematic review and meta-analysis of energy and macronutrient intake responses to physical activity interventions in children and adolescents with obesity: Food Intake and exercise in obese youth. *Pediatric Obesity.* juin 2017;12(3):179-94.
- 470
- 472
11. Miguet M, Fillon A, Khammassi M, Masurier J, Julian V, Pereira B, et al. Appetite, energy intake and food reward responses to an acute High Intensity Interval Exercise in adolescents with obesity. *Physiol Behav.* 15 oct 2018;195:90-7.
- 474

- 476 12. Martín-García M, Alegre Durán LM, García-Cuartero B, Bryant EJ, Gutin B, Royo IA. Effects  
478 of a 3-month vigorous physical activity intervention on eating behaviors and body  
composition in overweight and obese boys and girls. *Journal of Sport and Health Science*  
[Internet]. sept 2017 [cité 19 déc 2017]; Disponible sur:  
480 <http://linkinghub.elsevier.com/retrieve/pii/S2095254617301230>
- 482 13. Beaulieu K, Hopkins M, Blundell J, Finlayson G. Does Habitual Physical Activity Increase  
the Sensitivity of the Appetite Control System? A Systematic Review. *Sports Med.*  
2016;46(12):1897-919.
- 484 14. Oustric P, Gibbons C, Beaulieu K, Blundell J, Finlayson G. Changes in food reward during  
weight management interventions - a systematic review. *Obes Rev.* 2018;19(12):1642 -  
486 58.
- 488 15. Thivel D, Finlayson G, Blundell JE. Homeostatic and neurocognitive control of energy  
intake in response to exercise in pediatric obesity: a psychobiological framework. *Obes*  
*Rev.* 2019;20(2):316-24.
- 490 16. Boff R de M, Liboni RPA, Batista IP de A, de Souza LH, Oliveira M da S. Weight loss  
interventions for overweight and obese adolescents: a systematic review. *Eat Weight*  
492 *Disord.* juin 2017;22(2):211-29.
- 494 17. Knöpfli BH, Radtke T, Lehmann M, Schätzle B, Eisenblätter J, Gachnang A, et al. Effects of  
a multidisciplinary inpatient intervention on body composition, aerobic fitness, and  
quality of life in severely obese girls and boys. *J Adolesc Health.* févr 2008;42(2):119-27.
- 496 18. Bianchini JAA, da Silva DF, Nardo CCS, Carolino IDR, Hernandez F, Nardo N.  
Multidisciplinary therapy reduces risk factors for metabolic syndrome in obese  
498 adolescents. *Eur J Pediatr.* févr 2013;172(2):215-21.
- 500 19. Fonvig CE, Hamann SA, Nielsen TRH, Johansen MØ, Grønbæk HN, Møllerup PM, et al.  
Subjective evaluation of psychosocial well-being in children and youths with overweight  
or obesity: the impact of multidisciplinary obesity treatment. *Qual Life Res.*  
502 2017;26(12):3279-88.
- 504 20. Stice E, Palmrose CA, Burger KS. Elevated BMI and Male Sex Are Associated with Greater  
Underreporting of Caloric Intake as Assessed by Doubly Labeled Water. *J Nutr.* oct  
2015;145(10):2412-8.
- 506 21. Miguet M, Fearnbach SN, Metz L, Khammassi M, Julian V, Cardenoux C, et al. Effect of  
HIIT versus MICT on body composition and energy intake in dietary restrained and  
508 unrestrained adolescents with obesity. *Applied Physiology, Nutrition, and Metabolism.*  
2019;
- 510 22. Miguet M, Masurier J, Chaput JP, Pereira B, Lambert C, Dâmaso AR, et al. Cognitive  
restriction accentuates the increased energy intake response to a 10-month  
512 multidisciplinary weight loss program in adolescents with obesity. *Appetite* [Internet]. 18  
déc 2018 [cité 20 déc 2018]; Disponible sur:  
514 <http://www.sciencedirect.com/science/article/pii/S0195666318310547>

- 516 23. Polivy J, Herman CP. Dieting and bingeing. A causal analysis. *Am Psychol.* févr 1985;40(2):193-201.
- 518 24. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ.* 6 mai 2000;320(7244):1240-3.
- 520 25. Cappelleri JC, Bushmakin AG, Gerber RA, Leidy NK, Sexton CC, Lowe MR, et al. Psychometric analysis of the Three-Factor Eating Questionnaire-R21: results from a large diverse sample of obese and non-obese participants. *Int J Obes (Lond).* juin 2009;33(6):611-20.
- 524 26. Pradalie, L. Alimentation et santé des lycéens et des collégiens. Agence Méditerranéenne de l'Environnement; 2003.
- 526 27. Thivel D, Genin PM, Mathieu M-E, Pereira B, Metz L. Reproducibility of an in-laboratory test meal to assess ad libitum energy intake in adolescents with obesity. *Appetite.* 01 528 2016;105:129-33.
- 530 28. Flint A, Raben A, Blundell JE, Astrup A. Reproducibility, power and validity of visual analogue scales in assessment of appetite sensations in single test meal studies. *Int J Obes Relat Metab Disord.* janv 2000;24(1):38-48.
- 532 29. Dalton M, Finlayson G. Psychobiological examination of liking and wanting for fat and sweet taste in trait binge eating females. *Physiology & Behavior.* 1 sept 534 2014;136(Supplement C):128-34.
- 536 30. Murphy SP, Poos MI. Dietary Reference Intakes: summary of applications in dietary assessment. *Public Health Nutr.* déc 2002;5(6A):843-9.
- 538 31. INPES. Adolescents | Manger Bouger [Internet]. Programme National Nutrition Santé; 2015 [cité 9 mai 2019]. Disponible sur: <http://www.mangerbouger.fr/Manger-Mieux/Manger-mieux-a-tout-age/Adolescents>
- 540 32. Feise RJ. Do multiple outcome measures require p-value adjustment? *BMC Med Res Methodol.* 17 juin 2002;2:8.
- 542 33. Singh M. Mood, food, and obesity. *Front Psychol* [Internet]. 1 sept 2014 [cité 1 mars 2015];5. Disponible sur: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4150387/>
- 544 34. Clark JE. Diet, exercise or diet with exercise: comparing the effectiveness of treatment options for weight-loss and changes in fitness for adults (18–65 years old) who are overfat, or obese; systematic review and meta-analysis. *J Diabetes Metab Disord* [Internet]. 17 avr 2015 [cité 20 juin 2019];14. Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4429709/>
- 546 548 35. Blundell JE, Gibbons C, Caudwell P, Finlayson G, Hopkins M. Appetite control and energy balance: impact of exercise. *Obes Rev.* févr 2015;16 Suppl 1:67-76.

- 552 36. Stefanik PA, Heald FP, Mayer J. Caloric intake in relation to energy output of obese and non-obese adolescent boys. *Am J Clin Nutr.* févr 1959;7(1):55-62.
- 554 37. Beaulieu K, Hopkins M, Blundell J, Finlayson G. Homeostatic and non-homeostatic appetite control along the spectrum of physical activity levels: An updated perspective. *Physiol Behav.* 01 2018;192:23-9.
- 556 38. Melby CL, Paris HL, Foright RM, Peth J. Attenuating the Biologic Drive for Weight Regain Following Weight Loss: Must What Goes Down Always Go Back Up? *Nutrients* [Internet]. 6 mai 2017 [cité 25 juin 2019];9(5). Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5452198/>
- 560 39. Doucet É, McInis K, Mahmoodianfard S. Compensation in response to energy deficits induced by exercise or diet. *Obes Rev.* 2018;19 Suppl 1:36-46.
- 562 40. Martins C, Aschehoug I, Ludviksen M, Holst J, Finlayson G, Wisloff U, et al. High-Intensity Interval Training, Appetite, and Reward Value of Food in the Obese: *Medicine & Science in Sports & Exercise.* sept 2017;49(9):1851-8.
- 564 41. Sumithran P, Prendergast LA, Delbridge E, Purcell K, Shulkes A, Kriketos A, et al. Long-term persistence of hormonal adaptations to weight loss. *N Engl J Med.* 27 oct 2011;365(17):1597-604.
- 568 42. Cameron JD, Goldfield GS, Cyr M-J, Doucet E. The effects of prolonged caloric restriction leading to weight-loss on food hedonics and reinforcement. *Physiol Behav.* 9 juin 2008;94(3):474-80.
- 570 43. Klem ML, Wing RR, Lang W, McGuire MT, Hill JO. Does weight loss maintenance become easier over time? *Obes Res.* sept 2000;8(6):438-44.
- 572 44. Cameron JD, Goldfield GS, Finlayson G, Blundell JE, Doucet É. Fasting for 24 Hours Heightens Reward from Food and Food-Related Cues. Andrews Z, éditeur. *PLoS ONE.* 16 janv 2014;9(1):e85970.
- 574 45. Charlot K, Malgoyre A, Bourrilhon C. Proposition for a shortened version of the Leeds Food Preference Questionnaire (LFPQ). *Physiology & Behavior.* 1 févr 2019;199:244-51.
- 576 46. Stice E, Spoor S, Bohon C, Veldhuizen MG, Small DM. Relation of reward from food intake and anticipated food intake to obesity: a functional magnetic resonance imaging study. *J Abnorm Psychol.* nov 2008;117(4):924-35.
- 578 47. Stice E, Burger K. Neural vulnerability factors for obesity. *Clin Psychol Rev.* mars 2019;68:38-53.
- 580 48. Oustric P, Beaulieu K, Casanova N, Husson F, Gibbons C, Hopkins M, et al. Exploring the effect of weight loss on food reward at the individual level. In: *Obesity Abstracts.* 2019.
- 584

- 586 49. Beaulieu K, Hopkins M, Gibbons C, Caudwell P, Blundell J, Finlayson G. Impact of exercise  
training on food reward and eating behaviour traits that promote overconsumption in  
individuals with overweight and obesity. In: Obesity Abstracts. 2019.
- 588 50. Finlayson G, Dalton M. Hedonics of Food Consumption: Are Food 'Liking' and 'Wanting'  
590 Viable Targets for Appetite Control in the Obese? Curr Obes Rep. 1 mars 2012;1(1):42-  
9.

592

For Review Only

## TABLES

594

**Table 1.** Anthropometric and body composition changes throughout the 10-month weight loss multidisciplinary program.

596

	<b>Baseline Mean (SD)</b>	<b>5 months Mean (SD)</b>	<b>10 months Mean (SD)</b>	<b>Time effect</b>
<b>Weight (kg)</b>	94.3 (13.9)	88.9 (13.0)***	85.4 (14.4)*** <sup>\$\$\$</sup>	<0.001
<b>BMI (kg/m<sup>2</sup>)</b>	35.7 (4.5)	33.4 (4.7)***	30.9 (5.0)*** <sup>\$\$\$</sup>	<0.001
<b>FM (%)</b>	38.7 (3.3)	34.4 (4.1)***	32.0 (5.5)*** <sup>\$\$\$</sup>	<0.001
<b>FFM (kg)</b>	55.5 (7.1)	55.9 (6.9)	55.4 (6.8)** <sup>\$\$</sup>	0.003

598 BMI: Body Mass Index; FM: Fat Mass; FFM: Fat-Free Mass; SD: Standard Deviation; \*: Significantly different from T0 (\*p<0.05; \*\*p<0.01; \*\*\*p<0.001); \$: significantly different  
600 from T1 (<sup>\$</sup>p<0.05; <sup>\$\$</sup>p<0.01; <sup>\$\$\$</sup>p<0.001).

602

604

**Table 2.** Energy intake, macronutrient consumption, appetite sensations and eating behavior trait changes throughout the 10-month weight loss multidisciplinary program.

606

		<b>Baseline (T0) Mean (SD)</b>	<b>5 months (T1) Mean (SD)</b>	<b>10 months (T2) Mean (SD)</b>	<b>Time effect</b>
<b>Energy intake</b>	Lunch EI (kcal)	1205.9 (284.1)	1290.5 (295.8)	1007.7 (250.6)** <sup>\$\$\$</sup>	<0.001
	Dinner EI (kcal)	894.1 (254.9)	849.7 (187.5)	952.6 (200.5)	0.18
	Total EI (kcal)	2699.1 (435.8)	2740.1 (420.0)	2560.2 (367.3)	0.15
<b>Macronutrients</b>	Proteins (%)	22.7 (3.7)	23.1 (2.8)	21.9 (3.5)	0.19
	Fat (%)	32.2 (6.1)	31.4 (4.1)	21.8 (3.4)*** <sup>\$\$\$</sup>	<0.001
	CHO (%)	45.2 (8.0)	45.1 (4.5)	56.0 (3.7)*** <sup>\$\$\$</sup>	<0.001
<b>Eating behaviors</b>	Restrained eating	2.2 (0.4)	2.4 (0.5)	2.2 (0.5)	0.10
	Uncontrolled eating	2.6 (0.5)	2.4 (0.6)*	2.1 (0.5)*** <sup>\$\$\$</sup>	<0.001
	Emotional eating	2.4 (0.8)	2.3 (0.7)	1.9 (0.8)*** <sup>\$\$\$</sup>	<0.001

608 EI: energy intake; SD: Standard Deviation; \*: Significantly different from T0 (\*p<0.05; \*\*p<0.01; \*\*\*p<0.001); \$: significantly different from T1 (<sup>\$</sup>p<0.05; <sup>\$\$</sup>p<0.01; <sup>\$\$\$</sup>p<0.001).

610



612


**Table 3.** Food reward throughout the 10-month weight loss multidisciplinary program.

			Baseline (T0) Mean (SD)	5 months (T1) Mean (SD)	10 months (T2) Mean (SD)	Time effect	Meal effect	Interaction time x meal effect		
								T0 vs T1	T0 vs T2	T1 vs T2
Exp L	HFSA	Before	43.1 (27.6)	52.4 (23.9)*	38.1 (26.8) <sup>\$\$</sup>	0.01	<0.001	<0.001	0.04	0.07
		After	39.7 (22.1)	21.5 (23.5) <sup>#####</sup>	19.9 (21.3) <sup>#####</sup>	<0.001	<0.001	<0.001	0.04	0.07
	LFSA	Before	32.1 (22.1)	38.3 (19.1)*	24.3 (20.1) <sup>\$\$\$</sup>	0.003	<0.001	0.001	0.20	0.06
		After	31.2 (20.1)	18.3 (20.6) <sup>####</sup>	15.6 (21.7) <sup>**#</sup>	0.002	<0.001	0.001	0.20	0.06
	HFSW	Before	48.2 (27.0)	59.0 (21.5)**	48.5 (26.9) <sup>\$</sup>	0.01	<0.001	0.001	0.22	0.07
		After	39.7 (22.4)	28.9 (25.2) <sup>####</sup>	31.2 (32.9) <sup>#</sup>	0.05	<0.001	0.001	0.22	0.07
	LFSW	Before	42.7 (23.3)	51.8 (23.8)**	35.9 (23.8) <sup>\$\$\$</sup>	<0.001	<0.001	<0.001	0.02	0.07
		After	45.4 (25.2)	25.8 (25.2) <sup>#####</sup>	22.2 (25.6) <sup>#####</sup>	<0.001	<0.001	<0.001	0.02	0.07
Imp W	HFSA	Before	3.0 (25.1)	-3.2 (26.9)	0.6 (19.1)	0.48	0.79	0.75	0.73	0.52
		After	2.0 (34.7)	-1.5 (26.9)	-2.7 (32.4)	0.77	0.79	0.75	0.73	0.52
	LFSA	Before	-18.9 (25.5)	-20.3 (30.2)	-23.7 (22.8)	0.80	0.11	0.11	0.51	0.40
		After	-6.4 (42.4)	-21.4 (32.8)*	-16.9 (29.0) <sup>#</sup>	0.11	0.11	0.11	0.51	0.40
	HFSW	Before	14.1 (24.5)	17.7 (25.4)	19.2 (29.8)	0.84	0.72	0.07	0.09	0.97
		After	6.0 (30.3)	20.8 (36.3)*	24.4 (26.7)*	0.03	0.72	0.07	0.09	0.97
	LFSW	Before	5.0 (20.9)	6.3 (25.2)	5.0 (20.3)	0.82	0.60	0.86	0.68	0.79
		After	5.4 (33.2)	4.6 (30.4)	0.9 (35.8)	0.83	0.60	0.86	0.68	0.79


614

HFSA: high fat savory; LFSA: low fat savory; HFSW: high fat sweet; LFSW: low fat sweet;  
 616 Exp L: explicit liking; Imp W: implicit wanting; SD: Standard Deviation; \*: Significantly  
 different from T0 (\*p<0.05; \*\*p<0.01; \*\*\*p<0.001); \$: Significantly different from T1  
 618 (\$p<0.05; \$\$p<0.01; \$\$\$p<0.001); #: Significantly different from pre-meal (#p<0.05; ##p<0.01;  
 ###p<0.001).

620



**Figure 1.** Fasting (A) and daily (B) appetite sensations throughout the 10-month multidisciplinary weight loss program. Solid bar: mean; Error bar: standard deviation; VAS: visual analogue scale; AUC: area under the curves; DTE: desire to eat; PFC: prospective food consumption; \*: time effect (\* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ ).



**Figure 2.** Variability among explicit liking and implicit wanting changes in the hungry state in response to a 10-month multidisciplinary weight loss program. A: Changes between T0 and T1; B: Changes between T1 and T2. T0: 0 month; T1: 5 months; T2: 10 months; HFSA: high fat savory; LFSA: low fat savory; HFSW: high fat sweet; LFSW: low fat sweet; \*: significant differences between T0 and T1 or between T1 and T2 (\* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ ); the diagram shows the median, the lower and upper quartile and the individual variations.