

1 **SUPPLEMENTARY MATERIALS: A Variational Model Dedicated to Joint
2 Segmentation, Registration and Atlas Generation for Shape Analysis***

3 Noémie Debroux[†], John Aston[‡], Fabien Bonardi[§], Alistair Forbes[¶], Carole Le Guyader^{||},
4 Marina Romanchikova^{||}, and Carola-Bibiane Schönlieb[†]

6 **SM1. Detailed proof of Theorem 2.6.**

7 *Proof.* The proof follows the calculus of variations arguments. We first have that \mathcal{F}_1 is

8 bounded below by 0 and that $\mathcal{F}_1(\theta_R, \{\theta_{T_i}, \varphi_i\}_{i=1}^M) = \frac{1}{M} \sum_{i=1}^M (\|T_i - \bar{T}_i\|_{L^2(\Omega)}^2 + \|T_i - \bar{T}\|_{L^2(\Omega)}^2) <$
9 $+\infty$ for $\theta_R = (1, 0, \dots, 0)$, $\theta_{T_i} = (1, 0, \dots, 0)$, $\forall i = 1, \dots, M$, $\varphi_i = \text{Id}$, $\forall i = 1, \dots, M$ and
10 with $\bar{T}_i = \frac{\int_{\Omega} T_i dx}{\text{meas}(\Omega)}$ and $\bar{T} = \frac{1}{M} \sum_{i=1}^M \frac{\int_{\Omega} T_i dx}{\text{meas}(\Omega)}$. Therefore, the infimum exists and is finite.

11 Let $(\theta_R^k, \{\theta_{T_i}^k, \varphi_i^k\}_{i=1}^M)_k \in \mathcal{U} \times \mathcal{U}^M \times \hat{\mathcal{W}}^M$ such that $\theta_{T_i, l}^k \circ \varphi_i^k - \theta_{R, l}^k \in BV(\Omega)$ for all $l \in$
12 $\{1, \dots, N\}$, and for all $i \in \{1, \dots, M\}$ be a minimizing sequence such that $\lim_{k \rightarrow +\infty} \mathcal{F}_1(\theta_R^k,$
13 $\{\theta_{T_i}^k, \varphi_i^k\}_{i=1}^M) = \inf \mathcal{F}_1(\theta_R, \{\theta_{T_i}, \varphi_i\}_{i=1}^M) < +\infty$. Hence there exists $K \in \mathbb{N}$ such that $\forall k \in \mathbb{N}$,
14 $k \geq K \Rightarrow \mathcal{F}_1(\theta_R^k, \{\theta_{T_i}^k, \varphi_i^k\}_{i=1}^M) \leq \inf \mathcal{F}_1(\theta_R, \{\theta_{T_i}, \varphi_i\}_{i=1}^M) + 1 < +\infty$. From now on, we consider
15 $k \geq K$.

16 One thus has

$$\begin{aligned} 17 \quad & \frac{1}{M} \sum_{i=1}^M \left(\frac{\gamma_T}{2} \sum_{l=1}^N TV(\theta_{T_i, l}^k) + \frac{\gamma_R}{2} \sum_{l=1}^N TV(\theta_{R, l}^k) + a_1 \|\nabla \varphi_i\|_{L^4(\Omega, M_3(\mathbb{R}))}^4 - 9a_1 \text{meas}(\Omega) \right. \\ 18 \quad & + a_2 \|\text{Cof} \nabla \varphi_i\|_{L^4(\Omega, M_3(\mathbb{R}))}^4 - 9a_2 \text{meas}(\Omega) + \frac{a_3}{2} \|\det \nabla \varphi_i\|_{L^2(\Omega)}^2 - (a_3 + a_4) \text{meas}(\Omega) \\ 19 \quad & \left. + \mathbf{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha\}}(\nabla \varphi_i) + \mathbf{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta\}}((\nabla \varphi_i)^{-1}) \right) \\ 20 \quad & \leq \mathcal{F}_1(\theta_R^k, \{\theta_{T_i}^k, \varphi_i^k\}_{i=1}^M) \leq \inf \mathcal{F}_1(\theta_R, \{\theta_{T_i}, \varphi_i\}_{i=1}^M) + 1 < +\infty. \end{aligned}$$

22 We deduce from this inequality that $\forall i = 1, \dots, M$,

*Submitted to the editors DATE.

Funding: This project is co-financed by the European Union with the European regional development fund (ERDF, HN0002137) and by the Normandie Regional Council via the M2NUM project.

[†]Department of Applied Mathematics and Theoretical Physics (DAMPT), Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, UK (nd448@cam.ac.uk, cbs31@cam.ac.uk).

[‡]Statslab, Department of Pure Mathematics and Mathematical Statistics, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, UK (j.aston@statslab.cam.ac.uk).

[§]IBISC, Université d'Évry, 36, Rue du Pelvoux, CE1455 Courcouronnes, 91020 Evry Cédex, France (fabien.bonardi@univ-evry.fr).

[¶]National Physical Laboratory, Hampton Road, Teddington, Middlesex, TW11 0LW UK (alistair.forbes, marina.romanchikova@npl.co.uk).

^{||}Normandie Univ, Institut National des Sciences Appliquées de Rouen, Laboratory of Mathematics, 76000 Rouen, France (carole.le-guyader@insa-rouen.fr).

- (φ_i^k) is uniformly bounded according to k in $W^{1,4}(\Omega)$ by using the generalized Poincaré's inequality and the fact that $\varphi_i^k = \text{Id}$ on $\partial\Omega$.
 - $(\nabla\varphi_i^k)$ is uniformly bounded according to k in $L^\infty(\Omega, M_3(\mathbb{R}))$.
 - $((\nabla\varphi_i^k)^{-1})$ is uniformly bounded according to k in $L^\infty(\Omega, M_3(\mathbb{R}))$.
 - $(\text{Cof}\nabla\varphi_i^k)$ is uniformly bounded according to k in $L^4(\Omega, M_3(\mathbb{R}))$.
 - $(\det\nabla\varphi_i^k)$ is uniformly bounded according to k in $L^2(\Omega)$.
 - $\forall l = 1, \dots, N$, $(\theta_{T_i,l}^k)$ is uniformly bounded according to k in $BV(\Omega)$ since $(TV(\theta_{T_i,l}^k))$ is uniformly bounded according to k and $\|\theta_{T_i,l}^k\|_{L^1(\Omega)} \leq \text{meas}(\Omega)$ for all $k \in \mathbb{N}$.
 - $\forall l = 1, \dots, N$, $(\theta_{R,l}^k)$ is uniformly bounded according to k in $BV(\Omega)$ since $(TV(\theta_{R,l}^k))$ is uniformly bounded according to k and $\|\theta_{R,l}^k\|_{L^1(\Omega)} \leq \text{meas}(\Omega)$ for all $k \in \mathbb{N}$.
- Since we have assumed enough regularity on the boundary of Ω , we can apply [?, Theorem 136] and extract subsequences still denoted $(\theta_{T_i,l}^k)$ and $(\theta_{R,l}^k)$ such that

$$\theta_{T_i,l}^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{T_i,l} \in BV(\Omega) \text{ in } L^1(\Omega),$$

$$\theta_{R,l}^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{R,l} \in BV(\Omega) \text{ in } L^1(\Omega).$$

As the convergence is strong in $L^1(\Omega)$, then we can extract subsequences of $(\theta_{T_i,l}^k)_k$ and $(\theta_{R,l}^k)_k$ still denoted $(\theta_{T_i,l}^k)_k$ and $(\theta_{R,l}^k)_k$ such that $\theta_{T_i,l}^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{T_i,l}$ almost everywhere in Ω and $\theta_{R,l}^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{R,l}$ almost everywhere in Ω and thus $\bar{\theta}_{T_i,l} \in BV(\Omega, \{0, 1\})$ and $\bar{\theta}_{R,l} \in BV(\Omega, \{0, 1\})$. Besides, $\sum_{l=1}^N \theta_{T_i,l}^k(x) = 1 \xrightarrow[k \rightarrow +\infty]{} \sum_{l=1}^N \bar{\theta}_{T_i,l}(x) = 1$ almost everywhere in Ω and $\sum_{l=1}^N \theta_{R,l}^k(x) = 1 \xrightarrow[k \rightarrow +\infty]{} \sum_{l=1}^N \bar{\theta}_{R,l}(x) = 1$ so that $\bar{\theta}_{T_i} \in \mathcal{U}$ and $\bar{\theta}_R \in \mathcal{U}$.

There also exist subsequences still denoted $(\varphi_i^k)_k$ such that

$$\varphi_i^k \xrightarrow[k \rightarrow +\infty]{} \bar{\varphi}_i \text{ in } W^{1,4}(\Omega, \mathbb{R}^3).$$

- By continuity of the trace operator, we deduce that $\bar{\varphi}_i \in \text{Id} + W_0^{1,4}(\Omega, \mathbb{R}^3)$. We can also extract subsequences still denoted $(\text{Cof}\nabla\varphi_i^k)_k$ and $(\det\nabla\varphi_i^k)_k$ such that

$$\text{Cof}\nabla\varphi_i^k \xrightarrow[k \rightarrow +\infty]{} \bar{X}_i \text{ in } L^4(\Omega, M_3(\mathbb{R})),$$

$$\det\nabla\varphi_i^k \xrightarrow[k \rightarrow +\infty]{} \bar{\delta}_i \text{ in } L^2(\Omega),$$

and by [?, Theorem VI.3.3], we deduce that $\bar{X}_i = \text{Cof}\nabla\bar{\varphi}_i$ and $\bar{\delta}_i = \det\nabla\bar{\varphi}_i$.

There exist subsequences still denoted $(\nabla\varphi_i^k)$ and $((\nabla\varphi_i^k)^{-1})$ such that

$$(\nabla\varphi_i^k)^{-1} \xrightarrow[k \rightarrow +\infty]{*} u_i \text{ in } L^\infty(\Omega, M_3(\mathbb{R})),$$

$$\nabla\varphi_i^k \xrightarrow[k \rightarrow +\infty]{*} \nabla\bar{\varphi}_i \text{ in } L^\infty(\Omega, M_3(\mathbb{R})),$$

48 by uniqueness of the weak limit in $L^4(\Omega, M_3(\mathbb{R}))$ and the continuous embedding of $L^\infty(\Omega, M_3(\mathbb{R}))$
 49 into $L^4(\Omega, M_3(\mathbb{R}))$, and thus $\bar{\varphi}_i \in \text{Id} + W_0^{1,\infty}(\Omega, \mathbb{R}^3)$.
 50 Let us now prove that $u_i = (\nabla \bar{\varphi}_i)^{-1}$ for each i . For all $p \in L^1(\Omega, M_3(\mathbb{R}))$, we have that

$$\begin{aligned} 51 \quad & \int_{\Omega} ((\nabla \varphi_i^k)^{-1} \nabla \bar{\varphi}_i - \text{I}_3) : p \, dx = \int_{\Omega} ((\nabla \varphi_i^k)^{-1} \nabla \bar{\varphi}_i - u_i \nabla \bar{\varphi}_i + u_i \nabla \bar{\varphi}_i - u_i \nabla \varphi_i^k \\ 52 \quad & \quad + u_i \nabla \varphi_i^k - (\nabla \varphi_i^k)^{-1} \nabla \varphi_i^k) : p \, dx, \\ 53 \quad & = \int_{\Omega} (((\nabla \varphi_i^k)^{-1} - u_i) \nabla \bar{\varphi}_i) : p \, dx + \int_{\Omega} (u_i (\nabla \bar{\varphi}_i - \nabla \varphi_i^k)) : p \, dx \\ 54 \quad & \quad + \int_{\Omega} ((u_i - (\nabla \varphi_i^k)^{-1}) \nabla \varphi_i^k) : p \, dx, \\ 55 \quad & = \int_{\Omega} ((\nabla \varphi_i^k)^{-1} - u_i) : p (\nabla \bar{\varphi}_i)^T \, dx + \int_{\Omega} (\nabla \bar{\varphi}_i - \nabla \varphi_i^k) : u_i^T p \, dx \\ 56 \quad & \quad + \int_{\Omega} (u_i - (\nabla \varphi_i^k)^{-1}) : p (\nabla \varphi_i^k)^T \, dx. \\ 57 \end{aligned}$$

58 Since $\nabla \bar{\varphi}_i \in L^\infty(\Omega, M_3(\mathbb{R}))$, $u_i \in L^\infty(\Omega, M_3(\mathbb{R}))$ and for all $k \in \mathbb{N}$, $\|\nabla \varphi_i^k\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq$
 59 α , then we have that $p(\nabla \bar{\varphi}_i)^T \in L^1(\Omega, M_3(\mathbb{R}))$ so $\int_{\Omega} ((\nabla \varphi_i^k)^{-1} - u_i) : p(\nabla \bar{\varphi}_i)^T \, dx \xrightarrow[k \rightarrow +\infty]{} 0$,
 60 $u_i^T p \in L^1(\Omega, M_3(\mathbb{R}))$ so $\int_{\Omega} (\nabla \bar{\varphi}_i - \nabla \varphi_i^k) : u_i^T p \, dx \xrightarrow[k \rightarrow +\infty]{} 0$, and $|\int_{\Omega} (u_i - (\nabla \varphi_i^k)^{-1}) :$
 61 $p(\nabla \varphi_i^k)^T \, dx| \leq \alpha |\int_{\Omega} (u_i - (\nabla \varphi_i^k)^{-1}) : p' \, dx| \xrightarrow[k \rightarrow +\infty]{} 0$ with $p' \in L^1(\Omega, M_3(\mathbb{R}))$. Therefore,
 62 we get $(\nabla \varphi_i^k)^{-1} \nabla \bar{\varphi}_i \xrightarrow[k \rightarrow +\infty]{*} \text{I}_3$ in $L^\infty(\Omega, M_3(\mathbb{R}))$. But we know that $(\nabla \varphi_i^k)^{-1} \xrightarrow[k \rightarrow +\infty]{*} u_i$ in
 63 $L^\infty(\Omega, M_3(\mathbb{R}))$, so that $(\nabla \varphi_i^k)^{-1} \nabla \bar{\varphi}_i \xrightarrow[k \rightarrow +\infty]{*} u_i \nabla \bar{\varphi}_i = \text{I}_3$ in $L^\infty(\Omega, M_3(\mathbb{R}))$ and $u_i = (\nabla \bar{\varphi}_i)^{-1}$.
 64 Since $\theta_{T_i,l}^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{T_i,l}$ almost everywhere in Ω , then $T_i \theta_{T_i,l}^k \xrightarrow[k \rightarrow +\infty]{} T_i \bar{\theta}_{T_i,l}$ almost everywhere
 65 in Ω . Furthermore, $|\theta_{T_i,l}^k| \leq 1 \in L^1(\Omega)$ almost everywhere in Ω as Ω is bounded, and
 66 $|T_i \theta_{T_i,l}^k| \leq \|T_i\|_{L^\infty(\Omega)} \in L^1(\Omega)$ almost everywhere in Ω . We can thus apply the dominated
 67 convergence theorem and get

$$\begin{aligned} 68 \quad & \int_{\Omega} \theta_{T_i,l}^k \, dx \xrightarrow[k \rightarrow +\infty]{} \int_{\Omega} \bar{\theta}_{T_i,l} \, dx, \\ 69 \quad & \int_{\Omega} \theta_{T_i,l}^k T_i \, dx \xrightarrow[k \rightarrow +\infty]{} \int_{\Omega} \bar{\theta}_{T_i,l} T_i \, dx, \\ 70 \quad & c_{T_i,l}^k \xrightarrow[k \rightarrow +\infty]{} \bar{c}_{T_i,l}. \\ 71 \end{aligned}$$

72 As $T_i \in W^{1,\infty}(\Omega)$, we denote κ_i the Lipschitz constant related to T_i . Then the chain rule
 73 applies and we have that $T_i \circ \varphi_i^k \in W^{1,\infty}(\Omega)$, $T_i \circ \bar{\varphi}_i \in W^{1,\infty}(\Omega)$ and $\|T_i \circ \varphi_i^k - T_i \circ \bar{\varphi}_i\|_{C^0(\Omega)} \leq$
 74 $\kappa_i \|\varphi_i^k - \bar{\varphi}_i\|_{C^0(\Omega)} \xrightarrow[k \rightarrow +\infty]{} 0$ using the Sobolev compact embedding $W^{1,\infty}(\Omega) \hookrightarrow C^0(\Omega)$. Therefore,
 75 up to a subsequence, $\theta_{R,l}^k T_i \circ \varphi_i^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{R,l} T_i \circ \bar{\varphi}_i$ almost everywhere in Ω , $\theta_{R,l}^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{R,l}$
 76 almost everywhere in Ω with $|\theta_{R,l}^k T_i \circ \varphi_i^k| \leq \|T_i\|_{L^\infty(\Omega)} \in L^1(\Omega)$ and $|\theta_{R,l}^k| \leq 1 \in L^1(\Omega)$ almost

77 everywhere in Ω . Thus the dominated convergence theorem applies and we get

$$78 \quad \frac{1}{M} \sum_{i=1}^M \int_{\Omega} \theta_{R,l}^k T_i \circ \varphi_i^k dx \xrightarrow{k \rightarrow +\infty} \frac{1}{M} \sum_{i=1}^M \int_{\Omega} \bar{\theta}_{R,l} T_i \circ \bar{\varphi}_i dx,$$

$$79 \quad \int_{\Omega} \theta_{R,l}^k dx \xrightarrow{k \rightarrow +\infty} \int_{\Omega} \bar{\theta}_{R,l} dx,$$

$$80 \quad c_{R,l}^k \xrightarrow{k \rightarrow +\infty} \bar{c}_{R,l}.$$

82 We also know that W_{Op} is continuous and convex. If $\psi_n \xrightarrow{n \rightarrow +\infty} \psi$ in $W^{1,4}(\Omega, \mathbb{R}^3)$, then
83 $\nabla \psi_n \xrightarrow{n \rightarrow +\infty} \nabla \psi$ in $L^4(\Omega, M_3(\mathbb{R}))$ and one can extract a subsequence still denoted $(\nabla \psi_n)$ such
84 that $\nabla \psi_n \xrightarrow{n \rightarrow +\infty} \nabla \psi$ almost everywhere in Ω . If $\alpha_n \xrightarrow{n \rightarrow +\infty} \alpha$ in $L^4(\Omega, M_3(\mathbb{R}))$, then one can
85 extract a subsequence still denoted (α_n) such that $\alpha_n \xrightarrow{n \rightarrow +\infty} \alpha$ almost everywhere in Ω . If
86 $\delta_n \xrightarrow{n \rightarrow +\infty} \delta$ in $L^2(\Omega)$ then one can extract a subsequence still denoted (δ_n) such that $\delta_n \xrightarrow{n \rightarrow +\infty} \delta$
87 almost everywhere in Ω . Then by continuity of W_{Op} , we get that $W_{Op}(\nabla \psi_n(x), \alpha_n(x), \delta_n(x))$
88 $\xrightarrow{n \rightarrow +\infty} W_{Op}(\nabla \psi(x), \alpha(x), \delta(x))$ almost everywhere in Ω . We then apply Fatou's lemma and get
89 $\liminf_{n \rightarrow +\infty} \int_{\Omega} W_{Op}(\nabla \psi_n(x), \alpha_n(x), \delta_n(x)) dx \geq \int_{\Omega} W_{Op}(\nabla \psi(x), \alpha(x), \delta(x)) dx$. As W_{Op} is convex,
90 so is $\int_{\Omega} W_{Op}(\xi(x), \alpha(x), \delta(x)) dx$ and we can apply [?, Corollaire III.8]. Therefore $\int_{\Omega} W_{Op}(\xi(x),$
91 $\alpha(x), \delta(x)) dx$ is also weakly lower semicontinuous in $L^4(\Omega, M_3(\mathbb{R})) \times L^4(\Omega, M_3(\mathbb{R})) \times L^2(\Omega)$.
92 We then deduce that

$$93 \quad +\infty > \liminf_{k \rightarrow +\infty} \int_{\Omega} W_{Op}(\nabla \varphi_i^k, \text{Cof} \nabla \varphi_i^k, \det \nabla \varphi_i^k) dx \geq \int_{\Omega} W_{Op}(\nabla \bar{\varphi}_i, \text{Cof} \nabla \bar{\varphi}_i, \det \nabla \bar{\varphi}_i) dx.$$

95 Let $\bar{q} = q \frac{1+s}{q+s} = 4 \frac{1+10}{4+10} = \frac{44}{14} > 3$. We then compute

$$96 \quad \int_{\Omega} \|(\nabla \bar{\varphi}_i)^{-1}\|_F^{\bar{q}} \det \nabla \bar{\varphi}_i dx = \int_{\Omega} \left\| \frac{\text{Cof} \nabla \bar{\varphi}_i^T}{\det \nabla \bar{\varphi}_i} \right\|_F^{\bar{q}} \det \nabla \bar{\varphi}_i dx,$$

$$97 \quad = \int_{\Omega} \|\text{Cof} \nabla \bar{\varphi}_i\|_F^{\bar{q}} (\det \nabla \bar{\varphi}_i)^{1-\bar{q}} dx,$$

$$99 \quad \leq \|\text{Cof} \nabla \bar{\varphi}_i\|_{L^4(\Omega)}^{\bar{q}} \|(\det \nabla \bar{\varphi}_i)^{-1}\|_{L^{10}(\Omega)}^{1-\bar{q}} < +\infty,$$

100 from Hölder's inequality and the finiteness of $\int_{\Omega} W_{Op}(\nabla \bar{\varphi}_i, \text{Cof} \nabla \bar{\varphi}_i, \det \nabla \bar{\varphi}_i) dx$. Thus Ball's
101 conditions are satisfied [?, Theorems 1 and 2] and $\bar{\varphi}_i$ are bi-Hölder homeomorphisms.

102 By the weak-* lower semicontinuity of $\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))}$, we have that $\|\nabla \bar{\varphi}_i\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq$
103 $\liminf_{k \rightarrow +\infty} \|\nabla \varphi_i^k\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha$ and $\|(\nabla \bar{\varphi}_i)^{-1}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \liminf_{k \rightarrow +\infty} \|(\nabla \varphi_i^k)^{-1}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta$
104 so that $\mathbf{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha\}}(\nabla \bar{\varphi}_i) = 0 \leq \liminf_{k \rightarrow +\infty} \mathbf{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha\}}(\nabla \varphi_i^k)$ and $\mathbf{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta\}}$
105 $((\nabla \bar{\varphi}_i)^{-1}) = 0 \leq \liminf_{k \rightarrow +\infty} \mathbf{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta\}}((\nabla \varphi_i^k)^{-1})$. The same reasoning applies for each
106 $i = 1, \dots, M$.

107 We also have

$$108 \quad \|\nabla(\bar{\varphi}_i^{-1})\|_{L^\infty(\Omega, M_3(\mathbb{R}))} = \|(\nabla \bar{\varphi}_i)^{-1}(\bar{\varphi}_i^{-1})\|_{L^\infty(\Omega, M_3(\mathbb{R}))} = \|(\nabla \bar{\varphi}_i)^{-1}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta < +\infty,$$

110 since $\bar{\varphi}_i$ is an homeomorphism from Ω to $\text{Id}(\Omega) = \Omega$, according to Ball's results, meaning that
111 $\bar{\varphi}_i^{-1}$ is a one to one mapping from Ω to Ω . So, using Poincaré-Wirtinger inequality, we deduce
112 that $\bar{\varphi}_i$ is a bi-Lipschitz homeomorphism. The same reasoning applies for the sequence $(\varphi_i^k)_k$.

113 We now have that $\forall i \in \{1, \dots, M\}$, $\forall l \in \{1, \dots, N\}$,

$$114 \bullet \theta_{T_i,l}^k \xrightarrow{k \rightarrow +\infty} \bar{\theta}_{T_i,l} \text{ in } L^1(\Omega) \text{ with } \bar{\theta}_{T_i,l} \in BV(\Omega, \{0, 1\}).$$

$$115 \bullet \varphi_i^k \xrightarrow{k \rightarrow +\infty} \bar{\varphi}_i \text{ in } W^{1,4}(\Omega, \mathbb{R}^3) \text{ with } (\varphi_i^k) \text{ bi-Lipschitz homeomorphisms } (\varphi_i^k \in C^{0,\alpha}(\Omega, \mathbb{R}^3), \\ 116 (\varphi_i^k)^{-1} \in C^{0,\alpha'}(\Omega, \mathbb{R}^3)), \bar{\varphi}_i \text{ bi-Lipschitz homeomorphisms } (\bar{\varphi}_i \in C^{0,\alpha}(\Omega, \mathbb{R}^3), (\bar{\varphi}_i)^{-1} \in \\ 117 C^{0,\alpha'}(\Omega, \mathbb{R}^3)) \text{ and } \varphi_i^k \xrightarrow{k \rightarrow +\infty} \bar{\varphi}_i \text{ in } C^{0,\alpha}(\Omega, \mathbb{R}^3), \alpha < \frac{1}{4} \text{ and } \alpha' < 1 - \frac{3}{q} \text{ using the Sobolev} \\ 118 \text{embeddings properties.}$$

119 We then can prove that $\varphi_i^k \circ \bar{\varphi}_i^{-1} \xrightarrow{k \rightarrow +\infty} \text{Id}$ in $C^{0,\alpha''}(\Omega, \mathbb{R}^3)$ with $\alpha'' = \alpha\alpha'$ for all $i \in \{1, \dots, M\}$
120 and all $l \in \{1, \dots, N\}$. Indeed one has

$$121 \sup_{x \in \Omega} |\varphi_i^k \circ \bar{\varphi}_i^{-1}(x) - \bar{\varphi}_i \circ \bar{\varphi}_i^{-1}(x)| + \sup_{x \neq y} \frac{|\varphi_i^k \circ \bar{\varphi}_i^{-1}(x) - \varphi_i^k \circ \bar{\varphi}_i^{-1}(y) - \bar{\varphi}_i \circ \bar{\varphi}_i^{-1}(x) + \bar{\varphi}_i \circ \bar{\varphi}_i^{-1}(y)|}{|x - y|^{\alpha''}} \\ 122 \leq \sup_{x \in \Omega} |\varphi_i^k(x) - \bar{\varphi}_i(x)| + \sup_{x \neq y} \frac{|\varphi_i^k \circ \bar{\varphi}_i^{-1}(x) - \varphi_i^k \circ \bar{\varphi}_i^{-1}(y) - \bar{\varphi}_i \circ \bar{\varphi}_i^{-1}(x) + \bar{\varphi}_i \circ \bar{\varphi}_i^{-1}(y)|}{|\bar{\varphi}_i^{-1}(x) - \bar{\varphi}_i^{-1}(y)|^\alpha} \\ 123 \cdot \frac{|\bar{\varphi}_i^{-1}(y) - \bar{\varphi}_i^{-1}(x)|^\alpha}{|x - y|^{\alpha''}} \text{ as } \bar{\varphi}_i^{-1} \text{ is a one to one mapping from } \Omega \text{ to } \Omega, \\ 124 \leq \sup_{x \in \Omega} |\varphi_i^k(x) - \bar{\varphi}_i(x)| + \sup_{x \neq y} \frac{|\varphi_i^k(x) - \varphi_i^k(y) - \bar{\varphi}_i(x) + \bar{\varphi}_i(y)|}{|x - y|^\alpha} \cdot \sup_{x \neq y} \frac{|\bar{\varphi}_i^{-1}(y) - \bar{\varphi}_i^{-1}(x)|^\alpha}{|x - y|^{\alpha''}}, \\ 125 \leq (1 + \|\bar{\varphi}_i^{-1}\|_{C^{0,\alpha'}(\Omega, \mathbb{R}^3)}^\alpha) \|\varphi_i^k - \bar{\varphi}_i\|_{C^{0,\alpha}(\Omega, \mathbb{R}^3)} \xrightarrow{k \rightarrow +\infty} 0. \quad \blacksquare$$

127 Let us now prove that $\theta_{T_i,l}^k \circ \varphi_i^k \xrightarrow{k \rightarrow +\infty} \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i$ in $L^1(\Omega)$ based on the proof of [?, Lemma 5.1].

128 Since $\varphi_i^k \circ \bar{\varphi}_i^{-1} \xrightarrow{k \rightarrow +\infty} \text{Id}$ in $C^{0,\alpha''}(\Omega)$ and so uniformly, we can write

$$129 \forall \varepsilon > 0, \exists K \in \mathbb{N}, \forall k \in \mathbb{N}, k \geq K \Rightarrow \sup_{x \in \Omega} |\varphi_i^k \circ \bar{\varphi}_i^{-1}(x) - x| \leq \varepsilon.$$

131 Also, given $\eta > 0$, by Egorov's theorem, there exists a set S_η with Lebesgue measure
132 $\text{meas}(S_\eta) \leq \eta$ such that $\theta_{T_i,l}^k \xrightarrow{k \rightarrow +\infty} \bar{\theta}_{T_i,l}$ uniformly in $\Omega \setminus S_\eta$.

133 Now, we choose K large enough such that $\forall k \in \mathbb{N}, k \geq K$ then

$$134 |\varphi_i^k \circ \bar{\varphi}_i(x) - x| \leq \eta \text{ in } \Omega,$$

$$135 |\theta_{T_i,l}^k(x) - \bar{\theta}_{T_i,l}(x)| \leq \eta \text{ in } \Omega \setminus S_\eta.$$

137 We also denote $\mathcal{O} = \{x \in \Omega \mid \bar{\theta}_{T_i,l}(x) = 1\}$, $(\mathcal{O} \setminus S_\eta)_\eta = \{x \in \mathcal{O} \setminus S_\eta \mid \text{dist}(x, \partial(\mathcal{O} \setminus S_\eta)) > \eta\}$,
138 $\mathcal{O}^\eta = \{x \in \Omega \mid \text{dist}(x, \partial\mathcal{O}) \leq \eta\}$. Then $\forall x \in (\mathcal{O} \setminus S_\eta)_\eta$, $\varphi_i^k \circ \bar{\varphi}_i^{-1}(x) \in \mathcal{O} \setminus S_\eta$ and $\forall x \in$
139 $(\Omega \setminus (S_\eta \cup \mathcal{O}))_\eta$, $\varphi_i^k \circ \bar{\varphi}_i^{-1}(x) \in \Omega \setminus (S_\eta \cup \mathcal{O})$. We then have

$$140 \int_{\Omega} |\theta_{T_i,l}^k \circ \varphi_i^k - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i| dx = \int_{\Omega} \frac{|\theta_{T_i,l}^k \circ \varphi_i^k \circ \bar{\varphi}_i^{-1}(x) - \bar{\theta}_{T_i,l}(x)|}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx,$$

142

$$\begin{aligned}
 143 & \leq \int_{(\mathcal{O} \setminus S_\eta)_\eta} \frac{|\theta_{T_i,l}^k \circ \varphi_i^k \circ \bar{\varphi}_i^{-1}(x) - 1|}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx + \int_{(\Omega \setminus (\mathcal{O} \cup S_\eta))_\eta} \frac{|\theta_{T_i,l}^k \circ \varphi_i^k \circ \bar{\varphi}_i^{-1}(x)|}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx \\
 144 & + \int_{S_\eta \cup S_\eta^\eta \cup \mathcal{O}^\eta \cup \Omega^\eta} \frac{|\theta_{T_i,l}^k \circ \varphi_i^k \circ \bar{\varphi}_i^{-1}(x) - \bar{\theta}_{T_i,l}(x)|}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx, \\
 145 & \leq \eta \int_{(\mathcal{O} \setminus S_\eta)_\eta} \frac{1}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx + \eta \int_{(\Omega \setminus (\mathcal{O} \cup S_\eta))_\eta} \frac{1}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx \\
 146 & + 2 \int_{S_\eta \cup S_\eta^\eta \cup \mathcal{O}^\eta \cup \Omega^\eta} \frac{1}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx, \\
 147 & \leq (\eta \text{meas}(\Omega)^{\frac{10}{11}} + \eta \text{meas}(\Omega)^{\frac{10}{11}} + 2(\text{meas}(S_\eta) + \text{meas}(S_\eta^\eta) + \text{meas}(\mathcal{O}^\eta) + \text{meas}(\Omega^\eta))^{\frac{10}{11}}) \\
 148 & (\int_{\Omega} (\det \nabla \bar{\varphi}_i(x))^{-10} dx)^{\frac{1}{11}}, \\
 149 & \leq (2\eta \text{meas}(\Omega)^{\frac{10}{11}} + 2(\frac{4\pi}{3}\eta^3(\text{meas}(\partial S_\eta) + \text{meas}(\partial \mathcal{O}) + \text{meas}(\partial \Omega)) + \eta)^{\frac{10}{11}}) \|\det \nabla \bar{\varphi}_i(x)^{-1}\|_{L^{10}(\Omega)}^{\frac{10}{11}}, \\
 150 & \blacksquare
 \end{aligned}$$

152 with $\|\det \nabla \bar{\varphi}_i(x)^{-1}\|_{L^{10}(\Omega)}^{\frac{10}{11}} < +\infty$. By letting η tend to 0, we obtain that $\theta_{T_i,l}^k \circ \varphi_i^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i$ in $L^1(\Omega)$, for all $i \in \{1, \dots, M\}$ and for all $l \in \{1, \dots, N\}$.
 153 Since $\theta_{T_i,l}^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{T_i,l}$ in $L^1(\Omega)$ and $\theta_{R,l}^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{R,l}$ in $L^1(\Omega)$ and $\theta_{T_i,l}^k \circ \varphi_i^k - \theta_{R,l}^k \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i - \bar{\theta}_{R,l}$ for all $l \in \{1, \dots, N\}$ and for all $i \in \{1, \dots, M\}$, we deduce by the lower
 154 semicontinuity of the total variation that $\frac{1}{M} \sum_{i=1}^M \sum_{l=1}^N \frac{\gamma_T}{2} TV(\bar{\theta}_{T_i,l}) + \frac{\gamma_R}{2} TV(\bar{\theta}_{R,l}) + \frac{\lambda}{2} TV(\bar{\theta}_{T_i,l} \circ \bar{\varphi}_i - \bar{\theta}_{R,l}) \leq \liminf_{k \rightarrow +\infty} \frac{1}{M} \sum_{i=1}^M \sum_{l=1}^N \frac{\gamma_T}{2} TV(\theta_{T_i,l}^k) + \frac{\gamma_R}{2} TV(\theta_{R,l}^k) + \frac{\lambda}{2} TV(\theta_{T_i,l}^k \circ \varphi_i^k - \theta_{R,l}^k)$.
 155 Since from what precedes $\theta_{T_i,l}^k (c_{T_i,l}^k - T_i)^2 \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{T_i,l} (\bar{c}_{T_i,l} - T_i)^2$ almost everywhere in Ω and
 156 $\theta_{R,l}^k (c_{R,l}^k - T_i \circ \varphi_i^k)^2 \xrightarrow[k \rightarrow +\infty]{} \bar{\theta}_{R,l} (\bar{c}_{R,l} - T_i \circ \bar{\varphi}_i)^2$ almost everywhere in Ω with $|\theta_{T_i,l}^k (c_{T_i,l}^k - T_i)^2| \leq 4 \|T_i\|_{L^\infty(\Omega)}^2$ and $|\theta_{R,l}^k (c_{R,l}^k - T_i \circ \varphi_i^k)^2| \leq 4 \sum_{i=1}^M \|T_i\|_{L^\infty(\Omega)}^2$, we can apply the dominated convergence
 157 theorem and get $\int_{\Omega} \theta_{T_i,l}^k (c_{T_i,l}^k - T_i)^2 dx \xrightarrow[k \rightarrow +\infty]{} \int_{\Omega} \bar{\theta}_{T_i,l} (\bar{c}_{T_i,l} - T_i)^2 dx$ and $\int_{\Omega} \theta_{R,l}^k (c_{R,l}^k - T_i \circ \varphi_i^k)^2 dx \xrightarrow[k \rightarrow +\infty]{} \int_{\Omega} \bar{\theta}_{R,l} (\bar{c}_{R,l} - T_i \circ \bar{\varphi}_i)^2 dx$.
 158 By combining all the results, we get that

$$\begin{aligned}
 164 & \mathcal{F}_1(\bar{\theta}_R, \{\bar{\theta}_{T_i}, \bar{\varphi}_i\}_{i=1}^M) \leq \liminf_{k \rightarrow +\infty} \mathcal{F}_1(\theta_R^k, \{\theta_{T,i}^k, \varphi_i^k\}_{i=1}^M) < +\infty. \\
 165
 \end{aligned}$$

166 By finiteness of the functional, we deduce that $\bar{\theta}_R \in \mathcal{U}$, $\bar{\theta}_{T_i} \in \mathcal{U}$ for all $i \in \{1, \dots, M\}$ and
 167 $\bar{\varphi}_i \in \hat{\mathcal{W}}$ for all $i \in \{1, \dots, M\}$ with $\bar{\theta}_{T_i,l} \circ \bar{\varphi}_i - \bar{\theta}_{R,l} \in BV(\Omega)$ for all $l \in \{1, \dots, N\}$ and for all
 168 $i \in \{1, \dots, M\}$ and there exists a minimizer to our problem. \blacksquare

169 **SM2. Detailed proof of Theorem 3.1.**

170 *Proof.* Let $(\gamma_j)_{j \geq 0}$ be an increasing sequence of positive real numbers such that $\lim_{j \rightarrow +\infty} \gamma_j = +\infty$.
 171 Let $(\{\varphi_{i,k_j}, \theta_{T_i,k_j}, V_{i,k_j}, W_{i,k_j}\}_{i=1}^M, (\theta_{\tilde{T}_i,l,k_j})_{l=1,\dots,N}^{i=1,\dots,M}, \theta_{R,k_j})$ be a minimizing sequence of this
 172 problem $\mathcal{F}_{1,\gamma}$ for $\gamma = \gamma_j$. By definition, there exists $((\theta_{R,n}, \{\theta_{T_i,n}, \varphi_{i,n}\}_{i=1}^M) \in \mathcal{U} \times \mathcal{U}^M \times \hat{\mathcal{W}}^M$
 173 such that $\theta_{T_i,l,n} \circ \varphi_{i,n} - \theta_{R,l,n} \in BV(\Omega, \{-1, 0, 1\})$ for any $l \in \{1, \dots, N\}$ and all $i = 1, \dots, M$,
 174 and

$$\begin{aligned} 175 \quad & \mathcal{F}_1(\theta_{R,n}, \{\varphi_{i,n}, \theta_{T_i,n}\}_{i=1}^M) \\ 176 \quad & = \mathcal{F}_{1,\gamma_j}(\{\varphi_{i,n}, \theta_{T_i,n}, \nabla \varphi_{i,n}, (\nabla \varphi_{i,n})^{-1}\}_{i=1}^M, (\theta_{T_i,l,n} \circ \varphi_{i,n} - \theta_{R,l,n})_{l=1,\dots,N}^{i=1,\dots,M}, \theta_{R,n}) \\ 177 \quad & \leq \inf \mathcal{F}_1(\theta_R, \{\theta_{T_i}, \varphi_i\}_{i=1}^M) + \frac{1}{n} < +\infty. \end{aligned}$$

179 Then by definition of a minimizing sequence we get that

$$\begin{aligned} 180 \quad & \forall \varepsilon > 0, \exists N(\varepsilon, \gamma_j) \in \mathbb{N}, \forall k \in \mathbb{N}, k \geq N(\varepsilon, \gamma_j) \Rightarrow \mathcal{F}_{1,\gamma_j}(\{\varphi_{i,k_j}, \theta_{T_i,k_j}, V_{i,k_j}, W_{i,k_j}\}_{i=1}^M, \\ 181 \quad & (\theta_{\tilde{T}_i,l,k_j})_{l=1,\dots,N}^{i=1,\dots,M}, \theta_{R,k_j}) \leq \inf \mathcal{F}_{1,\gamma_j}(\{\varphi_i, \theta_{T_i}, V_i, W_i\}_{i=1}^M, (\theta_{\tilde{T}_i,l})_{l=1,\dots,N}^{i=1,\dots,M}, \theta_R) + \varepsilon, \\ 182 \quad & \leq \mathcal{F}_1(\theta_{R,n}, \{\theta_{T_i,n}, \varphi_{i,n}\}_{i=1}^M) + \varepsilon, \\ 183 \quad & = \mathcal{F}_{1,\gamma_j}(\{\varphi_{i,n}, \theta_{T_i,n}, \nabla \varphi_{i,n}, (\nabla \varphi_{i,n})^{-1}\}_{i=1}^M, (\theta_{T_i,l,n} \circ \varphi_{i,n} - \theta_{R,l,n})_{l=1,\dots,N}^{i=1,\dots,M}, \theta_{R,n}) + \varepsilon, \\ 184 \quad & \leq \inf \mathcal{F}_1(\theta_R, \{\theta_{T_i}, \varphi_i\}_{i=1}^M) + \frac{1}{n} + \varepsilon < +\infty. \end{aligned}$$

186 Let us set in particular $\varepsilon = \frac{1}{\gamma_j}$, then there exists $N_j \in \mathbb{N}$ such that for all $k \in \mathbb{N}, k \geq N_j \Rightarrow$

$$\begin{aligned} 187 \quad & \mathcal{F}_{1,\gamma_j}(\{\varphi_{i,k_j}, \theta_{T_i,k_j}, V_{i,k_j}, W_{i,k_j}\}_{i=1}^M, (\theta_{\tilde{T}_i,l,k_j})_{l=1,\dots,N}^{i=1,\dots,M}, \theta_{R,k_j}) \leq \inf \mathcal{F}_1(\theta_R, \{\theta_{T_i}, \varphi_i\}_{i=1}^M) + \frac{1}{n} + \frac{1}{\gamma_j} \\ 188 \quad & \leq \inf \mathcal{F}_1(\theta_R, \{\theta_{T_i}, \varphi_i\}_{i=1}^M) + \frac{1}{n} + \frac{1}{\gamma_0} \\ 189 \quad & < +\infty. \end{aligned}$$

191 So according to the previous inequality and by setting, for the sake of simplicity, for all
 192 $i = 1, \dots, M$,

$$\begin{aligned} 193 \quad & \varphi_{i,j} = \varphi_{i,N_j,j}, \\ 194 \quad & V_{i,j} = V_{i,N_j,j}, \\ 195 \quad & \theta_{T_i,j} = \theta_{T_i,N_j,j}, \\ 196 \quad & \theta_{R,j} = \theta_{R,N_j,j}, \\ 197 \quad & \theta_{\tilde{T}_i,l,j} = \theta_{\tilde{T}_i,l,N_j,j}, \\ 198 \quad & W_{i,j} = W_{i,N_j,j}, \end{aligned}$$

200 we get that

- $(\varphi_{i,j})$ is uniformly bounded according to j in $W^{1,4}(\Omega)$ by using the generalized Poincaré's inequality since $(V_{i,j})$ is uniformly bounded according to j in $L^4(\Omega, M_3(\mathbb{R}))$, for all $i = 1, \dots, M$.
- $(V_{i,j})$ is uniformly bounded according to j in $L^4(\Omega, M_3(\mathbb{R}))$ and in $L^\infty(\Omega, M_3(\mathbb{R}))$, for all $i = 1, \dots, M$.
- $(\theta_{T_i,l,j})$ is uniformly bounded according to j in $BV(\Omega)$ since $\|\theta_{T_i,l,j}\|_{L^1(\Omega)} \leq \text{meas}(\Omega) < +\infty$ and $(TV(\theta_{T_i,l,j}))$ is uniformly bounded according to j for all $l \in \{1, \dots, N\}$, and for all $i = 1, \dots, M$.
- $(\theta_{R,l,j})$ is uniformly bounded according to j in $BV(\Omega)$ since $\|\theta_{R,l,j}\|_{L^1(\Omega)} \leq \text{meas}(\Omega) < +\infty$ and $(TV(\theta_{R,l,j}))$ is uniformly bounded according to j for all $l \in \{1, \dots, N\}$.
- $(\theta_{\tilde{T}_i,l,j})$ is uniformly bounded according to j in $BV(\Omega)$ since $\forall l \in \{1, \dots, N\}$ and all $i = 1, \dots, M$, $\|\theta_{\tilde{T}_i,l,j}\|_{L^1(\Omega)} \leq \text{meas}(\Omega)$ and $(TV(\theta_{\tilde{T}_i,l,j}))$ is uniformly bounded according to j .
- $(\text{Cof}V_{i,j})$ is uniformly bounded according to j in $L^4(\Omega, M_3(\mathbb{R}))$ for all $i = 1, \dots, M$.
- $(\det V_{i,j})$ is uniformly bounded according to j in $L^2(\Omega)$ for all $i = 1, \dots, M$.
- $(W_{i,j})$ is uniformly bounded according to j in $L^\infty(\Omega, M_3(\mathbb{R}))$ and in $L^2(\Omega, M_3(\mathbb{R}))$ since $(V_{i,j})$ is uniformly bounded according to j in $L^4(\Omega, M_3(\mathbb{R}))$ and thus in $L^2(\Omega, M_3(\mathbb{R}))$, for all $i = 1, \dots, M$.

219 So, up to subsequences, we get that

- 220 $\varphi_{i,j} \xrightarrow[j \rightarrow +\infty]{} \bar{\varphi}_i$ in $W^{1,4}(\Omega, \mathbb{R}^3)$, for all $i = 1, \dots, M$,
- 221 $V_{i,j} \xrightarrow[j \rightarrow +\infty]{} \bar{V}_i$ in $L^4(\Omega, M_3(\mathbb{R}))$, and $V_{i,j} \xrightarrow[j \rightarrow +\infty]{*} \bar{V}_i$ in $L^\infty(\Omega, M_3(\mathbb{R}))$, for all $i = 1, \dots, M$,
by uniqueness of the weak limit in $L^4(\Omega, M_3(\mathbb{R}))$ and by the continuous embedding of
 $L^\infty(\Omega, M_3(\mathbb{R})) \subset L^4(\Omega, M_3(\mathbb{R}))$,
- 224 $\text{Cof}V_{i,j} \xrightarrow[j \rightarrow +\infty]{} \bar{X}_i$ in $L^4(\Omega, M_3(\mathbb{R}))$, for all $i = 1, \dots, M$,
- 225 $\det V_{i,j} \xrightarrow[j \rightarrow +\infty]{} \bar{\delta}_i$ in $L^2(\Omega)$, for all $i = 1, \dots, M$,
- 226 $W_{i,j} \xrightarrow[j \rightarrow +\infty]{} \bar{W}_i$ in $L^2(\Omega, M_3(\mathbb{R}))$, and $W_{i,j} \xrightarrow[j \rightarrow +\infty]{*} \bar{W}_i$ in $L^\infty(\Omega, M_3(\mathbb{R}))$, for all $i = 1, \dots, M$,
by uniqueness of the weak limit in $L^2(\Omega, M_3(\mathbb{R}))$ and by the continuous embedding of
 $L^\infty(\Omega, M_3(\mathbb{R})) \subset L^2(\Omega, M_3(\mathbb{R}))$,
- 229 $\theta_{T_i,l,j} \xrightarrow[j \rightarrow +\infty]{} \bar{\theta}_{T_i,l}$ in $L^1(\Omega)$, with $\bar{\theta}_{T_i,l} \in BV(\Omega)$, $\forall l \in \{1, \dots, N\}$, $\forall i \in \{1, \dots, M\}$,
- 230 $\theta_{R,l,j} \xrightarrow[j \rightarrow +\infty]{} \bar{\theta}_{R,l}$ in $L^1(\Omega)$, with $\bar{\theta}_{R,l} \in BV(\Omega)$, $\forall l \in \{1, \dots, N\}$,
- 231 $\theta_{\tilde{T}_i,l,j} \xrightarrow[j \rightarrow +\infty]{} \bar{\theta}_{\tilde{T}_i,l}$ in $L^1(\Omega)$, with $\bar{\theta}_{\tilde{T}_i,l} \in BV(\Omega)$, $\forall l \in \{1, \dots, N\}$, $\forall i \in \{1, \dots, M\}$.

Since the convergence is strong in $L^1(\Omega)$ then it is also almost everywhere up to subsequences and we deduce that $\bar{\theta}_{T_i,l} \in BV(\Omega, \{0, 1\})$, $\bar{\theta}_{R,l} \in BV(\Omega, \{0, 1\})$ and $\bar{\theta}_{\tilde{T}_i,l} \in BV(\Omega, \{-1, 0, 1\})$ for all $l \in \{1, \dots, N\}$ and all $i \in \{1, \dots, M\}$ and $\sum_{l=1}^M \bar{\theta}_{R,l}(x) = 1$ almost everywhere in Ω ,

$\sum_{l=1}^M \bar{\theta}_{T_i,l}(x) = 1$ almost everywhere in Ω for all $i = 1, \dots, M$, so that $\bar{\theta}_R \in \mathcal{U}$ and $\bar{\theta}_{T_i} \in \mathcal{U}$ for each $i = 1, \dots, M$.

Let us set $z_{i,j,l} = \theta_{\tilde{T}_i,l,j} - (\theta_{T_i,l,j} \circ \varphi_{i,j} - \theta_{R,l,j}) \xrightarrow{j \rightarrow +\infty} 0$ in $L^1(\Omega)$ for all $l \in \{1, \dots, N\}$ and all $i \in \{1, \dots, M\}$, $w_{i,j} = V_{i,j} - \nabla \varphi_{i,j} \xrightarrow{j \rightarrow +\infty} 0$ in $L^4(\Omega, M_3(\mathbb{R}))$ for all $i = 1, \dots, M$, and $y_{i,j} = W_{i,j} - V_{i,j}^{-1} \xrightarrow{j \rightarrow +\infty} 0$ in $L^2(\Omega, M_3(\mathbb{R}))$, for all $i = 1, \dots, M$.

Since strong convergence implies weak convergence, we have that

$$\forall i = 1, \dots, M, \forall \Phi \in L^{\frac{4}{3}}(\Omega, M_3(\mathbb{R})), \int_{\Omega} (V_{i,j} - \nabla \varphi_{i,j}) : \Phi dx \xrightarrow{j \rightarrow +\infty} 0.$$

But we also know that $\int_{\Omega} V_{i,j} : \Phi dx \xrightarrow{j \rightarrow +\infty} \int_{\Omega} \bar{V}_i : \Phi dx$, and we get $\nabla \varphi_{i,j} \xrightarrow{j \rightarrow +\infty} \bar{V}_i$ in $L^4(\Omega, M_3(\mathbb{R}))$, for all $i = 1, \dots, M$. By uniqueness of the weak limit in $L^4(\Omega, M_3(\mathbb{R}))$, we have that $\nabla \bar{\varphi}_i = \bar{V}_i \in L^\infty(\Omega, M_3(\mathbb{R}))$ for each $i = 1, \dots, M$.

We also have that

$$\forall i \in \{1, \dots, M\}, \forall \Phi \in L^2(\Omega, M_3(\mathbb{R})), \int_{\Omega} W_{i,j} - V_{i,j}^{-1} : \Phi dx \xrightarrow{j \rightarrow +\infty} 0.$$

233 But we know that $\int_{\Omega} W_{i,j} : \Phi dx \xrightarrow{j \rightarrow +\infty} \int_{\Omega} \bar{W}_i : \Phi dx$, and we get $V_{i,j}^{-1} \xrightarrow{j \rightarrow +\infty} \bar{W}_i$ in $L^2(\Omega, M_3(\mathbb{R}))$.

234 Let us now show that $\bar{W}_i = \bar{V}_i^{-1} = \nabla \bar{\varphi}_i^{-1}$, for all $i = 1, \dots, M$. We have that $\forall \Phi \in$
235 $L^2(\Omega, M_3(\mathbb{R}))$:

$$\begin{aligned} 236 \quad & \int_{\Omega} V_{i,j}^{-1} \bar{V}_i - I : \Phi dx = \int_{\Omega} V_{i,j}^{-1} \bar{V}_i - \bar{W}_i \bar{V}_i + \bar{W}_i \bar{V}_i - \bar{W}_i V_{i,j} + \bar{W}_i V_{i,j} - V_{i,j}^{-1} V_{i,j} : \Phi dx, \\ 237 \quad & = \int_{\Omega} (V_{i,j}^{-1} - \bar{W}_i) \bar{V}_i : \Phi dx + \int_{\Omega} \bar{W}_i (\bar{V}_i - V_{i,j}) : \Phi dx + \int_{\Omega} (\bar{W}_i - V_{i,j}^{-1}) V_{i,j} : \Phi dx, \\ 238 \quad & = \int_{\Omega} (V_{i,j}^{-1} - \bar{W}_i) : \Phi \bar{V}_i^T dx + \int_{\Omega} (\bar{V}_i - V_{i,j}) : \bar{W}_i^T \Phi dx + \int_{\Omega} (\bar{W}_i - V_{i,j}^{-1}) : \Phi V_{i,j}^T dx, \\ 239 \end{aligned}$$

240 with $\Phi \bar{V}_i^T \in L^2(\Omega, M_3(\mathbb{R}))$ since $\bar{V}_i \in L^\infty(\Omega, M_3(\mathbb{R}))$ so that $\int_{\Omega} (V_{i,j}^{-1} - \bar{W}_i) : \Phi \bar{V}_i^T dx \xrightarrow{j \rightarrow +\infty} 0$,

241 $\bar{W}_i^T \Phi \in L^2(\Omega, M_3(\mathbb{R})) \subset L^{\frac{4}{3}}(\Omega, M_3(\mathbb{R}))$ since $\bar{W}_i \in L^\infty(\Omega, M_3(\mathbb{R}))$ so that $\int_{\Omega} (V_{i,j}^{-1} - \bar{V}_i) : \Phi V_{i,j}^T dx \xrightarrow{j \rightarrow +\infty} 0$, and $\int_{\Omega} (\bar{W}_i - V_{i,j}^{-1}) : \Phi V_{i,j}^T dx \leq \alpha \int_{\Omega} (\bar{W}_i - V_{i,j}^{-1}) : \Phi' dx$ with $\Phi' \in L^2(\Omega)$

243 since $\forall j \in \mathbb{N}$, $\forall i \in \{1, \dots, M\}$, $\|V_{i,j}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha$ so that $\int_{\Omega} (\bar{W}_i - V_{i,j}^{-1}) : \Phi V_{i,j}^T dx \xrightarrow{j \rightarrow +\infty} 0$.

244 We therefore obtain that $V_{i,j}^{-1} \bar{V}_i \xrightarrow{j \rightarrow +\infty} I_3$ in $L^2(\Omega, M_3(\mathbb{R}))$ and since $V_{i,j}^{-1} \xrightarrow{j \rightarrow +\infty} \bar{W}_i$

245 in $L^2(\Omega, M_3(\mathbb{R}))$, we deduce that $\bar{W}_i \bar{V}_i = I_3$ and consequently $\bar{W}_i = \bar{V}_i^{-1} = (\nabla \bar{\varphi}_i)^{-1} \in$
246 $L^\infty(\Omega, M_3(\mathbb{R}))$, for all $i = 1, \dots, M$.

247 Now, we consider $V_{i,j} = w_{i,j} + \nabla \varphi_{i,j}$ and $\text{Cof}V_{i,j} = \text{Cof}(w_{i,j} + \nabla \varphi_{i,j})$.

248 For the purpose of illustration since the same reasoning applies to other indices, we focus on

249 the component of the first row, first column :

$$\begin{aligned}
 250 \quad (\text{Cof}V_{i,j})_{11} &= \left(w_{i,22} + \frac{\partial\varphi_{i,2}}{\partial x_2} \right) \left(w_{i,33} + \frac{\partial\varphi_{i,3}}{\partial x_3} \right) - \left(w_{i,23} + \frac{\partial\varphi_{i,2}}{\partial x_3} \right) \left(w_{i,32} + \frac{\partial\varphi_{i,3}}{\partial x_2} \right), \\
 251 \quad &= w_{i,22}w_{i,33} - w_{i,23}w_{i,32} + \frac{\partial\varphi_{i,3}}{\partial x_3} \frac{\partial\varphi_{i,2}}{\partial x_2} - \frac{\partial\varphi_{i,3}}{\partial x_2} \frac{\partial\varphi_{i,2}}{\partial x_3} + w_{i,22} \frac{\partial\varphi_{i,3}}{\partial x_3} + w_{i,33} \frac{\partial\varphi_{i,2}}{\partial x_2} \\
 252 \quad &- w_{i,23} \frac{\partial\varphi_{i,3}}{\partial x_2} - w_{i,32} \frac{\partial\varphi_{i,2}}{\partial x_3}, \\
 253 \quad &= (\text{Cof}w_{i,j})_{11} + (\text{Cof}\nabla\varphi_{i,j})_{11} + w_{i,22} \frac{\partial\varphi_{i,3}}{\partial x_3} + w_{i,33} \frac{\partial\varphi_{i,2}}{\partial x_2} - w_{i,23} \frac{\partial\varphi_{i,3}}{\partial x_2} - w_{i,32} \frac{\partial\varphi_{i,2}}{\partial x_3}.
 \end{aligned}$$

255 Let us set $d_{i,j} = (\text{Cof}w_{i,j})_{11} + w_{i,22} \frac{\partial\varphi_{i,3}}{\partial x_3} + w_{i,33} \frac{\partial\varphi_{i,2}}{\partial x_2} - w_{i,23} \frac{\partial\varphi_{i,3}}{\partial x_2} - w_{i,32} \frac{\partial\varphi_{i,2}}{\partial x_3} \in L^2(\Omega)$ by
256 generalized Hölder's inequality. Then using Hölder's inequality, one can prove that

$$\begin{aligned}
 257 \quad \|d_{i,j}\|_{L^2(\Omega)} &\leq \|w_{i,22}\|_{L^4(\Omega)} \|\frac{\partial\varphi_{i,3}}{\partial x_3}\|_{L^4(\Omega)} + \|w_{i,33}\|_{L^4(\Omega)} \|\frac{\partial\varphi_{i,2}}{\partial x_2}\|_{L^4(\Omega)} \\
 258 \quad &+ \|w_{i,23}\|_{L^4(\Omega)} \|\frac{\partial\varphi_{i,3}}{\partial x_2}\|_{L^4(\Omega)} + \|w_{i,32}\|_{L^4(\Omega)} \|\frac{\partial\varphi_{i,2}}{\partial x_3}\|_{L^4(\Omega)} \\
 259 \quad &+ \|w_{i,22}\|_{L^4(\Omega)} \|w_{i,33}\|_{L^4(\Omega)} + \|w_{i,23}\|_{L^4(\Omega)} \|w_{i,32}\|_{L^4(\Omega)}, \\
 260 \quad &\leq (\|w_{i,22}\|_{L^4(\Omega)}^4 + \|w_{i,33}\|_{L^4(\Omega)}^4 + \|w_{i,23}\|_{L^4(\Omega)}^4 + \|w_{i,32}\|_{L^4(\Omega)}^4)^{\frac{1}{4}} \\
 261 \quad &(\|\frac{\partial\varphi_{i,2}}{\partial x_3}\|_{L^4(\Omega)}^{\frac{4}{3}} + \|\frac{\partial\varphi_{i,3}}{\partial x_3}\|_{L^4(\Omega)}^{\frac{4}{3}} + \|\frac{\partial\varphi_{i,2}}{\partial x_2}\|_{L^4(\Omega)}^{\frac{4}{3}} + \|\frac{\partial\varphi_{i,3}}{\partial x_2}\|_{L^4(\Omega)}^{\frac{4}{3}})^{\frac{3}{4}} \\
 262 \quad &+ (\|w_{i,22}\|_{L^4(\Omega)}^4 + \|w_{i,33}\|_{L^4(\Omega)}^4 + \|w_{i,23}\|_{L^4(\Omega)}^4 + \|w_{i,32}\|_{L^4(\Omega)}^4)^{\frac{1}{4}} \\
 263 \quad &(\|w_{i,22}\|_{L^4(\Omega)}^{\frac{4}{3}} + \|w_{i,33}\|_{L^4(\Omega)}^{\frac{4}{3}} + \|w_{i,23}\|_{L^4(\Omega)}^{\frac{4}{3}} + \|w_{i,32}\|_{L^4(\Omega)}^{\frac{4}{3}})^{\frac{3}{4}}, \\
 264 \quad &\leq c_4 c_2^2 \|w_{i,j}\|_{L^4(\Omega, M_3(\mathbb{R}))} \|\nabla\varphi_{i,j}\|_{L^4(\Omega, M_3(\mathbb{R}))} + c_4 c_2^2 \|w_{i,j}\|_{L^4(\Omega, M_3(\mathbb{R}))}^2.
 \end{aligned}$$

Indeed, thanks to the property of equivalence of norms in finite dimension, there exist $(c_1, c_2, c_3, c_4) \in \mathbb{R}^4$ such that $\forall x \in \mathbb{R}^9$, $c_1\|x\|_2 \leq \|x\|_4 \leq c_2\|x\|_2$ and $c_3\|x\|_4 \leq \|x\|_{\frac{4}{3}} \leq c_4\|x\|_4 \leq c_4 c_2\|x\|_2$.

As $(\varphi_{i,j})$ is uniformly bounded according to j in $W^{1,4}(\Omega, \mathbb{R}^3)$ and as $w_{i,j} \xrightarrow[j \rightarrow +\infty]{} 0$ strongly in $L^4(\Omega)$, then

$$\|d_{i,j}\|_{L^2(\Omega)} \xrightarrow[j \rightarrow +\infty]{} 0,$$

and

$$\int_{\Omega} (\text{Cof}V_{i,j})_{11} \Phi \, dx = \int_{\Omega} d_{i,j} \Phi \, dx + \int_{\Omega} (\text{Cof}\nabla\varphi_{i,j})_{11} \Phi \, dx \xrightarrow[j \rightarrow +\infty]{} \int_{\Omega} (\text{Cof}\nabla\bar{\varphi}_i)_{11} \Phi \, dx, \forall \Phi \in L^2(\Omega),$$

266 $\forall i \in \{1, \dots, M\}$. Indeed, from [?, Theorem 8.20], as $\varphi_{i,j} \xrightarrow[j \rightarrow +\infty]{} \bar{\varphi}_i$ in $W^{1,4}(\Omega, \mathbb{R}^3)$, then
267 $\text{Cof}\nabla\varphi_{i,j} \xrightarrow[j \rightarrow +\infty]{} \text{Cof}\nabla\bar{\varphi}_i$ in $L^2(\Omega, M_3(\mathbb{R}))$. Thus $\text{Cof}V_{i,j} \xrightarrow[j \rightarrow +\infty]{} \text{Cof}\nabla\bar{\varphi}_i$ in $L^2(\Omega)$ and by
268 uniqueness of the weak limit in $L^2(\Omega)$, $\bar{X}_i = \text{Cof}\nabla\bar{\varphi}_i$ and $\text{Cof}V_{i,j} \xrightarrow[j \rightarrow +\infty]{} \text{Cof}\nabla\bar{\varphi}_i$ in $L^4(\Omega)$.

269 We then have from the generalized Hölder's inequality :

$$270 \quad \det V_{i,j} = V_{i,11}(\text{Cof } V_{i,j})_{11} + V_{i,21}(\text{Cof } V_{i,j})_{21} + V_{i,31}(\text{Cof } V_{i,j})_{31} \in L^{\frac{4}{3}}(\Omega), \\ 271 \quad = (w_{i,11} + \frac{\partial \varphi_{i,1}}{\partial x_1})((\text{Cof } \nabla \varphi_{i,j})_{11} + d_{i,j}) + \text{ similar components}.$$

273 Then $\forall \Phi \in L^4(\Omega)$, and for all $i = 1, \dots, M$,

$$274 \quad \int_{\Omega} \det V_{i,j} : \Phi \, dx = \int_{\Omega} w_{i,11}(\text{Cof } \nabla \varphi_{i,j})_{11} \Phi \, dx + \int_{\Omega} w_{i,11} d_{i,j} \Phi \, dx + \int_{\Omega} \frac{\partial \varphi_{i,1}}{\partial x_1} d_{i,j} \Phi \, dx \\ 275 \quad + \int_{\Omega} \frac{\partial \varphi_{i,1}}{\partial x_1} (\text{Cof } \nabla \varphi_{i,j})_{11} \Phi \, dx + \text{ similar components}.$$

277 But we have for all $i = 1, \dots, M$,

$$278 \quad \left| \int_{\Omega} w_{i,11}(\text{Cof } \nabla \varphi_{i,j})_{11} \Phi \, dx \right| \leq \|w_{i,11}\|_{L^4(\Omega)} \|\text{Cof } \nabla \varphi_{i,j}\|_{L^2(\Omega)} \|\Phi\|_{L^4(\Omega)}, \\ 279 \quad \left| \int_{\Omega} w_{i,11} d_{i,j} \Phi \, dx \right| \leq \|w_{i,11}\|_{L^4(\Omega)} \|d_{i,j}\|_{L^2(\Omega)} \|\Phi\|_{L^4(\Omega)}, \\ 280 \quad \left| \int_{\Omega} \frac{\partial \varphi_{i,1}}{\partial x_1} d_{i,j} \Phi \, dx \right| \leq \left\| \frac{\partial \varphi_{i,1}}{\partial x_1} \right\|_{L^4(\Omega)} \|d_{i,j}\|_{L^2(\Omega)} \|\Phi\|_{L^4(\Omega)},$$

282 with $\|w_{i,11}\|_{L^4(\Omega)} \xrightarrow{j \rightarrow +\infty} 0$, $\|d_{i,j}\|_{L^2(\Omega)} \xrightarrow{j \rightarrow +\infty} 0$ and $\|\text{Cof } \nabla \varphi_{i,j}\|_{L^2(\Omega)}$ uniformly bounded since
283 $\text{Cof } \nabla \varphi_{i,j} \xrightarrow{j \rightarrow +\infty} \text{Cof } \nabla \bar{\varphi}_i$ in $L^4(\Omega, M_3(\mathbb{R}))$ and thus in $L^2(\Omega, M_3(\mathbb{R}))$, $\|\frac{\partial \varphi_{i,1}}{\partial x_1}\|_{L^4(\Omega)}$ uniformly
284 bounded since $\varphi_{i,j} \xrightarrow{j \rightarrow +\infty} \bar{\varphi}_i$ in $W^{1,4}(\Omega)$ and $\|\Phi\|_{L^4(\Omega)}$ uniformly bounded.

285 Since $\varphi_{i,j} \xrightarrow{j \rightarrow +\infty} \bar{\varphi}_i$ in $W^{1,4}(\Omega, \mathbb{R}^3)$, then $\det \nabla \varphi_{i,j} \xrightarrow{j \rightarrow +\infty} \det \nabla \bar{\varphi}_i$ in $L^{\frac{4}{3}}(\Omega)$ and $\int_{\Omega} \frac{\partial \varphi_{i,1}}{\partial x_1}$
286 $(\text{Cof } \nabla \varphi_{i,j})_{11} \Phi \, dx \xrightarrow{j \rightarrow +\infty} \int_{\Omega} \frac{\partial \bar{\varphi}_{i,1}}{\partial x_1} (\text{Cof } \nabla \bar{\varphi}_i)_{11} \Phi \, dx$, for all $i = 1, \dots, M$.

287 Finally, $\int_{\Omega} \det V_{i,j} \Phi \, dx \xrightarrow{j \rightarrow +\infty} \int_{\Omega} \det \nabla \bar{\varphi}_i \Phi \, dx$, $\forall \Phi \in L^4(\Omega)$ and so $\det V_{i,j} \xrightarrow{j \rightarrow +\infty} \det \nabla \bar{\varphi}_i$ in
288 $L^{\frac{4}{3}}(\Omega)$ and by uniqueness of the weak limit, $\bar{\delta}_i = \det \nabla \bar{\varphi}_i$ and $\det V_{i,j} \xrightarrow{j \rightarrow +\infty} \det \nabla \bar{\varphi}_i$ in $L^2(\Omega)$,
289 for all $i = 1, \dots, M$.

290

We know that W_{Op} is convex and continuous. If $\psi_n \xrightarrow{n \rightarrow +\infty} \psi$ in $L^4(\Omega, M_3(\mathbb{R}))$ and we can extract a subsequence still denoted (ψ_n) such that $\nabla \psi_n \xrightarrow{n \rightarrow +\infty} \nabla \psi$ almost everywhere on Ω . If $\alpha_n \xrightarrow{n \rightarrow +\infty} \bar{\alpha}$ in $L^4(\Omega, M_3(\mathbb{R}))$, then we can extract a subsequence still denoted (α_n) such that $\alpha_n \xrightarrow{n \rightarrow +\infty} \bar{\alpha}$ almost everywhere on Ω . If $\delta_n \xrightarrow{n \rightarrow +\infty} \bar{\delta}$ in $L^2(\Omega)$, then we can extract a subsequence still denoted (δ_n) such that $\delta_n \xrightarrow{n \rightarrow +\infty} \bar{\delta}$ almost everywhere on Ω . Then by continuity of W_{Op} , we get that $W_{Op}(\psi_n, \alpha_n, \delta_n) \xrightarrow{j \rightarrow +\infty} W_{Op}(\psi, \bar{\alpha}, \bar{\delta})$ almost everywhere on Ω . Then by applying Fatou's lemma, we have that $\liminf_{n \rightarrow +\infty} \int_{\Omega} W_{Op}(\psi_n, \alpha_n, \delta_n) \, dx \geq \int_{\Omega} W_{Op}(\psi, \bar{\alpha}, \bar{\delta}) \, dx$. Since W_{Op} is convex, so is $\int_{\Omega} W_{Op}(\xi, \alpha, \delta) \, dx$ and [?, Corollaire III.8 p.38] applies. Therefore

$\int_{\Omega} W_{Op}(\xi, \alpha, \delta) dx$ is also weakly lower semicontinuous in $L^4(\Omega, M_3(\mathbb{R})) \times L^4(\Omega, M_3(\mathbb{R})) \times L^2(\Omega)$ and we deduce that $+\infty > \liminf_{j \rightarrow +\infty} \int_{\Omega} W_{Op}(V_{i,j}, \text{Cof}V_{i,j}, \det V_{i,j}) dx \geq \int_{\Omega} W_{Op}(\nabla \bar{\varphi}_i, \text{Cof} \nabla \bar{\varphi}_i, \det \nabla \bar{\varphi}_i) dx$.

Using the same arguments as in the previous proof, we get more regularity of $\bar{\varphi}_i$, that is to say, $\bar{\varphi}_i$ is a bi-Lipschitz homeomorphism for each $i = 1, \dots, M$.

By the weak-* lower semicontinuity of $\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))}$, we have that $\|\nabla \bar{\varphi}_i\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \liminf_{j \rightarrow +\infty} \|V_{i,j}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha$, $\|(\nabla \bar{\varphi}_i)^{-1}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \liminf_{j \rightarrow +\infty} \|W_{i,j}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta$ and

$$\mathbf{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha\}}(\nabla \bar{\varphi}_i) \leq \liminf_{j \rightarrow +\infty} \mathbf{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha\}}(V_{i,j}), \mathbf{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta\}}((\nabla \bar{\varphi}_i)^{-1}) \leq \liminf_{j \rightarrow +\infty} \mathbf{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta\}}(W_{i,j}).$$

We also know that $\varphi_{i,j} \xrightarrow{j \rightarrow +\infty} \bar{\varphi}_i$ in $C^{0,\alpha}(\Omega)$, $\bar{\varphi}_i^{-1} \in C^{0,\alpha'}(\Omega)$ and $\varphi_{i,j} \circ \bar{\varphi}_i^{-1} \xrightarrow{j \rightarrow +\infty} \text{Id}$ in $C^{0,\alpha''}(\Omega)$

with $\alpha'' = \alpha'\alpha$, $\alpha < \frac{1}{4}$ and $\alpha' < 1 - \frac{3}{q}$, using the same arguments as in the previous proof. Let us prove that $\theta_{T_i,l,j} \circ \varphi_{i,j} \xrightarrow{j \rightarrow +\infty} \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i$ in $L^1(\Omega)$ for all $i = 1, \dots, M$.

Since $\varphi_{i,j} \circ \bar{\varphi}_i^{-1} \xrightarrow{j \rightarrow +\infty} \text{Id}$ in $C^{0,\alpha''}(\Omega)$, then it converges uniformly and so

$$\forall \varepsilon > 0, \exists k \in \mathbb{N}, \forall j \in \mathbb{N}, j \geq k \Rightarrow \sup_{x \in \Omega} |\varphi_{i,j} \circ \bar{\varphi}_i^{-1}(x) - x| \leq \varepsilon.$$

- 291 Also given $\eta > 0$, by Egorov's theorem, there exists a set S_η with Lebesgue measure $\text{meas}(S_\eta) \leq \eta$
292 such that $\theta_{T_i,l,j} \xrightarrow{j \rightarrow +\infty} \bar{\theta}_{T_i,l}$ uniformly on $\Omega \setminus S_\eta$.
293 Now we choose $K \in \mathbb{N}$ large enough such that $\forall j \in \mathbb{N}$, if $j \geq K$, then

$$294 \quad |\varphi_{i,j} \circ \bar{\varphi}_i^{-1}(x) - x| \leq \eta, \text{ on } \Omega, \\ 295 \quad |\theta_{T_i,l,j}(x) - \bar{\theta}_{T_i,l}(x)| \leq \eta, \text{ on } \Omega \setminus S_\eta.$$

- 297 Then by denoting $\mathcal{O}_i = \{x \in \Omega : \bar{\theta}_{T_i,l}(x) = 1\}$,

$$298 \quad \|\theta_{T_i,l,j} \circ \varphi_{i,j} - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i\|_{L^1(\Omega)} = \int_{\bar{\varphi}_i^{-1}(\mathcal{O}_i)} |\theta_{T_i,l,j} \circ \varphi_{i,j}(x) - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i(x)| dx \\ 299 \quad + \int_{\bar{\varphi}_i^{-1}(\Omega \setminus \mathcal{O}_i)} |\theta_{T_i,l,j} \circ \varphi_{i,j}(x) - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i(x)| dx, \\ 300 \quad = \int_{\mathcal{O}_i} \frac{|\theta_{T_i,l,j} \circ \varphi_{i,j} \circ \bar{\varphi}_i^{-1}(x) - 1|}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx + \int_{\Omega \setminus \mathcal{O}_i} \frac{|\theta_{T_i,l,j} \circ \varphi_{i,j} \circ \bar{\varphi}_i^{-1}(x)|}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx, \\ 301 \quad \leq \int_{(\mathcal{O}_i \setminus S_\eta)_\eta} \frac{|\theta_{T_i,l,j} \circ \varphi_{i,j} \circ \bar{\varphi}_i^{-1}(x) - 1|}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx + \int_{((\Omega \setminus \mathcal{O}_i) \setminus S_\eta)_\eta} \frac{|\theta_{T_i,l,j} \circ \varphi_{i,j} \circ \bar{\varphi}_i^{-1}(x)|}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx \\ 302 \quad + \int_{S_\eta} \frac{|\theta_{T_i,l,j} \circ \varphi_{i,j} \circ \bar{\varphi}_i^{-1}(x) - \bar{\theta}_{T_i,l}(x)|}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx + \int_{\mathcal{O}_i^\eta \cup S_\eta^\eta \cup \Omega^\eta} \frac{|\theta_{T_i,l,j} \circ \varphi_{i,j} \circ \bar{\varphi}_i^{-1}(x) - \bar{\theta}_{T_i,l}(x)|}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx,$$

- 304 with $(\mathcal{O}_i \setminus S_\eta)_\eta = \{x \in \mathcal{O}_i \setminus S_\eta \mid \text{dist}(x, \partial(\mathcal{O}_i \setminus S_\eta)) > \eta\}$, $S_\eta^\eta = \{x \in \Omega \mid \text{dist}(x, \partial S_\eta) \leq \eta\}$. So
305 $\forall x \in (\mathcal{O}_i \setminus S_\eta)_\eta$, $\varphi_{i,j} \circ \bar{\varphi}_i^{-1}(x) \in \mathcal{O}_i \setminus S_\eta$, and $\forall x \in ((\Omega \setminus \mathcal{O}_i) \setminus S_\eta)_\eta$, $\varphi_{i,j} \circ \bar{\varphi}_i^{-1}(x) \in (\Omega \setminus \mathcal{O}_i) \setminus S_\eta$.

306 Then

$$\begin{aligned}
 307 \quad & \| \theta_{T_i,l,j} \circ \varphi_{i,j} - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i \|_{L^1(\Omega)} \leq \int_{(\mathcal{O}_i \setminus S_\eta)_\eta} \frac{\eta}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx \\
 308 \quad & + \int_{((\Omega \setminus \mathcal{O}_i) \setminus S_\eta)_\eta} \frac{\eta}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx \\
 309 \quad & + \int_{S_\eta} \frac{1}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx \\
 310 \quad & + \int_{\mathcal{O}_i^\eta \cup S_\eta^\eta \cup \Omega^\eta} \frac{1}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))} dx,
 \end{aligned}$$

312 taking into account that $\theta_{T_i,l,j} \in BV(\Omega, \{0, 1\})$ and $\bar{\theta}_{T_i,l} \in BV(\Omega, \{0, 1\})$. Using Hölder's
313 inequality, we get

$$\begin{aligned}
 314 \quad & \| \theta_{T_i,l,j} \circ \varphi_{i,j} - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i \|_{L^1(\Omega)} \leq \eta \left(\int_{\Omega} dx \right)^{\frac{s}{s+1}} \left(\int_{\Omega} \frac{1}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))^{s+1}} dx \right)^{\frac{1}{s+1}} \\
 315 \quad & + \eta \left(\int_{\Omega} dx \right)^{\frac{s}{s+1}} \left(\int_{\Omega} \frac{1}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))^{s+1}} dx \right)^{\frac{1}{s+1}} \\
 316 \quad & + \left(\int_{S_\eta} dx \right)^{\frac{s}{s+1}} \left(\int_{\Omega} \frac{1}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))^{s+1}} dx \right)^{\frac{1}{s+1}} \\
 317 \quad & + \left(\int_{S_\eta^\eta \cup \mathcal{O}_i^\eta \cup \Omega^\eta} dx \right)^{\frac{s}{s+1}} \left(\int_{\Omega} \frac{1}{\det \nabla \bar{\varphi}_i(\bar{\varphi}_i^{-1}(x))^{s+1}} dx \right)^{\frac{1}{s+1}}, \\
 318 \quad & \| \theta_{T_i,l,j} \circ \varphi_{i,j} - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i \|_{L^1(\Omega)} \leq \eta (\text{meas}(\Omega))^{\frac{s}{s+1}} \left(\int_{\Omega} \frac{1}{\det \nabla \bar{\varphi}_i(x)^s} dx \right)^{\frac{1}{s+1}} \\
 319 \quad & + \eta (\text{meas}(\Omega))^{\frac{s}{s+1}} \left(\int_{\Omega} \frac{1}{\det \nabla \bar{\varphi}_i(x)^s} dx \right)^{\frac{1}{s+1}} \\
 320 \quad & + (\text{meas}(S_\eta))^{\frac{s}{s+1}} \left(\int_{\Omega} \frac{1}{\det \nabla \bar{\varphi}_i(x)^s} dx \right)^{\frac{1}{s+1}} \\
 321 \quad & + (\text{meas}(S_\eta^\eta) + \text{meas}(\mathcal{O}_i^\eta) + \text{meas}(\Omega^\eta))^{\frac{s}{s+1}} \left(\int_{\Omega} \frac{1}{\det \nabla \bar{\varphi}_i(x)^s} dx \right)^{\frac{1}{s+1}}. \\
 322 \quad & \blacksquare
 \end{aligned}$$

324 With $s = 10$ and the finiteness of $\int_{\Omega} W_{Op}(\nabla \bar{\varphi}_i, \text{Cof} \nabla \bar{\varphi}_i, \det \nabla \bar{\varphi}_i) dx$, there exists $C \in \mathbb{R}_*^+$ such
325 that

$$\begin{aligned}
 326 \quad & \| \theta_{T_i,l,j} \circ \varphi_{i,j} - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i \|_{L^1(\Omega)} \leq \eta (\text{meas}(\Omega))^{\frac{s}{s+1}} C^{\frac{1}{s+1}} + \eta (\text{meas}(\Omega))^{\frac{s}{s+1}} C^{\frac{1}{s+1}} + (\text{meas}(S_\eta))^{\frac{s}{s+1}} \\
 327 \quad & C^{\frac{1}{s+1}} + \left(\int_{\partial S_\eta} \int_{B(x,\eta)} dy dx + \int_{\partial \mathcal{O}_i} \int_{B(x,\eta)} dy dx + \int_{\partial \Omega} \int_{B(x,\eta)} dy dx \right)^{\frac{s}{s+1}} \\
 328 \quad & C^{\frac{1}{s+1}}, \quad \blacksquare
 \end{aligned}$$

$$\begin{aligned}
 331 \quad & \|\theta_{T_i,l,j} \circ \varphi_{i,j} - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i\|_{L^1(\Omega)} \leq \eta (\text{meas}(\Omega))^{\frac{s}{s+1}} C^{\frac{1}{s+1}} \\
 332 \quad & + \eta (\text{meas}((\Omega))^{\frac{s}{s+1}} C^{\frac{1}{s+1}} \\
 333 \quad & + (\eta)^{\frac{s}{s+1}} C^{\frac{1}{s+1}} \\
 334 \quad & + \left(\frac{4\pi}{3} \eta^3 (\text{meas}(\partial S_\eta) + \text{meas}(\partial \mathcal{O}_i) + \text{meas}(\partial \Omega)) \right)^{\frac{s}{s+1}} C^{\frac{1}{s+1}}. \\
 335
 \end{aligned}$$

336 By letting η tend to 0 and so j to infinity, we finally get for all $i = 1, \dots, M$

$$337 \quad \|\theta_{T_i,l,j} \circ \varphi_{i,j} - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i\|_{L^1(\Omega)} \xrightarrow{j \rightarrow +\infty} 0. \\
 338$$

339 This is true for all $l \in \{1, \dots, N\}$.

340 Finally, for all $i = 1, \dots, M$ and all $l = 1, \dots, N$:

$$\begin{aligned}
 341 \quad & \|\bar{\theta}_{\tilde{T}_i,l} - \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i + \bar{\theta}_{R,l}\|_{L^1(\Omega)} \leq \|\theta_{\tilde{T}_i,l,j} - \theta_{T_i,l,j} \circ \varphi_{i,j} + \theta_{R,l,j}\|_{L^1(\Omega)} \\
 342 \quad & + \|\bar{\theta}_{\tilde{T}_i,l} - \theta_{\tilde{T}_i,l,j}\|_{L^1(\Omega)} + \|\bar{\theta}_{T_i,l} \circ \bar{\varphi}_i - \theta_{T_i,l,j} \circ \varphi_{i,j}\|_{L^1(\Omega)} \\
 343 \quad & + \|\bar{\theta}_{R,l} - \theta_{R,l,j}\|_{L^1(\Omega)}, \\
 344 \quad & \xrightarrow{j \rightarrow +\infty} 0, \\
 345
 \end{aligned}$$

346 from what precedes. We deduce that $\bar{\theta}_{\tilde{T}_i,l} = \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i - \bar{\theta}_{R,l}$ in $L^1(\Omega)$ and so almost everywhere
347 in Ω and $\theta_{\tilde{T}_i,l,j} \xrightarrow{j \rightarrow +\infty} \bar{\theta}_{T_i,l} \circ \bar{\varphi}_i - \bar{\theta}_{R,l}$ in $L^1(\Omega)$ for all $i = 1, \dots, M$ and all $l = 1, \dots, N$. By
348 the semicontinuity of the total variation we get that

$$349 \quad \frac{\lambda}{2} \sum_{l=1}^N TV(\bar{\theta}_{T_i,l} \circ \bar{\varphi}_i - \bar{\theta}_{R,l}) \leq \frac{\lambda}{2} \liminf_{j \rightarrow +\infty} \sum_{l=1}^N TV(\theta_{\tilde{T}_i,l,j}). \\
 350$$

351 Furthermore, we have that $\theta_{T_i,l,j} \xrightarrow{j \rightarrow +\infty} \bar{\theta}_{T_i,l}$ in $L^1(\Omega)$ and so almost everywhere in Ω up
352 to a subsequence. Then $\theta_{T_i,l,j} T_i \xrightarrow{j \rightarrow +\infty} \bar{\theta}_{T_i,l} T_i$ almost everywhere in Ω with $|\theta_{T_i,l,j} T_i| \leq$
353 $\|T_i\|_{L^\infty(\Omega)} \in L^1(\Omega)$ since Ω is bounded, and by the dominated convergence theorem one
354 gets $\int_{\Omega} \theta_{T_i,l,j} T_i dx \xrightarrow{j \rightarrow +\infty} \int_{\Omega} \bar{\theta}_{T_i,l} T_i dx$. Using the same reasoning, we get $\int_{\Omega} \theta_{T_i,l,j} dx \xrightarrow{j \rightarrow +\infty}$
355 $\int_{\Omega} \bar{\theta}_{T_i,l} dx$. As $T_i \in W^{1,\infty}(\Omega)$, we denote κ_i the Lipschitz constant related to T_i . Then the chain
356 rule applies and we have that $T_i \circ \varphi_i^k \in W^{1,4}(\Omega)$, $T_i \circ \bar{\varphi}_i \in W^{1,4}(\Omega)$ and $\|T_i \circ \varphi_{i,j} - T_i \circ \bar{\varphi}_i\|_{C^0(\Omega)} \leq$
357 $\kappa_i \|\varphi_{i,j} - \bar{\varphi}_i\|_{C^0(\Omega)} \xrightarrow{j \rightarrow +\infty} 0$ using the Sobolev compact embedding $W^{1,4}(\Omega) \hookrightarrow C^0(\Omega)$. Therefore,
358 up to a subsequence $\theta_{R,l,j} T_i \circ \varphi_{i,j} \xrightarrow{j \rightarrow +\infty} \bar{\theta}_{R,l} T_i \circ \bar{\varphi}_i$ almost everywhere in Ω , $\theta_{R,l,j} \xrightarrow{j \rightarrow +\infty} \bar{\theta}_{R,l}$
359 almost everywhere in Ω with $|\theta_{R,l,j} T_i \circ \varphi_{i,j}| \leq \|T_i\|_{L^\infty(\Omega)} \in L^1(\Omega)$ and $|\theta_{R,l,j}| \leq 1 \in L^1(\Omega)$

360 almost everywhere in Ω . Thus the dominated convergence theorem applies and we get

$$361 \quad \frac{1}{M} \sum_{i=1}^M \int_{\Omega} \theta_{R,l,j} T_i \circ \varphi_{i,j} dx \xrightarrow{j \rightarrow +\infty} \frac{1}{M} \sum_{i=1}^M \int_{\Omega} \bar{\theta}_{R,l} T_i \circ \bar{\varphi}_i dx,$$

$$362 \quad \int_{\Omega} \theta_{R,l,j} dx \xrightarrow{k \rightarrow +\infty} \int_{\Omega} \bar{\theta}_{R,l} dx,$$

$$363 \quad c_{R,l,j} \xrightarrow{k \rightarrow +\infty} \bar{c}_{R,l}.$$

365

366 Then $\forall l \in \{1, \dots, N\}$ and for all $i = 1, \dots, M$,

$$367 \quad \theta_{T_i,l,j}(c_{T_i,l,j} - T_i)^2 \xrightarrow{j \rightarrow +\infty} \bar{\theta}_{T_i,l}(\bar{c}_{T_i,l} - T_i)^2,$$

$$368 \quad \theta_{R,l,j}(c_{R,l,j} - R)^2 \xrightarrow{j \rightarrow +\infty} \bar{\theta}_{R,l}(\bar{c}_{R,l} - R)^2,$$

370 almost everywhere on Ω with $|\theta_{T_i,l,j}(c_{T_i,l,j} - T_i)^2| \leq 4\|T_i\|_{L^\infty(\Omega)}^2 \in L^1(\Omega)$ and $|\theta_{R,l,j}(c_{R,l,j} - R)^2| \leq 4\|T_i\|_{L^\infty(\Omega)}^2 \in L^1(\Omega)$. By applying the dominated convergence theorem, we have for all $l \in \{1, \dots, N\}$:

$$373 \quad \|\theta_{T_i,l,j}(c_{T_i,l,j} - T_i)^2\|_{L^1(\Omega)} \xrightarrow{j \rightarrow +\infty} \|\bar{\theta}_{T_i,l}(\bar{c}_{T_i,l} - T_i)^2\|_{L^1(\Omega)},$$

$$374 \quad \|\theta_{R,l,j}(c_{R,l,j} - R)^2\|_{L^1(\Omega)} \xrightarrow{j \rightarrow +\infty} \|\bar{\theta}_{R,l}(\bar{c}_{R,l} - R)^2\|_{L^1(\Omega)}.$$

375 By combining all the results, we finally get

$$377 \quad \frac{1}{M} \sum_{i=1}^M \frac{\gamma_T}{2} \sum_{l=1}^N TV(\bar{\theta}_{T_i,l}) + \int_{\Omega} \sum_{l=1}^N \bar{\theta}_{T_i,l}(\bar{c}_{T_i,l} - T_i)^2 dx \\ 378 \quad + \frac{\gamma_R}{2} \sum_{l=1}^N TV(\bar{\theta}_{R,l}) + \int_{\Omega} \sum_{l=1}^N \bar{\theta}_{R,l}(\bar{c}_{R,l} - R)^2 dx \\ 379 \quad + \frac{\lambda}{2} \sum_{l=1}^N TV(\bar{\theta}_{T_i,l} \circ \bar{\varphi}_i - \bar{\theta}_{R,l}) + \int_{\Omega} W_{Op}(\nabla \bar{\varphi}_i, \text{Cof} \nabla \bar{\varphi}_i, \det \nabla \bar{\varphi}_i) dx \\ 380 \quad + \mathbb{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha\}}(\nabla \bar{\varphi}_i) + \mathbb{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta\}}((\nabla \bar{\varphi}_i)^{-1}) \\ 381 \quad \leq \liminf_{j \rightarrow +\infty} \mathcal{F}_{1,\gamma_j}(\{\varphi_{i,j}, \theta_{T_i,j}, V_{i,j}, W_{i,j}\}_{i=1}^M, (\theta_{T_i,l,j})_{l=1, \dots, N}^{i=1, \dots, M}, \theta_{R,j}), \\ 382 \quad \leq \limsup_{j \rightarrow +\infty} \mathcal{F}_{1,\gamma_j}(\{\varphi_{i,j}, \theta_{T_i,j}, V_{i,j}, W_{i,j}\}_{i=1}^M, (\theta_{T_i,l,j})_{l=1, \dots, N}^{i=1, \dots, M}, \theta_{R,j}), \\ 383 \quad \leq \limsup_{j \rightarrow +\infty} \inf \mathcal{F}_1(\theta_R, \{\theta_{T_i}, \varphi_i\}_{i=1}^M) + \frac{2}{\gamma_j} = \inf \mathcal{F}_i(\theta_R, \{\theta_{T_i}, \varphi_i\}_{i=1}^M).$$

384 By finiteness of $\int_{\Omega} W_{Op}(\nabla \bar{\varphi}_i, \text{Cof} \nabla \bar{\varphi}_i, \det \nabla \bar{\varphi}_i) dx$ and by continuity of the trace operator, we have that $\bar{\varphi}_i \in \hat{\mathcal{W}}$. By strong convergence in $L^1(\Omega)$ and so almost everywhere, we get that $\bar{\theta}_{T_i} \in \mathcal{U}$ and $\bar{\theta}_R \in \mathcal{U}$ with $\bar{\theta}_{T_i,l} \circ \bar{\varphi}_i - \bar{\theta}_{R,l} \in BV(\Omega, \{-1, 0, 1\})$ for all $l \in \{1, \dots, N\}$ and all $i \in \{1, \dots, N\}$. Therefore our decoupled problem (DP) converges to the initial problem (P) as γ tends to infinity. \blacksquare

Algorithm SM3.1 Alternating scheme of resolution.

1. Define $k := 1$, $T_i := i$ -th template image, $\theta_{T_i} :=$ Potts segmentation of T_i ,

$$\theta_R := \text{Potts segmentation of } \frac{1}{M} \sum_{i=1}^M T_i, \quad \theta_{\tilde{T}_i} := \theta_{T_i} - \theta_R, \quad V_i = \begin{pmatrix} V_{i,11} & V_{i,12} \\ V_{i,21} & V_{i,22} \end{pmatrix} := I, \quad W_i := \begin{pmatrix} W_{i,11} & W_{i,12} \\ W_{i,21} & W_{i,22} \end{pmatrix} := I, \quad a_1, a_2, a_3, \lambda_T, \lambda_R, \gamma_T, \gamma_R, \gamma_1, \gamma_2, \gamma_3, nbIter, \alpha, \beta, U_i = (U_{i,1}, U_{i,2}) :=$$

0, displacements associated to the deformation φ_i , for $i = 1, \dots, M$.

while $k < nbIter$ **do**

if $k \% 10 == 0$ **then**

 2.1. Compute for $i = 1, \dots, M$ the inverse deformation φ_i^{-1} using a Delaunay triangulation and linear interpolation.

 2.2. For $i = 1, \dots, M$, update θ_{T_i} by solving the Potts model with algorithm in [?]:

$$\inf_u \gamma_T \|\nabla u\|_{L^0(\Omega)} + \|\sqrt{\lambda_T + (\det \nabla \varphi_i)^{-1} \frac{\gamma_1}{2}} u - \frac{\lambda_T T_i + \frac{\gamma_1}{2} (\det \nabla \varphi_i)^{-1} (\theta_{\tilde{T}_i} \circ \varphi_i^{-1} + \theta_R \circ \varphi_i^{-1})}{\sqrt{\lambda_T + (\det \nabla \varphi_i)^{-1} \frac{\gamma_1}{2}}} \|_{L^2(\Omega)}^2 \text{ end for 2.2.}$$

 2.3. For $i = 1, \dots, M$, update $\theta_{\tilde{T}_i}$ by solving the Potts model with algorithm in [?]:

$$\inf_u \gamma_{\tilde{T}_i} \|\nabla u\|_{L^0(\Omega)} + \frac{\gamma_1}{2} \|(\theta_{T_i} \circ \varphi_i - \theta_R) - u\|_{L^2(\Omega)}^2 \text{ end for 2.3.}$$

 2.4. Update θ_R by solving the Potts model with algorithm in [?]: $\inf_u \gamma_R \|\nabla u\|_{L^0(\Omega)} +$

$$(\lambda_R + \frac{\gamma_1}{2}) \|u - \frac{\frac{\lambda_R}{M} \sum_{i=1}^M T_i \circ \varphi_i + \frac{\gamma_1}{2M} \sum_{i=1}^M (\theta_{T_i} \circ \varphi - \theta_{\tilde{T}_i})}{\lambda_R + \frac{\gamma_1}{2}}\|_{L^2(\Omega)}^2.$$

end if

2.5. For each $i = 1, \dots, M$, for each pixel (l, j) , — c playing a role similar to the one of a step size in a gradient method— update V_i using the following equations:

$$\left\{ \begin{array}{l} \text{temp}_1(l, j) = V_{i,11}(l, j) + c \left(\frac{10a_3}{(\det V_i(l, j))^{11}} V_{i,22}(l, j) - 4a_1 V_{i,11}(l, j) \right. \\ \quad \left. \|V_i(l, j)\|^2 - 2a_2(\det V_i(l, j) - 1)V_{i,22}(l, j) + \gamma_2 \left(\frac{\partial \varphi_{i,1}}{\partial x}(l, j) \right. \right. \\ \quad \left. \left. - V_{i,11}(l, j) \right) - \gamma_3 \left(W_{i,11}(l, j) - \frac{V_{i,22}(l, j)}{\det V_i(l, j)} \right) \left(\frac{V_{i,22}(l, j)}{(\det V_i(l, j))^2} \right) \right. \\ \quad \left. + \gamma_3 \left(W_{i,12}(l, j) + \frac{V_{i,12}(l, j)}{\det V_i(l, j)} \right) \left(\frac{V_{i,22}(l, j)V_{i,12}(l, j)}{(\det V_i(l, j))^2} \right) \right. \\ \quad \left. + \gamma_3 \left(W_{i,21}(l, j) + \frac{V_{i,21}(l, j)}{\det V_i(l, j)} \right) \left(\frac{V_{i,22}(l, j)V_{i,21}(l, j)}{(\det V_i(l, j))^2} \right) \right. \\ \quad \left. - \gamma_3 \left(W_{i,22}(l, j) - \frac{V_{i,11}(l, j)}{\det V_i(l, j)} \right) \left(-\frac{1}{\det V_i(l, j)} + \frac{V_{i,11}(l, j)V_{i,22}(l, j)}{(\det V_i(l, j))^2} \right) \right), \\ V_{i,11}(l, j) = \begin{cases} -\alpha & \text{if } \text{temp}_1(l, j) < -\alpha, \\ \text{temp}_1(l, j) & \text{if } |\text{temp}_1(l, j)| \leq \alpha, \\ \alpha & \text{if } \text{temp}_1(l, j) > \alpha \end{cases}, \\ \text{temp}_2(l, j) = V_{i,12}(l, j) + c \left(\frac{10a_3}{(\det V_i(l, j))^{11}} (-V_{i,21}(l, j)) - 4a_1(V_{i,12}(l, j)) \right. \\ \quad \left. \|V_i(l, j)\|^2 + 2a_2(\det V_i(l, j) - 1)(V_{i,21}(l, j)) + \gamma_2 \left(\frac{\partial \varphi_{i,1}}{\partial y}(l, j) \right. \right. \\ \quad \left. \left. - V_{i,12}(l, j) \right) + \gamma_3 \left(W_{i,11}(l, j) - \frac{V_{i,22}(l, j)}{\det V_i(l, j)} \right) \left(\frac{V_{i,22}(l, j)V_{i,21}(l, j)}{(\det V_i(l, j))^2} \right) \right. \\ \quad \left. - \gamma_3 \left(W_{i,12}(l, j) + \frac{V_{i,12}(l, j)}{\det V_i(l, j)} \right) \left(\frac{1}{\det V_i(l, j)} + \frac{V_{i,21}(l, j)V_{i,12}(l, j)}{(\det V_i(l, j))^2} \right) \right. \\ \quad \left. - \gamma_3 \left(W_{i,21}(l, j) + \frac{V_{i,21}(l, j)}{\det V_i(l, j)} \right) \left(\frac{V_{i,21}(l, j)^2}{(\det V_i(l, j))^2} \right) \right. \\ \quad \left. + \gamma_3 \left(W_{i,22}(l, j) - \frac{V_{i,11}(l, j)}{\det V_i(l, j)} \right) \left(\frac{V_{i,11}(l, j)V_{i,21}(l, j)}{(\det V_i(l, j))^2} \right) \right), \\ V_{i,12}(l, j) = \begin{cases} -\alpha & \text{if } \text{temp}_2(l, j) < -\alpha, \\ \text{temp}_2(l, j) & \text{if } |\text{temp}_2(l, j)| \leq \alpha, \\ \alpha & \text{if } \text{temp}_2(l, j) > \alpha \end{cases} \end{array} \right.$$

$$\left\{
 \begin{array}{l}
 \text{temp}_3(l, j) = V_{i,21}(l, j) + c \left(\frac{10a_3}{(\det V_i(l, j))^{\frac{1}{11}}} (-V_{i,12}(l, j)) - 4a_1(V_{i,21}(l, j)) \right. \\
 \quad \|V_i(l, j)\|^2 - 2a_2(\det V_i(l, j) - 1)(-V_{i,12}(l, j)) + \gamma_2 \left(\frac{\partial \varphi_{i,2}}{\partial x}(l, j) \right. \\
 \quad \left. - V_{i,21}(l, j) \right) + \gamma_3 \left(W_{i,11}(l, j) - \frac{V_{i,22}(l, j)}{\det V_i(l, j)} \right) \\
 \quad \left(\frac{V_{i,22}(l, j)V_{i,12}(l, j)}{(\det V_i(l, j))^2} \right) - \gamma_3 \left(W_{i,12}(l, j) + \frac{V_{i,12}(l, j)}{\det V_i(l, j)} \left(\frac{V_{i,12}(l, j)^2}{(\det V_i(l, j))^2} \right) \right. \\
 \quad \left. - \gamma_3 \left(W_{i,21}(l, j) + \frac{V_{i,21}(l, j)}{\det V_i(l, j)} \right) \left(\frac{1}{\det V_i(l, j)} + \frac{V_{i,12}(l, j)V_{i,21}(l, j)}{(\det V_i(l, j))^2} \right) \right. \\
 \quad \left. + \gamma_3 \left(W_{i,22}(l, j) - \frac{V_{i,11}(l, j)}{\det V_i(l, j)} \right) \left(\frac{V_{i,11}(l, j)(V_{i,12}(l, j))}{(\det V_i(l, j))^2} \right) \right), \\
 V_{i,21}(l, j) = \begin{cases} -\alpha & \text{if } \text{temp}_3(l, j) < -\alpha, \\ \text{temp}_3(l, j) & \text{if } |\text{temp}_3(l, j)| \leq \alpha, \\ \alpha & \text{if } \text{temp}_3(l, j) > \alpha \end{cases}, \\
 \text{temp}_4(l, j) = V_{i,22}(l, j) + c \left(\frac{10a_3}{(\det V_i(l, j))^{\frac{1}{11}}} V_{i,11}(l, j) - 4a_1(V_{i,22}(l, j)) \right. \\
 \quad \|V_i(l, j)\|^2 - 2a_2(\det V_i(l, j) - 1)V_{i,11}(l, j) \\
 \quad + \gamma_2 \left(\frac{\partial \varphi_{i,2}}{\partial y}(l, j) - V_{i,22}(l, j) \right) - \gamma_3 \left(W_{i,11}(l, j) - \frac{V_{i,22}(l, j)}{\det V_i(l, j)} \right) \\
 \quad \left(-\frac{1}{\det V_i(l, j)} + \frac{V_{i,22}(l, j)V_{i,11}(l, j)}{(\det V_i(l, j))^2} \right) + \gamma_3 \left(W_{i,12}(l, j) \right. \\
 \quad \left. + \frac{V_{i,12}(l, j)}{\det V_i(l, j)} \left(\frac{V_{i,11}(l, j)V_{i,12}(l, j)}{(\det V_i(l, j))^2} \right) \right) + \gamma_3 \left(W_{i,21}(l, j) + \frac{V_{i,21}(l, j)}{\det V_i(l, j)} \right) \\
 \quad \left(\frac{V_{i,11}(l, j)V_{i,21}(l, j)}{(\det V_i(l, j))^2} \right) - \gamma_3 \left(W_{i,22}(l, j) - \frac{V_{i,11}(l, j)}{\det V_i(l, j)} \right) \\
 \quad \left(\frac{(V_{i,11}(l, j))^2}{(\det V_i(l, j))^2} \right), \\
 V_{i,22}(l, j) = \begin{cases} -\alpha & \text{if } \text{temp}_4(l, j) < -\alpha, \\ \text{temp}_4(l, j) & \text{if } |\text{temp}_4(l, j)| \leq \alpha, \\ \alpha & \text{if } \text{temp}_4(l, j) > \alpha \end{cases}. \end{array} \right.$$

end for 2.5.

2.6. For each $i = 1, \dots, M$, for each pixel (l, j) , update W_i with this closed form:

$$\left\{
 \begin{array}{l}
 W_{i,11}(l, j) = \begin{cases} -\beta & \text{if } \frac{V_{i,22}(l, j)}{\det V_i(l, j)} < -\beta \\ \frac{V_{i,22}(l, j)}{\det V_i(l, j)} & \text{if } \left| \frac{V_{i,22}(l, j)}{\det V_i(l, j)} \right| \leq \beta \\ \beta & \text{if } \frac{V_{i,22}(l, j)}{\det V_i(l, j)} > \beta \end{cases} \\
 W_{i,12}(l, j) = \begin{cases} -\beta & \text{if } \frac{-V_{i,12}(l, j)}{\det V_i(l, j)} < -\beta \\ \frac{-V_{i,12}(l, j)}{\det V_i(l, j)} & \text{if } \left| \frac{-V_{i,12}(l, j)}{\det V_i(l, j)} \right| \leq \beta \\ \beta & \text{if } \frac{-V_{i,12}(l, j)}{\det V_i(l, j)} > \beta \end{cases} \\
 W_{i,21}(l, j) = \begin{cases} -\beta & \text{if } \frac{-V_{i,21}(l, j)}{\det V_i(l, j)} < -\beta \\ \frac{-V_{i,21}(l, j)}{\det V_i(l, j)} & \text{if } \left| \frac{-V_{i,21}(l, j)}{\det V_i(l, j)} \right| \leq \beta \\ \beta & \text{if } \frac{-V_{i,21}(l, j)}{\det V_i(l, j)} > \beta \end{cases} \\
 W_{i,22}(l, j) = \begin{cases} -\beta & \text{if } \frac{V_{i,11}(l, j)}{\det V_i(l, j)} < -\beta \\ \frac{V_{i,11}(l, j)}{\det V_i(l, j)} & \text{if } \left| \frac{V_{i,11}(l, j)}{\det V_i(l, j)} \right| \leq \beta \\ \beta & \text{if } \frac{V_{i,11}(l, j)}{\det V_i(l, j)} > \beta \end{cases} \end{array} \right. \text{end for 2.6.}$$

2.7. Solve for all $i = 1, \dots, M$, the Euler-Lagrange equation in U_i using an implicit finite difference scheme: $0 = \gamma_1 \nabla \theta_{T_i} \circ \varphi_i (\theta_{T_i} \circ \varphi_i - \theta_R - \theta_{\tilde{T}_i}) + \lambda_R (T_i \circ \varphi_i - \theta_R) \nabla T_i \circ \varphi_i + \gamma_2 \left(\frac{\text{div} V_{i,1}}{\text{div} V_{i,2}} \right)$, where $V_{i,j}$ stands for the j^{th} row of V_i , and $\varphi_i = Id + U_i$. end for 2.7.

2.8. $k := k + 1$.

end while

390 **SM3. Overall algorithm.**

391 **SM4. Detailed presentation of the first two methods to perform PCA.**

392 **SM4.1. First Approach : Linearisation around the Identity.** Let us recall the definition
393 of the Cauchy-stress tensors and the Cauchy axiom relating, in a hyperelastic framework, the
394 deformations and the forces applied on the material.

Definition SM4.1 (Cauchy-stress tensor). *In a hyperelastic framework, the Cauchy stress tensor in the reference configuration, is defined by*

$$\sigma^{ref}(x) = \frac{\partial W}{\partial F}(x, F),$$

395 where W is the stored energy function of the material. In a deformed configuration (characterised by the transformation φ), the first Piola-Kirchhoff tensor is defined by $\sigma(x) = \sigma^{ref}(\varphi^{-1}(x))\text{Cof}\nabla\varphi^{-1}(x)$.

398 **Axiom SM4.2 (Cauchy axiom).** *In an equilibrium position the following relations hold*

$$399 \quad \forall y \in \Gamma_1 \subset \partial\Omega, t(y, \vec{n}) = \sigma^{ref}(y)\vec{n},$$

$$400 \quad \forall y \in \Omega, f(y) = -\text{div } \sigma^{ref}(y),$$

402 where $t(y, \vec{n})$ is the pressure applied to the material at the boundary point y in the normal
403 direction \vec{n} , $f(y)$ is the inner volumetric force applied at $y \in \Omega$.

404 **Remark SM4.3.** In our model, we aim at having $\theta_{T_i} \circ \varphi_i$ close to θ_R , and therefore φ_i is
405 the deformation from θ_R to θ_{T_i} . The deformation from θ_{T_i} to θ_R is φ_i^{-1} and thanks to Ball's
406 results [?], we have that

$$\begin{aligned} 407 \quad \int_{\Omega} W_{Op}(\nabla\varphi_i) dx &= \int_{\Omega} \tilde{W}(\nabla\varphi_i, \text{Cof}\nabla\varphi_i, \det\nabla\varphi_i) dx, \\ 408 &= \int_{\Omega} \tilde{W}\left(\frac{\text{Cof}\nabla(\varphi_i^{-1})(\varphi_i)^T}{\det\nabla(\varphi_i^{-1})(\varphi_i)}, \frac{\nabla(\varphi_i^{-1})(\varphi_i)^T}{\det\nabla(\varphi_i^{-1})(\varphi_i)}, \frac{1}{\det\nabla(\varphi_i^{-1})(\varphi_i)}\right) dx, \\ 409 &= \int_{\Omega} \det\nabla(\varphi_i^{-1})\tilde{W}\left(\frac{\text{Cof}\nabla(\varphi_i^{-1})^T}{\det\nabla(\varphi_i^{-1})}, \frac{\nabla(\varphi_i^{-1})^T}{\det\nabla(\varphi_i^{-1})}, \frac{1}{\det\nabla(\varphi_i^{-1})}\right) dx, \\ 410 &= \int_{\Omega} \tilde{W}_{Op}(\nabla(\varphi_i^{-1})) dx, \end{aligned}$$

412 with $\tilde{W}(\psi, \phi, \delta) = \begin{cases} a_1\|\psi\|^4 + a_2\|\phi\|^4 + a_3(\delta - 1)^2 + \frac{a_4}{\delta^{10}} - 3a_1 - 3a_2 - a_4 & \text{if } \delta > 0, \\ +\infty & \text{otherwise} \end{cases}$. Therefore, the Cauchy-stress tensor applied to the reference configuration θ_{T_i} associated to the
413 deformation φ_i^{-1} to transform θ_{T_i} into θ_R is $\sigma_i^{ref}(x) = \frac{\partial \tilde{W}_{Op}}{\partial F}(\nabla\varphi_i^{-1}(x))$. The corresponding
414 first Piola-Kirchhoff tensor applied to the deformed configuration θ_R is given by $\sigma_i = \frac{\partial \tilde{W}_{Op}}{\partial F}(\nabla\varphi_i^{-1}(\varphi_i))\text{Cof}\nabla\varphi_i$. The inner volumetric forces applied to θ_R that are necessary to
415 deform θ_{T_i} into θ_R are given by $-\text{div}\sigma_i$.

418 Equipped with this material and the previous observation, we consider our mean configuration
 419 θ_R as a free material (not pre-stressed) and look for the displacements v_i induced by a small
 420 increase of the impact of θ_{T_i} parameterised by δ and minimising the following energy for each
 421 $i = 1, \dots, M$:

$$\begin{aligned} 422 \quad \inf_{v_i \in \mathcal{V}} \left\{ \mathcal{F}_i^\delta(v_i) &= \int_{\Omega} \tilde{W}_{Op}((\mathbf{I} + \delta \nabla v_i)) dx - \delta^2 \int_{\Omega} \operatorname{div} \sigma_i : v_i dx, \\ 423 \quad &= \int_{\Omega} \tilde{W}_{Op}((\mathbf{I} + \delta \nabla v_i)) dx + \delta^2 \int_{\Omega} \sigma_i : \nabla v_i dx - \delta^2 \int_{\partial\Omega} \sigma_i v_i \vec{n}_{\partial\Omega} ds, \\ 424 \quad &= \int_{\Omega} \tilde{W}_{Op}((\mathbf{I} + \delta \nabla v_i)) dx + \delta^2 \int_{\Omega} \sigma_i : \nabla v_i dx \right\}, \\ 425 \end{aligned}$$

426 with $\mathcal{V} := \{\psi \in W_0^{1,4}(\Omega, \mathbb{R}^3) \mid \operatorname{Cof}(\mathbf{I} + \delta \nabla \psi) \in L^4(\Omega, M_3(\mathbb{R})), \det(\mathbf{I} + \delta \nabla \psi) \in L^2(\Omega), \frac{1}{\det(\mathbf{I} + \delta \nabla \psi)} \\ 427 \in L^{10}(\Omega), \det(I + \delta \psi) > 0 \text{ a.e. on } \Omega\}$. The first term corresponds to the stored energy of the
 428 chosen Ogden material, while the second component represents the work of the applied forces
 429 with the following boundary conditions: $v_i = 0$ on $\partial\Omega$. However, the solution of this problem
 430 still lives in a nonlinear space and we thus propose to linearise the stored energy function \tilde{W}_{Op}
 431 around the identity. Before linearising the functional using a Taylor development, we introduce
 432 the deformation Cauchy-Green tensor $b_i = (\mathbf{I} + \delta v_i)^T (\mathbf{I} + \delta v_i) = \mathbf{I} + \delta(\nabla v_i^T + \nabla v_i) + \delta^2 \nabla v_i^T \nabla v_i = \\ 433 \mathbf{I} + 2\delta\epsilon(v_i) + \delta^2 \nabla v_i^T \nabla v_i \approx \mathbf{I} + 2\delta\epsilon(v_i)$. We first linearise the deformation tensor then the stored
 434 energy function and will see that we come back to the linearised elasticity problem.

$$\begin{aligned} 435 \quad \mathcal{F}_i^\delta(v_i) &= \int_{\Omega} \tilde{W}_{Op}(\mathbf{I} + \delta \nabla v_i) dx + \delta^2 \int_{\Omega} \nabla v_i : \sigma_i dx, \\ 436 \quad &= \int_{\Omega} \bar{W}_{Op}(b_i) + \delta^2 \int_{\Omega} \nabla v_i : \sigma_i dx, \\ 437 \quad &= \int_{\Omega} \bar{W}_{Op}(\mathbf{I} + 2\delta\epsilon(v_i)) + \delta^2 \int_{\Omega} \nabla v_i : \sigma_i dx, \\ 438 \quad &= \int_{\Omega} \bar{W}_{Op}(\mathbf{I}) dx + 2\delta \int_{\Omega} \frac{\partial \bar{W}_{Op}}{\partial b}(\mathbf{I}) : \epsilon(v_i) dx + 2\delta^2 \int_{\Omega} \frac{\partial^2 \bar{W}_{Op}}{\partial b^2}(\mathbf{I}) : \epsilon(v_i) : \epsilon(v_i) dx \\ 439 \quad &\quad + \delta^2 \int_{\Omega} \nabla v_i : \sigma_i dx, \\ 440 \end{aligned}$$

441 with $\bar{W}_{Op}(b) = a_1 \frac{II_b^2}{\sqrt{III_b^3}} + a_2 \frac{I_b^2}{\sqrt{III_b^3}} + a_4 \sqrt{III_b}^{-11} + \frac{a_3}{\sqrt{III_b}} - 2a_3 + a_3 \sqrt{III_b} - (9a_1 + 9a_2 + a_4) \sqrt{III_b}$,
 442 $I_b = \operatorname{Tr}(b) = \|\mathbf{I} + \delta v_i\|_F^2, II_b = \frac{\operatorname{Tr}(b)^2 - \operatorname{Tr}(b^2)}{2} = \|\operatorname{Cof}(\mathbf{I} + \delta v_i)\|_F^2$ and $III_b = \det b = \det(\mathbf{I} + \delta v_i)^2$.
 443 The detailed computations of the derivatives are given in section SM5. For our configuration
 444 to be in equilibrium, we need $\frac{\partial \bar{W}_{Op}}{\partial b}|_{(I_b, II_b, III_b)=(3,3,1)} = 0$, implying $a_4 = \frac{12a_2 + 6a_1}{5}$. We also
 445 set $\mu = 12a_1 + 12a_2 > 0$ and $\lambda = 4(32a_1 + 74a_2 + \frac{a_3}{2}) > 0$. Since $\mathbf{I} \otimes \mathbf{I} : \epsilon(v) : \epsilon(v) = \operatorname{Tr}(\epsilon^2)$,
 446 $\mathbf{I} : \epsilon(v) : \epsilon(v) = \bar{\mathbf{I}} : \epsilon(v) : \epsilon(v) = \operatorname{Tr}(\epsilon^2)$ by symmetry of $\epsilon(v)$, we finally obtain a linearised

447 elasticity functional with $v_i \in H_0^1(\Omega, \mathbb{R}^3)$:

$$\begin{aligned} 448 \quad \mathcal{F}_i^\delta(v_i) &= \int_{\Omega} \tilde{W}_{Op}(\mathbf{I} + \delta v_i) dx + \delta^2 \int_{\Omega} \nabla v_i : \sigma_i dx, \\ 449 \quad &= \int_{\Omega} \bar{W}_{Op}(\mathbf{I}) dx + 2\delta \int_{\Omega} \frac{\partial \bar{W}_{Op}}{\partial b}(\mathbf{I}) : \epsilon(v_i) dx + 2\delta^2 \int_{\Omega} \frac{\partial^2 \bar{W}_{Op}}{\partial b^2}(\mathbf{I}) : \epsilon(v_i) : \epsilon(v_i) dx \\ 450 \quad &\quad + \delta^2 \int_{\Omega} \nabla v_i : \sigma_i dx, \\ 451 \quad &= 2\delta^2 \int_{\Omega} \mu \text{Tr}(\epsilon(v_i)^2) + \frac{\lambda}{2} \text{Tr}(\epsilon(v_i))^2 dx + \delta^2 \int_{\Omega} \sigma_i : \nabla v_i dx. \\ 452 \end{aligned}$$

453 We now consider the following linearised elasticity problem (LEP)

$$454 \quad (\text{LEP}) \quad \inf_{v_i \in H_0^1(\Omega, \mathbb{R}^3)} \{ \mathcal{F}_i^\delta(v_i) = 2\delta^2 \int_{\Omega} \mu \text{Tr}(\epsilon(v_i)^2) + \frac{\lambda}{2} \text{Tr}(\epsilon(v_i))^2 dx + \delta^2 \int_{\Omega} \sigma_i : \nabla v_i dx \}. \\ 455$$

456 By using the Lax-Milgram theorem, one can prove the following theorem.

457 **Theorem SM4.4 (Existence and uniqueness of a solution to problem (LEP)).** *There exists a
458 unique solution of (LEP) in the linear space $H_0^1(\Omega, \mathbb{R}^2)$.*

459 *Proof.* In order to prove the existence and uniqueness of a minimizer in $H_0^1(\Omega)$, we recall
460 the Lax-Milgram theorem :

461 **Theorem SM4.5 (Lax-Milgram).** *Let \mathcal{H} be a Hilbert space. Let $a(., .)$ be a continuous bilinear
462 form on $\mathcal{H} \times \mathcal{H}$ and coercive on \mathcal{H} . Let $L(.)$ be a linear continuous form on \mathcal{H} . Then there
463 exists a unique element of \mathcal{H} , u , such that for all $v \in \mathcal{H}$, $a(u, v) = L(v)$.
464 Furthermore, if $a(., .)$ is symmetric, then u minimizes $J(v) = \frac{1}{2}a(v, v) - L(v)$, that is to say
465 there exists a unique $u \in \mathcal{H}$ such that $J(u) = \min_{v \in \mathcal{H}} J(v)$.*

466 We now set $\mathcal{H} = H_0^1(\Omega, \mathbb{R}^3)$, $a(u, v) = \int_{\Omega} 4\delta^2(\mu \epsilon(u) : \epsilon(v) + \frac{\lambda}{2} \text{Tr}(\epsilon(u)) \text{Tr}(\epsilon(v))) dx$ for all
467 $(u, v) \in H_0^1(\Omega, \mathbb{R}^3) \times H_0^1(\Omega, \mathbb{R}^3)$, and $L(v) = -\delta^2 \int_{\Omega} \sigma_i : \nabla v_i dx$ for all $v \in H_0^1(\Omega, \mathbb{R}^3)$. L is
468 linear on $H_0^1(\Omega, \mathbb{R}^3)$ by linearity of the gradient operator and the scalar product in $M_3(\mathbb{R})$.
469 It is also continuous on $H_0^1(\Omega, M_3(\mathbb{R}))$ by using Cauchy-Schwartz inequality and Poincaré's
470 inequality, $\forall v \in H_0^1(\Omega, \mathbb{R}^3)$:

$$471 \quad L(v) = \int_{\Omega} \sigma_i : \nabla v dx \leq \|\sigma_i\|_{L^2(\Omega, M_3(\mathbb{R}))} \|\nabla v\|_{L^2(\Omega, M_3(\mathbb{R}))} \leq \|\sigma_i\|_{L^2(\Omega, M_3(\mathbb{R}))} \|v\|_{H^1(\Omega, \mathbb{R}^3)}, \\ 472$$

473 with $\|\sigma_i\|_{L^2(\Omega, M_3(\mathbb{R}))} < +\infty$ since we have proved earlier that $\sigma_i \in L^2(\Omega, M_3(\mathbb{R}))$.
474 $a(., .)$ is bilinear on $H_0^1(\Omega, \mathbb{R}^3) \times H_0^1(\Omega, \mathbb{R}^3)$ by linearity of ϵ coming from the linearity of the
475 gradient operator and the linearity of the scalar product in $M_3(\mathbb{R})$ and the trace operator.
476 $a(., .)$ is also symmetric by symmetry of $\epsilon(u)$ and $\epsilon(v)$. $a(., .)$ is also continuous on $H_0^1(\Omega, \mathbb{R}^3) \times$

477 $H_0^1(\Omega, \mathbb{R}^3)$ since we have :

$$\begin{aligned} 478 \quad & \forall (u, v) \in H_0^1(\Omega, \mathbb{R}^3) \times H_0^1(\Omega, \mathbb{R}^3), a(u, v) = \int_{\Omega} 4\delta^2(\mu\epsilon(u) : \epsilon(v) + \frac{\lambda}{2}Tr(\epsilon(u))Tr(\epsilon(v))) dx, \\ 479 \quad & \leq 4\delta^2\mu\|\epsilon(u)\|_{L^2(\Omega, M_3(\mathbb{R}))}\|\epsilon(v)\|_{L^2(\Omega, M_3(\mathbb{R}))} + 2\delta^2\lambda\|Tr(\epsilon(u))\|_{L^2(\Omega)}\|Tr(\epsilon(v))\|_{L^2(\Omega)}, \\ 480 \quad & \leq 4\delta^2\mu\|\nabla u\|_{L^2(\Omega, M_3(\mathbb{R}))}\|\nabla v\|_{L^2(\Omega, M_3(\mathbb{R}))} + 2\delta^2\lambda\|\nabla u\|_{L^2(\Omega, M_3(\mathbb{R}))}\|\nabla v\|_{L^2(\Omega, M_3(\mathbb{R}))}, \\ 481 \quad & \leq (4\delta^2\mu + 2\delta^2\lambda)\|u\|_{H^1(\Omega)}\|v\|_{H^1(\Omega)}, \text{ Poincaré's inequality.} \end{aligned}$$

483 We now need to prove that $a(., .)$ is coercive. Let $v \in H_0^1(\Omega, \mathbb{R}^3)$, then we have

$$484 \quad a(v, v) = \int_{\Omega} 4\delta^2(\mu\epsilon(v) : \epsilon(v) + \frac{\lambda}{2}Tr(\epsilon(v))^2) dx \geq 4\delta^2\mu\|\epsilon(v)\|_{L^2(\Omega, M_3(\mathbb{R}))}^2. \\ 485$$

486 Since $4\delta^2\mu > 0$, we just need to prove the coercivity of $\|\epsilon(v)\|_{L^2(\Omega, M_3(\mathbb{R}))}$. To do so, we
487 use a reductio ad absurdum reasoning as in [?] and assume there exists a sequence $(v_n) \in$
488 $H_0^1(\Omega, M_3(\mathbb{R}))$ such that $\|\epsilon(v_n)\|_{L^2(\Omega, M_3(\mathbb{R}))} \xrightarrow{n \rightarrow +\infty} 0$ and $\|v_n\|_{H_0^1(\Omega, \mathbb{R}^3)} = 1$. As (v_n) is uniformly
489 bounded according to n in $H^1(\Omega, \mathbb{R}^3)$, we can extract a subsequence still denoted (v_n) such
490 that $v_n \xrightarrow{n \rightarrow +\infty} v^*$ in $H^1(\Omega, \mathbb{R}^3)$ and by continuity of the trace operator, we get that $v^* \in$
491 $H_0^1(\Omega)$. We also have that $\epsilon(v_n) \xrightarrow{n \rightarrow +\infty} \epsilon(v^*)$ in $L^2(\Omega, M_3(\mathbb{R}))$. However, we have assumed
492 that $\epsilon(v_n) \xrightarrow{n \rightarrow +\infty} 0$ in $L^2(\Omega, M_3(\mathbb{R}))$ and so by uniqueness of the weak limit, we get that
493 $\epsilon(v^*) = 0$. Then by [?, Lemma IV.3.3], it comes that v^* is an infinitesimal rigid displacement.
494 But a rigid displacement can only be null on a straight line if $a \neq 0$, nowhere if $a \neq 0, b \neq 0$
495 or everywhere if $a = b = 0$. Since v^* is null on $\partial\Omega$ with $\text{meas}(\partial\Omega) > 0$, then we are necessary
496 in the last case and so $v^* = 0$. Then Rellich theorem allows us to deduce that $v_n \rightarrow 0$ in
497 $L^2(\Omega, \mathbb{R}^3)$. Then by using Korn's inequality ([?, Theorem IV.3.2]), we get that $\nabla v_n \xrightarrow{n \rightarrow +\infty} 0$
498 in $L^2(\Omega, M_3(\mathbb{R}))$ so that $v_n \xrightarrow{n \rightarrow +\infty} 0$ in $H^1(\Omega, \mathbb{R}^3)$ which contradicts our second hypothesis.
499 Therefore, $a(., .)$ is coercive on $H_0^1(\Omega, \mathbb{R}^3)$. Lax Milgram's theorem applies and there exists a
500 unique solution to our linearised elasticity problem (LEP) in $H_0^1(\Omega)$. ■

501 This problem can be solved in the finite element approximation framework which results in
502 the resolution of a sparse high-dimensional linear system. A classical PCA is then performed
503 on the obtained displacement fields using the $L^2(\Omega, \mathbb{R}^3)$ scalar product for the covariance
504 operator. We now address the second method based on the Cauchy-stress tensors.

505 **SM4.2. Second Approach : PCA on the Cauchy-stress Tensors.** This approach relies
506 on the following lemma.

507 **Lemma SM4.6.** *The Cauchy-stress tensors σ_i^{ref} and σ_i are in $L^\infty(\Omega, M_3(\mathbb{R})) \subset L^2(\Omega, M_3(\mathbb{R}))$.* ■

508 **Proof.** Let us compute σ_i^{ref} and σ_i and make sure they belong to $L^\infty(\Omega, M_3(\mathbb{R}))$. We first
509 recall that $\sigma_i^{ref} = \frac{\partial \tilde{W}_{Op}}{\partial F}(\nabla(\varphi_i^{-1})) = \frac{\partial}{\partial F}(\det F(a_1 \frac{\|\text{Cof } F\|_F^4}{\det F^4} + a_2 \frac{\|F\|_F^4}{\det F^4} + a_4 \det F^{10} - a_3 (\frac{1}{\det F} - 1)^2 -$

510 $9a_1 - 9a_2 - a_4)(\nabla(\varphi_i^{-1}))$. We compute the following terms :

511 $F = \begin{pmatrix} F_{11} & F_{12} & F_{13} \\ F_{21} & F_{22} & F_{23} \\ F_{31} & F_{32} & F_{33} \end{pmatrix},$

512 $\text{Cof } F = \begin{pmatrix} F_{22}F_{33} - F_{32}F_{23} & F_{31}F_{23} - F_{21}F_{33} & F_{21}F_{32} - F_{31}F_{22} \\ F_{32}F_{13} - F_{12}F_{33} & F_{11}F_{33} - F_{13}F_{31} & F_{31}F_{12} - F_{11}F_{32} \\ F_{12}F_{23} - F_{22}F_{13} & F_{21}F_{13} - F_{11}F_{23} & F_{11}F_{22} - F_{21}F_{12} \end{pmatrix},$

513 $\det F = F_{11}(F_{22}F_{33} - F_{32}F_{23}) + F_{12}(F_{31}F_{23} - F_{21}F_{33}) + F_{13}(F_{21}F_{32} - F_{31}F_{22}),$

514 $= F_{21}(F_{32}F_{13} - F_{12}F_{33}) + F_{22}(F_{11}F_{33} - F_{13}F_{31}) + F_{23}(F_{31}F_{12} - F_{11}F_{32}),$

515 $= F_{31}(F_{12}F_{23} - F_{22}F_{13}) + F_{32}(F_{21}F_{13} - F_{11}F_{23}) + F_{33}(F_{11}F_{22} - F_{21}F_{12}),$

516 $\frac{\partial \|F\|_F^2}{\partial F} = 2F,$

517 $(\frac{\partial \|\text{Cof } F\|_F^2}{\partial F})_1^T = 2 \begin{pmatrix} F_{33}\text{Cof } F_{22} - F_{32}\text{Cof } F_{23} - F_{23}\text{Cof } F_{32} + F_{22}\text{Cof } F_{33} \\ -F_{33}\text{Cof } F_{21} + F_{31}\text{Cof } F_{23} + F_{23}\text{Cof } F_{31} - F_{21}\text{Cof } F_{33} \\ F_{32}\text{Cof } F_{21} - F_{31}\text{Cof } F_{22} - F_{22}\text{Cof } F_{31} + F_{21}\text{Cof } F_{32} \end{pmatrix},$

518 $(\frac{\partial \|\text{Cof } F\|_F^2}{\partial F})_2^T = 2 \begin{pmatrix} -F_{31}\text{Cof } F_{12} + F_{32}\text{Cof } F_{13} + F_{13}\text{Cof } F_{32} - F_{12}\text{Cof } F_{33} \\ F_{33}\text{Cof } F_{11} - F_{31}\text{Cof } F_{13} - F_{13}\text{Cof } F_{31} + F_{11}\text{Cof } F_{33} \\ -F_{32}\text{Cof } F_{11} + F_{31}\text{Cof } F_{12} + F_{12}\text{Cof } F_{31} - F_{11}\text{Cof } F_{32} \end{pmatrix},$

519 $(\frac{\partial \|\text{Cof } F\|_F^2}{\partial F})_3^T = 2 \begin{pmatrix} F_{23}\text{Cof } F_{12} - F_{22}\text{Cof } F_{13} - F_{13}\text{Cof } F_{22} + F_{12}\text{Cof } F_{23} \\ -F_{23}\text{Cof } F_{11} + F_{21}\text{Cof } F_{13} + F_{13}\text{Cof } F_{21} - F_{11}\text{Cof } F_{23} \\ F_{22}\text{Cof } F_{11} - F_{21}\text{Cof } F_{12} - F_{12}\text{Cof } F_{21} + F_{11}\text{Cof } F_{22} \end{pmatrix},$

520 $\frac{\partial \det F}{\partial F} = \text{Cof } F,$

521 $\sigma_i^{ref} = 2a_1 \frac{\partial \|\text{Cof } F\|_F^2}{\partial F} \frac{\|\text{Cof } F\|_F^2}{\det F^3} - 3a_1 \frac{\|\text{Cof } F\|_F^4}{\det F^4} \frac{\partial \det F}{\partial F} - 9a_1 \frac{\partial \det F}{\partial F}$
 522 $+ 2a_2 \frac{\partial \|F\|_F^2}{\partial F} \frac{\|F\|_F^2}{\det F^3} - 9a_2 \frac{\partial \det F}{\partial F} - 3a_2 \frac{\|F\|_F^4}{\det F^4} \frac{\partial \det F}{\partial F}$
 523 $- \frac{a_3}{\det F^2} \frac{\partial \det F}{\partial F} + a_3 \frac{\partial \det F}{\partial F} + 11a_4 \det F^{10} \frac{\partial \det F}{\partial F} - a_4 \frac{\partial \det F}{\partial F},$

526 $= 2a_1 \frac{\|\text{Cof } F\|_F^2}{\det F^3} \frac{\partial \|\text{Cof } F\|_F^2}{\partial F} - 3a_1 \frac{\|\text{Cof } F\|_F^4}{\det F^4} \text{Cof } F - 9a_1 \text{Cof } F$
 527 $+ 4a_2 \frac{\|F\|_F^2}{\det F^3} F - 9a_2 \text{Cof } F - 3a_2 \frac{\|F\|_F^4}{\det F^4} \text{Cof } F$
 528 $- \frac{a_3}{\det F^2} \text{Cof } F + a_3 \text{Cof } F + 11a_4 \det F^{10} \text{Cof } F - a_4 \text{Cof } F.$

530 Since, for all $i \in \{1, \dots, M\}$, $\|\nabla \varphi_i\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha$ and $\|(\nabla \varphi_i)^{-1}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta$, we
 531 deduce that

532 $\|\nabla(\varphi_i^{-1})\|_{L^\infty(\Omega, M_3(\mathbb{R}))} = \|(\nabla \varphi_i)^{-1}(\varphi_i^{-1})\|_{L^\infty(\Omega, M_3(\mathbb{R}))} = \|(\nabla \varphi_i)^{-1}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \beta$

534 and

$$\begin{aligned} 535 \quad \|\nabla \varphi_i\|_{L^\infty(\Omega, M_3(\mathbb{R}))} &= \|(\nabla \varphi_i^{-1})^{-1}(\varphi_i)\|_{L^\infty(\Omega, M_3(\mathbb{R}))} = \|(\nabla \varphi_i^{-1})^{-1}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \\ 536 \quad &= \left\| \frac{\text{Cof} \nabla(\varphi_i)^{-1}}{\det \nabla(\varphi_i)^{-1}} \right\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha \\ 537 \end{aligned}$$

538 so that $\|\text{Cof} \nabla(\varphi_i)^{-1}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq 18\beta^2$, $\|\det \nabla(\varphi_i)^{-1}\|_{L^\infty(\Omega)} \leq 6\beta^3$ and $\|\text{Cof} \nabla \varphi_i\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq 18\alpha^2$, $\|\det \nabla \varphi_i\|_{L^\infty(\Omega)} = \left\| \frac{1}{\det \nabla(\varphi_i^{-1})(\varphi_i)} \right\|_{L^\infty(\Omega)} = \left\| \frac{1}{\det \nabla(\varphi_i)^{-1}} \right\|_{L^\infty(\Omega)} \leq 6\alpha^3$. From these observations, we deduce that $\sigma_i^{ref} \in L^\infty(\Omega, M_3(\mathbb{R})) \subset L^2(\Omega, M_3(\mathbb{R}))$ and therefore $\sigma_i \in L^\infty(\Omega, M_3(\mathbb{R})) \subset L^2(\Omega, M_3(\mathbb{R}))$. ■

542 We thus propose to perform a Principal Component Analysis on the tensors σ_i directly since
543 they belong to $L^2(\Omega, M_3(\mathbb{R}))$ and following [?]. To do so, let $V = L^2(\Omega, M_3(\mathbb{R}))$ and let \mathcal{V} be
544 a random variable with values in V . The associated covariance operator is thus defined by

$$545 \quad \mathcal{C}_V(\sigma) = \mathbb{E}[\langle \sigma, \mathcal{V} - \mathbb{E}\mathcal{V} \rangle_V (\mathcal{V} - \mathbb{E}\mathcal{V})], \forall \sigma \in V.$$

547 Then, the corresponding empirical estimation of this operator is given by

$$548 \quad \hat{\mathcal{C}}(\sigma) = \frac{1}{M} \sum_{i=1}^M \langle \sigma, \sigma_i - \bar{\sigma} \rangle_V (\sigma_i - \bar{v}), \sigma \in V, \bar{\sigma} = \frac{1}{M} \sum_{i=1}^M \sigma_i. \\ 549$$

550 We are now looking for the eigenvalues (λ_j) and eigenvectors (ψ_j) associated to this operator,
551 and then sort them in a decreasing order, i.e. $\lambda_1 \geq \lambda_2 \geq \dots$, so that (ψ_j) is the j -th principal
552 component function. Since the number of observations is small with respect to the size of the
553 space, an appropriate choice for the base of V is given by the collection of observations σ_i .
554 We then aim to solve

$$555 \quad \frac{1}{M} \sum_{i=1}^M (\sigma_i - \bar{V}) \sum_{j=1}^M b_{ji} \langle (\sigma_i - \bar{\sigma}), (\sigma_j - \bar{\sigma}) \rangle_V = \lambda_j \sum_{i=1}^M b_{ji} (\sigma_i - \bar{\sigma}), \\ 556 \quad \iff \frac{1}{M} W b_j = \lambda_j b_j, \\ 557$$

558 where $W = (W_{ij})_{i,j} = (\langle \sigma_i - \bar{\sigma}, \sigma_j - \bar{\sigma} \rangle_V)_{i,j} \in M_M(\mathbb{R})$. It results in finding the eigenvalues
559 and eigenvectors of the matrix W . Then the j -th PC function is given by $\psi_j = \sum_{i=1}^M b_{ji} (\sigma_i - \bar{\sigma})$.
560 In order to come back to the deformation space and to properly model the loaded configuration,
561 we compose the sought deformation with the previous deformations. We then solve the
562 following minimization problem with the last term ensuring that the sought displacements v_j

563 are Lipschitz continuous so that the chain rule applies:

$$\begin{aligned}
 564 \quad & \inf \left\{ \mathcal{F}(v_j) = \frac{1}{M} \sum_{k=1}^M \int_{\Omega} \tilde{W}_{Op}(\nabla((\text{Id} - \delta v_j)^{-1} \circ \varphi_k^{-1})) dx - \delta^2 \int_{\Omega} \psi_j : \nabla v_j dx \right. \\
 565 \quad & + \mathbb{1}_{\{\|\cdot\|_{L^\infty(\Omega)} \leq \alpha\}} (\mathbf{I} - \delta \nabla v_j), \\
 566 \quad & = \frac{1}{M} \sum_{k=1}^M \int_{\Omega} \tilde{W}_{Op}(\nabla(\varphi_k^{-1}) - \delta \nabla v_i(\varphi_k^{-1}) \nabla \varphi_k^{-1}) dx \\
 567 \quad & - \delta^2 \int_{\Omega} \psi_j : \nabla v_j dx + \mathbb{1}_{\{\|\cdot\|_{L^\infty(\Omega)} \leq \alpha\}} (\mathbf{I} - \delta \nabla v_j), \\
 568 \quad & = \frac{1}{M} \sum_{k=1}^M \int_{\Omega} \det \nabla \varphi_k \tilde{W}_{Op}(\nabla(\varphi_k^{-1})(\varphi_k) - \delta \nabla v_i \nabla \varphi_k^{-1}(\varphi_k)) dy \\
 569 \quad & - \delta^2 \int_{\Omega} \psi_j : \nabla v_j dx + \mathbb{1}_{\{\|\cdot\|_{L^\infty(\Omega)} \leq \alpha\}} (\mathbf{I} - \delta \nabla v_j), \\
 570 \quad & = \frac{1}{M} \sum_{k=1}^M \int_{\Omega} \det \nabla \varphi_k \tilde{W}_{Op}((\nabla \varphi_k)^{-1} - \delta \nabla v_i(\nabla \varphi_k)^{-1}) dy \\
 571 \quad & \left. - \delta^2 \int_{\Omega} \psi_j : \nabla v_j dx + \mathbb{1}_{\{\|\cdot\|_{L^\infty(\Omega)} \leq \alpha\}} (\mathbf{I} - \delta \nabla v_j) \right\}. \\
 572
 \end{aligned}$$

573 \tilde{W}_{Op} is continuous since $\lim_{\det \xi \rightarrow 0^+} \tilde{W}_{Op}(\xi) = +\infty$. Let us set $\hat{\mathcal{W}}_1 = \{\psi \in W_0^{1,\infty}(\Omega, \mathbb{R}^3), \det(\mathbf{I} - \delta \nabla \psi) > 0$ a.e. on $\Omega, \|\mathbf{I} - \delta \nabla \psi\|_{L^\infty(\Omega)} \leq \alpha, \forall k \in \{1, \dots, M\}, \frac{\|\text{Cof}((\nabla \varphi_k)^{-1} - \delta \nabla \psi(\nabla \varphi_k)^{-1})\|_F^4}{\det((\nabla \varphi_k)^{-1} - \delta \nabla \psi(\nabla \varphi_k)^{-1})^3} \in L^1(\Omega), \frac{\|(I - \delta \nabla \psi)(\nabla \varphi_k)^{-1}\|_F^4}{\det((\nabla \varphi_k)^{-1} - \delta \nabla \psi(\nabla \varphi_k)^{-1})^3} \in L^1(\Omega), \det((\nabla \varphi_k)^{-1} - \delta \nabla \psi(\varphi_k)^{-1}) \in L^{11}(\Omega)\}$. The minimizer is searched for $v_j \in \hat{\mathcal{W}}$.

577 **Theorem SM4.7 (Existence of minimizers).** *One can prove that this problem (P1) admits
578 at least one solution.*

579 **Proof.** We follow the classical steps of the direct method of the calculus of variations.

580 **1. Coercivity inequality :**

581 Let us first derive a coercivity inequality :

$$\begin{aligned}
 582 \quad & \mathcal{F}(v) \geq \frac{1}{M} \sum_{k=1}^M \int_{\Omega} \frac{1}{6\beta^3} (a_1 \frac{\|\text{Cof}((\nabla \varphi_k)^{-1} - \delta \nabla v(\nabla \varphi_k)^{-1})\|_F^4}{\det((\nabla \varphi_k)^{-1} - \delta \nabla v(\nabla \varphi_k)^{-1})^3} \\
 583 \quad & + a_2 \frac{\|(\nabla \varphi_k)^{-1} - \delta \nabla v(\nabla \varphi_k)^{-1}\|_F^4}{\det((\nabla \varphi_k)^{-1} - \delta \nabla v(\nabla \varphi_k)^{-1})^3} + a_4 \det(\nabla \varphi_k^{-1} - \delta \nabla v_i(\nabla \varphi_k)^{-1})^{11} \\
 584 \quad & - (9a_1 + 9a_2 + a_4)36\alpha^6 + \mathbb{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha\}} (\mathbf{I} - \delta \nabla v),
 \end{aligned}$$

586 since Ω is bounded, $\det((\nabla \varphi_k)^{-1} - \delta \nabla v(\nabla \varphi_k)^{-1}) = \det((\mathbf{I} - \delta \nabla v)(\nabla \varphi_k)^{-1}) = \det(\mathbf{I} - \delta \nabla v) \det((\nabla \varphi_k)^{-1}) \leq 36\alpha^6$, and $\|\det \nabla \varphi_k\|_{L^\infty(\Omega)} \leq 6\alpha^3$, $\|\frac{1}{\det \nabla \varphi_k}\|_{L^\infty(\Omega)} \leq 6\beta^3$ so
587 that for all $x \in \Omega$, $\frac{1}{6\beta^3} \leq \det \nabla \varphi_k(x) \leq 6\alpha^3$. Furthermore, by taking $v = 0$, then
588

589 $\mathcal{F}(v) = \frac{1}{M} \int_{\Omega} \tilde{W}_{Op}(\nabla \varphi_k^{-1}) dx < +\infty$ from what precedes. Thus the functional is
590 proper and coercive so that the infimum exists and is finite.

591 **2. Convergence of a minimizing sequence :**

592 Let (v^n) be a minimizing sequence so that $\lim_{n \rightarrow +\infty} \mathcal{F}(v^n) = \inf \mathcal{F}(v) < +\infty$ from what
593 precedes. So there exists $M \in \mathbb{N}$ such that $\forall n \in \mathbb{N}, n \geq M \Rightarrow \mathcal{F}(v^n) \leq \inf \mathcal{F}(v) + 1 <$
594 $+\infty$. From now on, we will consider $n \geq M$.

595 From the previous coercivity inequality, one has:

- 596 • $((I - \delta \nabla v^n)(\nabla \varphi_k)^{-1})$ is uniformly bounded according to n in $L^4(\Omega, M_3(\mathbb{R}))$ for
597 all $k \in \{1, \dots, M\}$.
- 598 • $(I - \delta \nabla v^n)$ and so (∇v^n) is uniformly bounded according to n in $L^\infty(\Omega, M_3(\mathbb{R}))$.
- 599 • $(\text{Cof}((I - \delta v^n)(\nabla \varphi_k)^{-1})) = (\text{Cof}(I - \delta \nabla v^n)\text{Cof}((\nabla \varphi_k)^{-1}) = (\text{Cof}(I - \delta \nabla v^n)$
600 $(\text{Cof} \nabla \varphi_k)^{-1})$ is uniformly bounded according to n in $L^4(\Omega, M_3(\mathbb{R}))$ for all
601 $k \in \{1, \dots, M\}$.
- 602 • $(\det((I - \delta \nabla v^n)(\nabla \varphi_k)^{-1})) = (\frac{\det(I - \delta \nabla v^n)}{\det \nabla \varphi_k})$ is uniformly bounded according to
603 n in $L^{11}(\Omega)$.

604 There exists a subsequence still denoted $((I - \delta \nabla v^n)(\nabla \varphi_k)^{-1})$ such that

$$605 \quad (I - \delta \nabla v^n)(\nabla \varphi_k)^{-1} \xrightarrow[n \rightarrow +\infty]{} \alpha_k \text{ in } L^4(\Omega, M_3(\mathbb{R})), \forall k \in \{1, \dots, M\},$$

$$606 \quad I - \delta \nabla v^n \xrightarrow[n \rightarrow +\infty]{} \alpha_k \nabla \varphi_k \text{ in } L^4(\Omega, M_3(\mathbb{R})), \forall k \in \{1, \dots, M\},$$

$$607 \quad \nabla v^n \xrightarrow[n \rightarrow +\infty]{} \bar{V} = \frac{1}{\delta}(I - \alpha_k \nabla \varphi_k), \forall k \in \{1, \dots, M\}, \text{ in } L^4(\Omega, M_3(\mathbb{R})).$$

609 We thus deduce that (∇v^n) is uniformly bounded according to n in $L^4(\Omega, M_3(\mathbb{R}))$ and
610 by Poincaré's inequality, we deduce that (v^n) is uniformly bounded according to n in
611 $W^{1,4}(\Omega, \mathbb{R}^3)$. We can thus extract a subsequence still denoted (v^n) such that

$$612 \quad v^n \xrightarrow[n \rightarrow +\infty]{} \bar{v} \text{ in } W^{1,4}(\Omega, \mathbb{R}^3).$$

614 Furthermore, we can extract a subsequence still denoted (∇v^n) such that :

$$615 \quad \nabla v^n \xrightarrow[n \rightarrow +\infty]{*} \nabla \bar{v} \text{ in } L^\infty(\Omega, M_3(\mathbb{R})),$$

617 by uniqueness of the weak limit in $L^4(\Omega, M_3(\mathbb{R}))$ and the continuous embedding of
618 $L^\infty(\Omega, M_3(\mathbb{R}))$ into $L^4(\Omega, M_3(\mathbb{R}))$. Besides, using Poincaré-Wirtinger's inequality,
619 there exists a constant C depending only on Ω such that

$$620 \quad \|v^n\|_{L^\infty(\Omega, \mathbb{R}^3)} - \frac{\int_{\Omega} v^n dx}{\text{meas}(\Omega)} \leq \|v^n - \frac{\int_{\Omega} v^n dx}{\text{meas}(\Omega)}\|_{L^\infty(\Omega, \mathbb{R}^3)} \leq C \|\nabla v^n\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq C\alpha,$$

$$621 \quad \|v^n\|_{L^\infty(\Omega, \mathbb{R}^3)} \leq C\alpha + \frac{\int_{\Omega} v^n dx}{\text{meas}(\Omega)} \leq C\alpha + \frac{\|v^n\|_{L^1(\Omega, \mathbb{R}^3)}}{\text{meas}(\Omega)} \leq C\alpha + \frac{\|v^n\|_{L^4(\Omega, \mathbb{R}^3)}}{\text{meas}(\Omega)},$$

$$622 \quad \leq C\alpha + \frac{C'}{\text{meas}(\Omega)} \|\nabla v^n\|_{L^4(\Omega, M_3(\mathbb{R}))},$$

624 with C' a constant depending only on Ω coming from Poincaré's inequality. Since,
625 (∇v^n) is uniformly bounded in $L^4(\Omega, M_3(\mathbb{R}))$, we deduce that (v^n) is uniformly bounded
626 according to n in $W^{1,\infty}(\Omega, \mathbb{R}^3)$ and there exists a subsequence still denoted (v^n) such
627 that

$$628 \quad v^n \xrightarrow[n \rightarrow +\infty]{*} \bar{v} \text{ in } W^{1,\infty}(\Omega, \mathbb{R}^3),$$

630 by uniqueness of the weak limit in $W^{1,4}(\Omega, \mathbb{R}^3)$ and the continuous embedding of
631 $W^{1,\infty}(\Omega, \mathbb{R}^3)$ into $W^{1,4}(\Omega, \mathbb{R}^3)$. By continuity of the trace operator, we get $\bar{v} \in$
632 $W_0^{1,\infty}(\Omega, \mathbb{R}^3)$.

633 We can also extract subsequences of $(\text{Cof}(\mathbf{I} - \delta \nabla v^n)(\text{Cof} \nabla \varphi_k)^{-1})$ and $(\frac{\det(\mathbf{I} - \delta \nabla v^n)}{\det \nabla \varphi_k})$
634 still denoted $(\text{Cof}(\mathbf{I} - \delta \nabla v^n)(\text{Cof} \nabla \varphi_k)^{-1})$ and $(\frac{\det(\mathbf{I} - \delta \nabla v^n)}{\det \nabla \varphi_k})$ for all $k \in \{1, \dots, M\}$,
635 such that

$$636 \quad \text{Cof}(\mathbf{I} - \delta \nabla v^n)(\text{Cof} \nabla \varphi_k)^{-1} \xrightarrow[n \rightarrow +\infty]{*} H_k \text{ in } L^4(\Omega, M_3(\mathbb{R})),$$

$$637 \quad \text{Cof}(\mathbf{I} - \delta \nabla v^n) \xrightarrow[n \rightarrow +\infty]{*} H_k \text{Cof} \nabla \varphi_k \text{ in } L^4(\Omega, \mathbb{R}),$$

$$638 \quad \frac{\det(\mathbf{I} - \delta \nabla v^n)}{\det \nabla \varphi_k} \xrightarrow[n \rightarrow +\infty]{*} \delta_k \text{ in } L^{11}(\Omega),$$

$$639 \quad \det(\mathbf{I} - \delta \nabla v^n) \xrightarrow[n \rightarrow +\infty]{*} \delta_k \det \nabla \varphi_k \text{ in } L^{11}(\Omega).$$

641 By uniqueness of the weak limit, we have that $H = H_k \text{Cof} \nabla \varphi_k$ and $\delta = \delta_k \det \nabla \varphi_k$ for
642 all $k \in \{1, \dots, M\}$. By [?, Theorem VI.3.3] and $\text{Id} - \delta v^n \xrightarrow[n \rightarrow +\infty]{*} \text{Id} - \delta \bar{v}$ in $W^{1,4}(\Omega, \mathbb{R}^3)$,
643 we have that $H = \text{Cof}(\mathbf{I} - \delta \nabla \bar{v})$ and $\delta = \det(\mathbf{I} - \delta \nabla \bar{v})$.

644 **3. Lower semi-continuity :** \tilde{W}_{Op} is convex and continuous. Indeed, let $f(x, y) =$
645 $\frac{x^4}{y^3}$ for all $(x, y) \in (\mathbb{R}^+)^2$. Then the Hessian matrix of f is given by $H(x, y) =$
646 $\frac{12x^2}{y^3} \begin{pmatrix} 1 & -\frac{x}{y} \\ -\frac{x}{y} & \frac{x^2}{y^2} \end{pmatrix}$ and is positive semi-definite : $\forall (a, b) \in \mathbb{R}^2$, $(a, b)H(x, y)(a, b)^T =$
647 $\frac{12x^2}{y^3} (bx - ay)^2 \geq 0$, for all $(x, y) \in (\mathbb{R}^+)^2$. Besides, $\|\cdot\|_F$ is also convex and we have
648 $\forall (A, B) \in M_3(\mathbb{R})^2$, $\forall (c, d) \in (\mathbb{R}^+)^2$, $\forall \lambda \in (0, 1)$, $\frac{(\|\lambda A + (1-\lambda)B\|_F^4)}{(\lambda c + (1-\lambda)d)^3} \leq \frac{(\lambda \|A\|_F + (1-\lambda)\|B\|_F)^4}{(\lambda c + (1-\lambda)d)^3} =$
649 $f(\lambda \|A\|_F + (1-\lambda)\|B\|_F, \lambda c + (1-\lambda)d) \leq \lambda f(\|A\|_F, c) + (1-\lambda)f(\|B\|_F, d) = \lambda \frac{\|A\|_F^4}{c^3} +$
650 $(1-\lambda) \frac{\|B\|_F^4}{d^3}$. If $\psi_n \xrightarrow[n \rightarrow +\infty]{*} \bar{\psi}$ in $W^{1,4}(\Omega, \mathbb{R}^3)$ then $\nabla \psi_n \xrightarrow[n \rightarrow +\infty]{*} \nabla \bar{\psi}$ in $L^4(\Omega, M_3(\mathbb{R}))$ and
651 we can extract a subsequence still denoted $(\nabla \psi_n)$ such that $\nabla \psi_n \xrightarrow[n \rightarrow +\infty]{*} \nabla \bar{\psi}$ almost
652 everywhere in Ω . If $\alpha_n \xrightarrow[n \rightarrow +\infty]{*} \bar{\alpha}$ in $L^4(\Omega, M_3(\mathbb{R}))$ then we can extract a subsequence
653 still denoted (α_n) such that $\alpha_n \xrightarrow[n \rightarrow +\infty]{*} \bar{\alpha}$ almost everywhere in Ω . If $\delta_n \xrightarrow[n \rightarrow +\infty]{*} \bar{\delta}$ in
654 $L^{11}(\Omega)$, then there exists a subsequence still denoted (δ_n) such that $\delta_n \xrightarrow[n \rightarrow +\infty]{*} \bar{\delta}$ almost
655 everywhere in Ω . Then by continuity of \tilde{W}_{Op} , we get

$$656 \quad \tilde{W}_{Op}(\nabla \psi_n, \alpha_n, \delta_n) \xrightarrow[n \rightarrow +\infty]{*} \tilde{W}_{Op}(\nabla \bar{\psi}, \bar{\alpha}, \bar{\delta}) \text{ almost everywhere in } \Omega.$$

658

Then, by applying Fatou's lemma, we have that

$$659 \quad \liminf_{n \rightarrow +\infty} \int_{\Omega} \det \nabla \varphi_k \tilde{W}_{Op}(\nabla \psi_n, \alpha_n, \delta_n) dx \geq \int_{\Omega} \det \nabla \varphi_k \tilde{W}_{Op}(\nabla \bar{v}, \bar{\alpha}, \bar{\delta}) dx.$$

$$660$$

661 Since \tilde{W}_{Op} is convex, so is $\int_{\Omega} \det \nabla \varphi_k \tilde{W}_{Op}(\xi, \alpha, \delta) dx$, and we can apply [?, Corollaire
662 III.8] to get that $\int_{\Omega} \det \nabla \varphi_k \tilde{W}_{Op}(\xi, \alpha, \delta) dx$ is lower semicontinuous in $L^4(\Omega, M_3(\mathbb{R})) \times$
663 $L^4(\Omega, M_3(\mathbb{R})) \times L^{11}(\Omega)$. We deduce that

$$664 \quad +\infty > \liminf_{n \rightarrow +\infty} \int_{\Omega} \det \nabla \varphi_k \tilde{W}_{Op}((I - \delta \nabla v^n)(\nabla \varphi_k)^{-1}, \text{Cof}(I - \delta \nabla v^n)(\text{Cof} \nabla \varphi_k)^{-1}),$$

$$665 \quad \frac{\det(I - \delta \nabla v^n)}{\det \nabla \varphi_k} dx$$

$$666 \quad \geq \int_{\Omega} \det \nabla \varphi_k(x) \tilde{W}_{Op}((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1}, \text{Cof}(I - \delta \nabla \bar{v})(\text{Cof} \nabla \varphi_k)^{-1},$$

$$667 \quad \frac{\det(I - \delta \nabla \bar{v})}{\det \nabla \varphi_k} dx.$$

$$668$$

By the weak-* lower semi-continuity of the $\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))}$, we get that

$$\|I - \delta \nabla \bar{v}\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \liminf_{n \rightarrow +\infty} \|I - \nabla v^n\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha,$$

so that

$$\mathbb{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha\}}(I - \nabla \bar{v}) = 0 \leq \liminf_{n \rightarrow +\infty} \mathbb{1}_{\{\|\cdot\|_{L^\infty(\Omega, M_3(\mathbb{R}))} \leq \alpha\}}(I - \delta \nabla v^n).$$

Since $\psi_j \in L^2(\Omega, M_3(\mathbb{R}))$ and $\nabla v^n \xrightarrow{n \rightarrow +\infty} \nabla \bar{v}$ in $L^4(\Omega, M_3(\mathbb{R}))$ and so in $L^2(\Omega, M_3(\mathbb{R}))$,
we have that $-\delta^2 \int_{\Omega} \psi_j : \nabla \bar{v} dx = \lim_{n \rightarrow +\infty} -\delta^2 \int_{\Omega} \psi_j : \nabla v^n dx$.

By combining all the results, we get that

$$\liminf_{n \rightarrow +\infty} \mathcal{F}(v^n) \geq \mathcal{F}(\bar{v}).$$

669 Also $\bar{v} \in W_0^{1,\infty}(\Omega, \mathbb{R}^3)$, $\frac{\|\text{Cof}((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1})\|_F^4}{\det((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1})^3} \in L^1(\Omega)$, $\frac{\|(I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1}\|_F^4}{\det((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1})^3} \in L^1(\Omega)$
670 by finiteness of $\mathcal{F}(\bar{v})$ inducing the finiteness of $\int_{\Omega} \det \nabla \varphi_k \tilde{W}_{Op}((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1},$
671 $\text{Cof}((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1}), \det((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1})) dx$ for all $k \in \{1, \dots, M\}$. Since
672 $\tilde{W}_{Op}((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1}, \text{Cof}((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1}), \det((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1})) = +\infty$ when
673 $\det((I - \delta \nabla \bar{v})(\nabla \varphi_k)^{-1}) = \frac{\det(I - \delta \nabla \bar{v})(x)}{\det \nabla \varphi_k(x)} \leq 0 \iff \det(I - \delta \nabla \bar{v}) \leq 0$ since $\det \nabla \varphi_k(x) > 0$
674 on Ω , the set on which it happens must be of null measure otherwise we would have
675 $\mathcal{F}(\bar{v}) = +\infty$. So $\det(I - \delta \nabla \bar{v}) > 0$ almost everywhere in Ω and there exists a minimizer
676 to our initial problem $\bar{v} \in \hat{\mathcal{W}}_1$. \blacksquare

677 This problem is hard to solve in practice and we follow the same strategy as previously by
678 adding additional variables and using an alternative optimisation scheme.

SM5. Computation of the derivatives.

$$\begin{aligned}
 679 \quad & \frac{\partial \bar{W}_{Op}}{\partial I_b} = \frac{2a_2 I_b}{\sqrt{III_b}^3} = 6a_2|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 680 \quad & \frac{\partial \bar{W}_{Op}}{\partial II_b} = \frac{2a_1 I_b}{\sqrt{III_b}^3} = 6a_1|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 681 \quad & \frac{\partial \bar{W}_{Op}}{\partial III_b} = \frac{-3a_1 III_b^2}{2\sqrt{III_b}^5} - \frac{3a_2 III_b^2}{2\sqrt{III_b}^5} + \frac{11}{2}a_4 \sqrt{III_b}^9 - \frac{a_3}{2III_b \sqrt{III_b}} + \frac{a_3}{2\sqrt{III_b}^3} \\
 682 \quad & - \frac{9a_1 + 9a_2 + a_4}{2\sqrt{III_b}} = -18a_1 - 18a_2 + 5a_4|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 683 \quad & \frac{\partial^2 \bar{W}_{Op}}{\partial I_b^2} = \frac{2a_2}{\sqrt{III_b}^3} = 2a_2|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 684 \quad & \frac{\partial^2 \bar{W}_{Op}}{\partial II_b \partial I_b} = 0, \\
 685 \quad & \frac{\partial^2 \bar{W}_{Op}}{\partial III_b \partial I_b} = -\frac{-3a_2 I_b}{III_b^2 \sqrt{III_b}} = -9a_2|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 686 \quad & \frac{\partial^2 \bar{W}_{Op}}{\partial I_b \partial II_b} = 0, \\
 687 \quad & \frac{\partial^2 \bar{W}_{Op}}{\partial II_b^2} = \frac{2a_1}{\sqrt{III_b}^3} = 2a_1|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 688 \quad & \frac{\partial^2 \bar{W}_{Op}}{\partial III_b \partial II_b} = -\frac{-3a_1 II_b}{III_b^2 \sqrt{III_b}} = -9a_1|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 689 \quad & \frac{\partial^2 \bar{W}_{Op}}{\partial I_b \partial III_b} = \frac{-3a_2 I_b}{III_b^2 \sqrt{III_b}} = -9a_2|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 690 \quad & \frac{\partial^2 \bar{W}_{Op}}{\partial II_b \partial III_b} = \frac{-3a_1 II_b}{III_b^2 \sqrt{III_b}} = -9a_1|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 691 \quad & \frac{\partial^2 \bar{W}_{Op}}{\partial III_b^2} = \frac{15a_1 III_b^2}{4\sqrt{III_b}^7} + \frac{15a_2 III_b^2}{4\sqrt{III_b}^7} + \frac{99a_4}{4} \sqrt{III_b}^7 \\
 692 \quad & + \frac{3a_3}{4\sqrt{III_b}^3} - \frac{a_3}{4III_b \sqrt{III_b}} + \frac{9a_1 + 9a_2 + a_4}{4\sqrt{III_b}^3}, \\
 693 \quad & = 36a_1 + 36a_2 + \frac{a_3}{2} + 25a_4|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 694 \quad & \frac{\partial I_b}{\partial b} = I, \\
 695 \quad & \frac{\partial II_b}{\partial b} = I_b I - b = 2I|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 696 \quad & \frac{\partial III_b}{\partial b} = III_b b^{-1} = I|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 697 \quad & \frac{\partial I}{\partial b} = 0,
 \end{aligned}$$

$$\begin{aligned}
 & \frac{\partial b}{\partial b} = \mathbf{I}, \text{ Identity tensor of order 4 such that } \mathbf{I} : A : A = \text{Tr}(A^T A), \\
 & \frac{\partial b^{-1}}{\partial b} = \frac{\mathbf{I} + \bar{\mathbf{I}}}{2}|_{(I_b, II_b, III_b)=(3,3,1)}, \text{ with } \bar{\mathbf{I}} : A : A = \text{Tr}(A^2), \\
 & \frac{\partial \bar{W}_{Op}}{\partial b} = \frac{\partial \bar{W}_{Op}}{\partial I_b} \frac{\partial I_b}{\partial b} + \frac{\partial \bar{W}_{Op}}{\partial II_b} \frac{\partial II_b}{\partial b} + \frac{\partial \bar{W}_{Op}}{\partial III_b} \frac{\partial III_b}{\partial b} = (-12a_1 - 6a_1 + 5a_4)I|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 & \frac{\partial^2 \bar{W}_{Op}}{\partial b^2} = \frac{\partial \bar{W}_{Op}}{\partial I_b} \frac{\partial I}{\partial b} + I \otimes \left(\frac{\partial^2 \bar{W}_{Op}}{\partial I_b^2} \frac{\partial I_b}{\partial b} + \frac{\partial^2 \bar{W}_{Op}}{\partial II_b \partial I_b} \frac{\partial II_b}{\partial b} + \frac{\partial^2 \bar{W}_{Op}}{\partial III_b \partial I_b} \frac{\partial III_b}{\partial b} \right) \\
 & \quad + \frac{\partial \bar{W}_{Op}}{\partial II_b} \frac{\partial (I_b I - b)}{\partial b} + (I_b I - b) \otimes \left(\frac{\partial^2 \bar{W}_{Op}}{\partial I_b \partial II_b} \frac{\partial I_b}{\partial b} + \frac{\partial^2 \bar{W}_{Op}}{\partial II_b^2} \frac{\partial II_b}{\partial b} + \frac{\partial^2 \bar{W}_{Op}}{\partial III_b \partial II_b} \frac{\partial III_b}{\partial b} \right) \\
 & \quad + \frac{\partial \bar{W}_{Op}}{\partial III_b} \frac{\partial (III_b b^{-1})}{\partial b} + (III_b b^{-1}) \otimes \left(\frac{\partial^2 \bar{W}_{Op}}{\partial I_b \partial III_b} \frac{\partial I_b}{\partial b} + \frac{\partial^2 \bar{W}_{Op}}{\partial II_b \partial III_b} \frac{\partial II_b}{\partial b} + \frac{\partial^2 \bar{W}_{Op}}{\partial III_b^2} \frac{\partial III_b}{\partial b} \right), \\
 & = I \otimes \left(\frac{\partial^2 \bar{W}_{Op}}{\partial I_b^2} I + \frac{\partial^2 \bar{W}_{Op}}{\partial II_b \partial I_b} (I_b I - b) + \frac{\partial^2 \bar{W}_{Op}}{\partial III_b \partial I_b} III_b b^{-1} \right) \\
 & \quad + \frac{\partial \bar{W}_{Op}}{\partial II_b} (I \otimes I - \mathbf{I}) + (I_b I - b) \otimes \left(\frac{\partial^2 \bar{W}_{Op}}{\partial I_b \partial II_b} I + \frac{\partial^2 \bar{W}_{Op}}{\partial II_b^2} (I_b I - b) + \frac{\partial^2 \bar{W}_{Op}}{\partial III_b \partial II_b} III_b b^{-1} \right) \\
 & \quad + \frac{\partial \bar{W}_{Op}}{\partial III_b} (b^{-1} \otimes III_b b^{-1} + III_b \frac{\partial b^{-1}}{\partial b}) \\
 & \quad + III_b b^{-1} \otimes \left(\frac{\partial^2 \bar{W}_{Op}}{\partial I_b \partial III_b} I + \frac{\partial^2 \bar{W}_{Op}}{\partial II_b \partial III_b} (I_b I - b) + \frac{\partial^2 \bar{W}_{Op}}{\partial III_b^2} III_b b^{-1} \right), \\
 & = \frac{\partial \bar{W}_{Op}}{\partial II_b} (I \otimes I - \mathbf{I}) + \frac{\partial \bar{W}_{Op}}{\partial III_b} III_b (b^{-1} \otimes b^1 + \frac{\partial b^{-1}}{\partial b}) \\
 & \quad + I \otimes \left(\frac{\partial^2 \bar{W}_{Op}}{\partial I_b^2} I + \frac{\partial^2 \bar{W}_{Op}}{\partial II_b \partial I_b} (I_b I - b) + \frac{\partial^2 \bar{W}_{Op}}{\partial III_b \partial I_b} III_b b^{-1} \right. \\
 & \quad \left. + (I_b I - b) \otimes \left(\frac{\partial^2 \bar{W}_{Op}}{\partial I_b \partial II_b} I + \frac{\partial^2 \bar{W}_{Op}}{\partial II_b^2} (I_b I - b) + \frac{\partial^2 \bar{W}_{Op}}{\partial III_b \partial II_b} III_b b^{-1} \right) \right. \\
 & \quad \left. + III_b b^{-1} \otimes \left(\frac{\partial^2 \bar{W}_{Op}}{\partial I_b \partial III_b} I + \frac{\partial^2 \bar{W}_{Op}}{\partial II_b \partial III_b} (I_b I - b) + \frac{\partial^2 \bar{W}_{Op}}{\partial III_b^2} III_b b^{-1} \right) \right), \\
 & = -\left(\frac{\partial \bar{W}_{Op}}{\partial II_b} \mathbf{I} + \frac{1}{2} \frac{\partial \bar{W}_{Op}}{\partial III_b} (\mathbf{I} + \bar{\mathbf{I}}) \right) + I \otimes I \left(\frac{\partial \bar{W}_{Op}}{\partial I_b} + \frac{\partial \bar{W}_{Op}}{\partial III_b} + \frac{\partial^2 \bar{W}_{Op}}{\partial I_b^2} + 4 \frac{\partial^2 \bar{W}_{Op}}{\partial I_b \partial II_b} \right. \\
 & \quad \left. + 2 \frac{\partial^2 \bar{W}_{Op}}{\partial I_b \partial III_b} + 4 \frac{\partial^2 \bar{W}_{Op}}{\partial II_b^2} + 4 \frac{\partial^2 \bar{W}_{Op}}{\partial III_b \partial II_b} + \frac{\partial^2 \bar{W}_{Op}}{\partial III_b^2} \right)|_{(I_b, II_b, III_b)=(3,3,1)}, \\
 & = (15a_1 + 9a_2 - \frac{5}{2}a_4)\mathbf{I} + (9a_1 + 9a_2 - \frac{5a_4}{2})\bar{\mathbf{I}} + I \otimes I (-4a_1 + 20a_2 + 30a_4 + \frac{a_3}{2}).
 \end{aligned}$$

718 **SM6. Proof of Theorem Theorem 4.2.**

719 *Proof.* Let us denote by a the mapping defined by :

$$720 \quad a : \begin{cases} H^3(\Omega, \mathbb{R}^2) \times H^3(\Omega, \mathbb{R}^2) \rightarrow \mathbb{R} \\ (u, v) \mapsto \langle \rho_0(u), \rho_0(v) \rangle_{\mathbb{R}^2, N_0} + \frac{\gamma}{2} \langle \rho_1(\nabla u + \nabla u^T), \rho_1(\nabla v + \nabla v^T) \rangle_{M_2(\mathbb{R}), N_1} \\ + \epsilon(u, v)_{3, \Omega, \mathbb{R}^2} \end{cases},$$

722 with $(., .)_{3, \Omega, \mathbb{R}^2}$ denoting the semi-norm in $H^3(\Omega, \mathbb{R}^2)$, and by L the mapping defined by :

$$723 \quad L : \begin{cases} H^3(\Omega, \mathbb{R}^2) \rightarrow \mathbb{R} \\ v \mapsto \langle \rho_0(f_k), \rho_0(v) \rangle_{\mathbb{R}^2, N_0} + \frac{\gamma}{2} \langle \rho_1(\nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k), \rho_1(\nabla v + \nabla v^T) \rangle_{M_2(\mathbb{R}), N_1} \end{cases}.$$

725 Let us notice that $\forall v \in H^3(\Omega, \mathbb{R}^2)$,

$$\begin{aligned} 726 \quad \mathcal{F}_{\epsilon, k}(v) &= \langle \rho_0(v) - \rho_0(f_k) \rangle_{\mathbb{R}^2, N_0}^2 + \frac{\gamma}{2} \langle \rho_1(\nabla v + \nabla v^T) - \rho_1(\nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R}), N_1}^2 \\ 727 \quad &\quad + \epsilon |v|_{3, \Omega, \mathbb{R}^2}^2, \\ 728 \quad &= \langle \rho_0(v) \rangle_{\mathbb{R}^2, N_0}^2 + \langle \rho_0(f_k) \rangle_{\mathbb{R}^2, N_0}^2 - 2 \langle \rho_0(v), \rho_0(f_k) \rangle_{\mathbb{R}^2, N_0} + \frac{\gamma}{2} \langle \rho_1(\nabla v + \nabla v^T) \rangle_{M_2(\mathbb{R}), N_1}^2 \\ 729 \quad &\quad + \frac{\gamma}{2} \langle \rho_1(\nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R}), N_1}^2 \\ 730 \quad &\quad - \gamma \langle \rho_1(\nabla v + \nabla v^T), \rho_1(\nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R}), N_1} + \epsilon |v|_{3, \Omega, \mathbb{R}^2}^2, \\ 731 \quad &= a(v, v) - 2L(v) + \langle \rho_0(f_k) \rangle_{\mathbb{R}^2, N_0}^2 + \frac{\gamma}{2} \langle \rho_1(\nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R}), N_1}^2. \end{aligned}$$

733 Also problem (4.2) is equivalent to :

$$734 \quad \left\{ \begin{array}{l} \text{Search for } u_{\epsilon, k} \in H^3(\Omega, \mathbb{R}^2) \text{ such that:} \\ \forall v \in H^3(\Omega, \mathbb{R}^2), \forall \mu \in \mathbb{R}, \mathcal{F}_{\epsilon, k}(u_{\epsilon, k}) \leq \mathcal{F}_{\epsilon, k}(u_{\epsilon, k} + \mu v) \end{array} \right..$$

736 Using the bilinearity and symmetry of the mapping a , and the linearity of the mapping L ,
737 one has :

$$\begin{aligned} 738 \quad \mathcal{F}_{\epsilon, k}(u_{\epsilon, k} + \mu v) &= a(u_{\epsilon, k}, u_{\epsilon, k}) + 2\mu a(u_{\epsilon, k}, v) + \mu^2 a(v, v) - 2L(u_{\epsilon, k}) - 2\mu L(v) + \langle \rho_0(f_k) \rangle_{\mathbb{R}^2, N_0}^2 \\ 739 \quad &\quad + \langle \rho_1(\nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R}), N_1}^2, \\ 740 \quad &= \mathcal{F}_{\epsilon, k}(u_{\epsilon, k}) + 2\mu [a(u_{\epsilon, k}, v) - L(v)] + \mu^2 a(v, v), \end{aligned}$$

742 which means, using the previous reformulation that :

$$743 \quad 2\mu [a(u_{\epsilon, k}, v) - L(v)] + \mu^2 a(v, v) \geq 0.$$

745 Assuming now that $\mu > 0$, and dividing the previous inequality by μ , and finally letting μ
746 tend to 0, we get :

$$747 \quad a(u_{\epsilon, k}, v) - L(v) \geq 0.$$

749 Assuming that $\mu < 0$, dividing the inequality by μ and letting μ tend to 0, we finally obtain :

$$750 \quad a(u_{\epsilon, k}, v) - L(v) \leq 0.$$

752 We then deduce that $a(u_{\epsilon, k}, v) = L(v)$, for all $v \in H^3(\Omega, \mathbb{R}^2)$. The inverse is readily obtained
753 thanks to the previous relations. ■

754 **SM7. Proof of Lemma Lemma 4.3.**

755 *Proof.* Let us take $f \in H^3(\Omega, \mathbb{R}^2)$ such that $\|f\|_{A_0, 3, \Omega, \mathbb{R}^2} = 0$. It implies that $|f|_{3, \Omega, \mathbb{R}^2} = 0$
 756 and taking into account the connectedness of Ω , it yields $f \in P^2(\Omega, \mathbb{R}^2)$. We also have
 757 that $\rho_1(\nabla f + \nabla f^T) = 0$ meaning that $\frac{\partial f_1}{\partial x}(b_i) = 0$, $\frac{\partial f_2}{\partial y}(b_i) = 0$, $\frac{\partial f_1}{\partial y}(b_i) + \frac{\partial f_2}{\partial x}(b_i) = 0$, for
 758 $i = 1, \dots, N_1$. Since $f \in P^2(\Omega, \mathbb{R}^2)$, it means that $\frac{\partial f_1}{\partial x} \in P^1(\Omega)$, and $\frac{\partial f_2}{\partial y} \in P^1(\Omega)$. Since A_1
 759 contains a $P^1(\Omega, \mathbb{R}^2)$ -unisolvant subset, it results that $\frac{\partial f_1}{\partial x} \equiv 0$ and $\frac{\partial f_2}{\partial y} \equiv 0$. We therefore
 760 have that

761
$$f_1(x, y) = g_1(y),$$

 762
$$f_2(x, y) = g_1(x).$$

764 Since, $\frac{\partial f_1}{\partial y} + \frac{\partial f_2}{\partial x} \in P^1(\Omega)$ and A_1 contains a $P^1(\Omega, \mathbb{R}^2)$ unisolvant subset, we deduce that

765
$$\frac{\partial f_1}{\partial y} + \frac{\partial f_2}{\partial x} \equiv 0 \Leftrightarrow \frac{\partial f_1}{\partial y} = g'_1(y) \equiv -\frac{\partial f_2}{\partial x} = -g'_2(x), \forall (x, y) \in \Omega.$$

 766

767 It thus means that $f_1(x, y) = ay + b$, $f_2(x, y) = -ax + c$ and are in $P^1(\Omega)$ with $f_1(a_i) =$
 768 $f_2(a_i) = 0$ for all $i = 1, \dots, N_0$. Since A_0 contains a $P^1(\Omega, \mathbb{R}^2)$ -unisolvant subset, we deduce
 769 that $f_1 \equiv f_2 \equiv 0$. It is now clear that $\|\cdot\|_{A_0, 3, \Omega, \mathbb{R}^2}$ is a norm on $H^3(\Omega, \mathbb{R}^2)$ associated with a
 770 scalar product. We now prove the equivalence of the norm $\|\cdot\|_{A_0, \Omega, 3, \mathbb{R}^2}$ with the norm $\|\cdot\|_{3, \Omega, \mathbb{R}^2}$.

771
 772 First, we have $\forall f \in H^3(\Omega, \mathbb{R}^2)$, $\forall a_i \in A_0$, $i = 1, \dots, N_0$, $\forall b_j \in A_1$, $j = 1, \dots, N_1$:

773
$$\langle f(a_i) \rangle_{\mathbb{R}^2} \leq \|f\|_{C^0(\Omega, \mathbb{R}^2)} \leq c \|f\|_{3, \Omega, \mathbb{R}^2}$$
 thanks to Sobolev's embedding,
 774
$$\langle \nabla f(b_j) + \nabla f(b_j)^T \rangle_{M_2(\mathbb{R})} \leq 2 \|f\|_{C^1(\Omega, \mathbb{R}^2)} \leq 2c' \|f\|_{3, \Omega, \mathbb{R}^2}$$
 thanks to Sobolev's embedding,
 775 so,
$$\|f\|_{A_0, 3, \Omega, \mathbb{R}^2} \leq (1 + c^2 N + 4c'^2 N)^{\frac{1}{2}} \|f\|_{3, \Omega, \mathbb{R}^2}.$$

777 Furthermore, let us take $k = 3 = m$, $p = 2$ in Nečas theorem [?, Chapter 2, section 7.1], and
 778 let us take $\rho_0(f)$ and $\rho_1(\nabla f + \nabla f^T)$ as functionals f_i (using the same reasoning as previously
 779 to show the property (7.1 bis) $\forall v \in P^2(\Omega, \mathbb{R}^2)$, $\sum_{i=1}^l |f_i(v)|^2 = 0 \Leftrightarrow v \equiv 0$). Then there exists a
 780 positive constant $c_1 > 0$ such that

781
$$c_1 \|f\|_{3, \Omega, \mathbb{R}^2} \leq [|f|_{3, \Omega, \mathbb{R}^2}^2 + \sum_{i=1}^{N_0} \langle f(a_i) \rangle_{\mathbb{R}^2}^2 + \sum_{j=1}^{N_1} \langle \nabla f(b_j) + \nabla f(b_j)^T \rangle_{M_2(\mathbb{R})}^2]^{\frac{1}{2}},$$

 782

783 which concludes the proof. ■

784 **SM8. Proof of Theorem Theorem 4.4.**

785 *Proof.* The mapping L is a linear and continuous form on $H^3(\Omega, \mathbb{R}^2)$. Indeed,

$$\begin{aligned}
786 \quad |L(v)| &= \left| \sum_{i=1}^{N_0} \langle f_k(a_i), v(a_i) \rangle_{\mathbb{R}^2} + \sum_{j=1}^{N_1} \frac{\gamma}{2} \langle \nabla f_k(b_j) + \nabla f_k(b_j)^T + \nabla f_k(b_j)^T \nabla f_k(b_j), \nabla v(b_j) \right. \\
787 \quad &\quad \left. + \nabla v(b_j)^T \rangle_{M_2(\mathbb{R})} \right|, \\
788 \quad &\leq \sum_{i=1}^{N_0} |\langle f_k(a_i), v(a_i) \rangle_{\mathbb{R}^2}| + \frac{\gamma}{2} \sum_{j=1}^{N_1} |\langle \nabla f_k(b_j) + \nabla f_k(b_j)^T + \nabla f_k(b_j)^T \nabla f_k(b_j), \nabla v(b_j) \rangle_{M_2(\mathbb{R})} \\
789 \quad &\quad + \nabla v(b_j)^T \rangle_{M_2(\mathbb{R})}|, \\
790 \quad &\leq \sum_{i=1}^{N_0} |\langle f_k(a_i) \rangle_{\mathbb{R}^2}| |\langle v(a_i) \rangle_{\mathbb{R}^2}| + \frac{\gamma}{2} \sum_{j=1}^{N_1} |\langle \nabla f_k(b_j) + \nabla f_k(b_j)^T + \nabla f_k(b_j)^T \nabla f_k(b_j) \rangle_{M_2(\mathbb{R})}| \\
791 \quad &\quad |\langle \nabla v(b_j) + \nabla v(b_j)^T \rangle_{M_2(\mathbb{R})}|, \\
792 \quad &\leq \sum_{i=1}^{N_0} |\langle f_k(a_i) \rangle_{\mathbb{R}^2}| \|v\|_{C^0(\Omega, \mathbb{R}^2)} + \gamma \sum_{j=1}^{N_1} |\langle \nabla f_k(b_j) + \nabla f_k(b_j)^T + \nabla f_k(b_j)^T \nabla f_k(b_j) \rangle_{M_2(\mathbb{R})}| \\
793 \quad &\quad \|v\|_{C^1(\Omega, \mathbb{R}^2)}, \\
794 \quad &\leq (cN_0 \|f_k\|_{C^0(\Omega, \mathbb{R}^2)} + c'N_1 \gamma \max_{i=1, \dots, N_1} \langle \nabla f_k(b_i) + \nabla f_k(b_i)^T + \nabla f_k(b_i)^T \nabla f_k(b_i) \rangle_{M_2(\mathbb{R})}) \\
795 \quad &\quad \|v\|_{3, \Omega, \mathbb{R}^2}.
\end{aligned}$$

797 Moreover, the mapping a is a symmetric, bilinear form, continuous on $H^3(\Omega, \mathbb{R}^2) \times H^3(\Omega, \mathbb{R}^2)$.
798 While symmetry and bilinearity are obvious, the continuity of a can be obtained by using the
799 equivalence of norms established in the previous lemma : $\forall (u, v) \in H^3(\Omega, \mathbb{R}^2) \times H^3(\Omega, \mathbb{R}^2)$,

$$\begin{aligned}
800 \quad |a(u, v)| &= \left| \sum_{i=1}^{N_0} \langle u(a_i), v(a_i) \rangle_{\mathbb{R}^2} + \frac{\gamma}{2} \sum_{j=1}^{N_1} \langle \nabla u(b_j) + \nabla u(b_j)^T, \nabla v(b_j) + \nabla v(b_j)^T \rangle_{M_2(\mathbb{R})} + \epsilon(u, v)_{3, \Omega, \mathbb{R}^2} \right|, \\
801 \quad &\leq \max(1, \epsilon, \frac{\gamma}{2}) \left[\sum_{i=1}^{N_0} |\langle u(a_i), v(a_i) \rangle_{\mathbb{R}^2}| + \sum_{j=1}^{N_1} |\langle \nabla u(b_j) + \nabla u(b_j)^T, \nabla v(b_j) + \nabla v(b_j)^T \rangle_{M_2(\mathbb{R})}| \right. \\
802 \quad &\quad \left. + |(u, v)_{3, \Omega, \mathbb{R}^2}| \right], \\
803 \quad &\leq 4 \max(1, \epsilon, \frac{\gamma}{2}) \left[\sum_{i=1}^{N_0} |\langle u(a_i) \rangle_{\mathbb{R}^2} \langle v(a_i) \rangle_{\mathbb{R}^2}| + \sum_{j=1}^{N_1} |\langle \nabla u(b_j) \rangle_{M_2(\mathbb{R})} \langle \nabla v(b_j) \rangle_{M_2(\mathbb{R})}| + |u|_{3, \Omega, \mathbb{R}^2} |v|_{3, \Omega, \mathbb{R}^2} \right].
\end{aligned}$$

■

805 This last inequality proves that a is continuous on $H^3(\Omega, \mathbb{R}^2) \times H^3(\Omega, \mathbb{R}^2)$.
806 To finish with, we prove that a is $H^3(\Omega, \mathbb{R}^3)$ -elliptic. Let $v \in H^3(\Omega, \mathbb{R}^2)$,

$$\begin{aligned}
807 \quad a(v, v) &= \langle \rho_0(u) \rangle_{\mathbb{R}^2, N_0}^2 + \frac{\gamma}{2} \langle \rho_1(\nabla v + \nabla v^T) \rangle_{M_2(\mathbb{R}), N_1}^2 + \epsilon |v|_{3, \Omega, \mathbb{R}^2}^2, \\
808 \quad &\geq \min(1, \epsilon, \frac{\gamma}{2}) \|v\|_{A, 3, \Omega, \mathbb{R}^2}^2.
\end{aligned}$$

810 Using again the equivalence of norms established above, we deduce that a is $H^3(\Omega, \mathbb{R}^2)$ -elliptic.
 811 The Lax-Milgram theorem enables us to conclude that the variational problem (4.3) has a
 812 unique solution denoted by $u_{\epsilon,k}$. ■

813 SM9. Proof of Lemma Lemma 4.5.

814 *Proof.* The Sobolev's embedding gives that

$$815 \quad \exists C_1 > 0, \forall d \in D, \forall f \in H^3(\Omega, \mathbb{R}^2), \|f\|_{A_0^d, 3, \Omega, \mathbb{R}^2} \leq C_1 \|f\|_{3, \Omega, \mathbb{R}^2},$$

817 by using the same arguments as previously, with C_1 independent of d and depending only
 818 on \mathcal{N} . Let us now find a constant C_2 independent of d such that the inequality $\|f\|_{3, \Omega, \mathbb{R}^2} \leq$
 819 $C_2 \|f\|_{A_0^d, 3, \Omega, \mathbb{R}^2}$ holds. For $f \in H^3(\Omega, \mathbb{R}^2)$, we have

$$\begin{aligned} 820 \quad & \frac{1}{2} \sum_{j=1}^{\mathcal{N}} \langle f(b_{0j}) \rangle_{\mathbb{R}^2}^2 = \frac{1}{2} \sum_{i=1}^{\mathcal{N}} \langle f(b_{0j}) - f(a_{0j}^d) + f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2, \\ 821 \quad & = \frac{1}{2} \sum_{i=1}^{\mathcal{N}} \langle f(b_{0j}) - f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2 + \langle f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2 + 2 \langle f(b_{0j}) - f(a_{0j}^d), f(a_{0j}^d) \rangle_{\mathbb{R}^2}, \\ 822 \quad & \leq \sum_{i=1}^{\mathcal{N}} \langle f(b_{0j}) - f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2 + \langle f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2. \end{aligned}$$

824 The open subset Ω having a Lipschitz continuous boundary, the space $H^3(\Omega, \mathbb{R}^2)$ verifies the
 825 Sobolev's Hölder embedding theorem namely $\exists \lambda \in]0, 1]$, $H^3(\Omega, \mathbb{R}^2) \hookrightarrow \mathcal{C}^{0,\lambda}(\bar{\Omega}, \mathbb{R}^2)$. Thus
 826 $f \in \mathcal{C}^{0,\lambda}(\bar{\Omega}, \mathbb{R}^2)$, and $\exists C > 0, \forall j = 1, \dots, \mathcal{N}, \forall d \in D$,

$$\begin{aligned} 827 \quad & \langle f(b_{0j}) - f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2 \leq \|f\|_{\mathcal{C}^{0,\lambda}(\bar{\Omega}, \mathbb{R}^2)}^2 \langle b_{0j} - a_{0j}^d \rangle_{\mathbb{R}^2}^{2\lambda}, \\ 828 \quad & \leq C^2 \|f\|_{H^3(\Omega, \mathbb{R}^2)}^2 \langle b_{0j} - a_{0j}^d \rangle_{\mathbb{R}^2}^{2\lambda} \end{aligned}$$

830 Besides, from the hypotheses, it comes that $\forall j = 1, \dots, \mathcal{N}$

$$831 \quad \forall \beta_j > 0, \exists \eta_{\beta_j} > 0, \forall d \in D, (d \leq \eta_{\beta_j} \Rightarrow \langle a_{0j}^d - b_{0j} \rangle_{\mathbb{R}^2} \leq \beta_{\eta_j}).$$

833 Then $\forall j = 1, \dots, \mathcal{N}$,

$$834 \quad \forall \beta_j > 0, \exists \eta_{\beta_j}, \forall d \in D, (d \leq \eta_{\beta_j} \Rightarrow \langle f(b_{0j}) - f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2 \leq C^2 \|f\|_{H^3(\Omega, \mathbb{R}^2)}^2).$$

836 Let $\beta > 0$ and let us take $\beta_j = \beta$ for all $j = 1, \dots, \mathcal{N}$, and $\eta = \min\{\eta_{\beta_1}, \dots, \eta_{\beta_N}\}$, then

$$\begin{aligned} 837 \quad & d \leq \eta \Rightarrow \sum_{j=1}^{\mathcal{N}} \langle f(b_{0j}) - f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2 \leq C^2 \beta^{2\lambda} \mathcal{N} \|f\|_{H^3(\Omega, \mathbb{R}^2)}^2. \end{aligned}$$

839 This implies that $\forall \beta > 0, \exists \eta > 0, \forall d \in D, \forall f \in H^3(\Omega, \mathbb{R}^2)$,

$$\begin{aligned} 840 \quad & d \leq \eta \Rightarrow \sum_{j=1}^{\mathcal{N}} \langle f(b_{0j}) - f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2 \leq C^2 \beta^{2\lambda} \mathcal{N} \|f\|_{H^3(\Omega, \mathbb{R}^2)}^2. \end{aligned}$$

842 This implies that $\forall \beta > 0, \exists \eta > 0, \forall d \in D, \forall f \in H^3(\Omega, \mathbb{R}^2)$,

$$843 \quad d \leq \eta \Rightarrow \frac{1}{2} \sum_{j=1}^{\mathcal{N}} \langle f(b_{0j}) \rangle_{\mathbb{R}^2}^2 + |f|_{3,\Omega,\mathbb{R}^2}^2 - C^2 \beta^{2\lambda} \mathcal{N} \|f\|_{H^3(\Omega, \mathbb{R}^2)}^2 \leq \sum_{i=1}^{\mathcal{N}} \langle f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2 + |f|_{3,\Omega,\mathbb{R}^2}^2,$$

$$844 \quad d \leq \eta \Rightarrow \frac{1}{2} \sum_{j=1}^{\mathcal{N}} \langle f(b_{0j}) \rangle_{\mathbb{R}^2}^2 + \sum_{i=1}^{N_1} \langle \nabla f(b_i) + \nabla f(b_i)^T \rangle_{M_2(\mathbb{R})}^2 |f|_{3,\Omega,\mathbb{R}^2}^2 - C^2 \beta^{2\lambda} \mathcal{N} \|f\|_{H^3(\Omega, \mathbb{R}^2)}^2$$

$$845 \quad \leq \sum_{i=1}^{\mathcal{N}} \langle f(a_{0j}^d) \rangle_{\mathbb{R}^2}^2 + \sum_{i=1}^{N_1} \langle \nabla f(b_i) + \nabla f(b_i)^T \rangle_{M_2(\mathbb{R})}^2 + |f|_{3,\Omega,\mathbb{R}^2}^2.$$

847 As previously shown, the mapping $f \in H^3(\Omega, \mathbb{R}^2) \mapsto (\frac{1}{2} \sum_{j=1}^{\mathcal{N}} \langle f(b_{0j}) \rangle_{\mathbb{R}^2}^2 + \sum_{i=1}^{N_1} \langle \nabla f(b_i) + \nabla f(b_i)^T \rangle_{M_2(\mathbb{R})}^2 + |f|_{3,\Omega,\mathbb{R}^2}^2)^{\frac{1}{2}}$

848 is a norm on $H^3(\Omega, \mathbb{R}^2)$ equivalent to the norm $\|\cdot\|_{3,\Omega,\mathbb{R}^2}$, so $\forall \beta > 0, \exists \eta > 0, \forall d \in D, \forall f \in H^3(\Omega, \mathbb{R}^2)$,

$$850 \quad d \leq \eta \Rightarrow (C'' - C^2 \beta^{2\lambda} \mathcal{N}) \|f\|_{H^3(\Omega, \mathbb{R}^2)}^2 \leq \|f\|_{A_0^d, 3, \Omega, \mathbb{R}^2}^2.$$

852 By choosing β adequately, the norm equivalence is obtained. ■

853 **SM10. Proof of Theorem Theorem 4.6.**

854 *Proof.* The proof is divided into 3 steps that we detail hereafter :

- **First step :** We start by proving that the sequence $(u_\epsilon^d)_{d \in D \cap [0, \eta]}$ is bounded in $H^3(\Omega, \mathbb{R}^2)$ for fixed N_1 , and ϵ . In the minimisation problem (4.4), let us take $v = f_k$. Then we have

$$\begin{aligned} 858 \quad & \langle \rho^d(u_\epsilon^d - f_k) \rangle_{\mathbb{R}^2, N(d)}^2 + \frac{\gamma}{2} \langle \rho_1(\nabla u_\epsilon^d + (\nabla u_\epsilon^d)^T - \nabla f_k - \nabla f_k^T - \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R}), N_1}^2 + \epsilon |u_\epsilon^d|_{3, \Omega, \mathbb{R}^2}^2 \\ 859 \quad & \leq \epsilon |f_k|_{3, \Omega, \mathbb{R}^2}^2 + \frac{\gamma}{2} \sum_{i=1}^{N_1} \langle \nabla f_k(b_i)^T \nabla f_k(b_i) \rangle_{M_2(\mathbb{R})}^2, \\ 860 \quad & \leq \epsilon |f_k|_{3, \Omega, \mathbb{R}^2}^2 + 2\gamma \sum_{i=1}^{N_1} \langle \nabla f_k(b_i) \rangle_{M_2(\mathbb{R})}^2, \\ 861 \quad & \leq \epsilon |f_k|_{3, \Omega, \mathbb{R}^2}^2 + 2\gamma N_1 \|f_k\|_{C^1(\bar{\Omega}, \mathbb{R}^2)}^2, \end{aligned}$$
■

863 from which we deduce that

$$\begin{cases} |u_\epsilon^d|_{3, \Omega, \mathbb{R}^2}^2 \leq |f_k|_{3, \Omega, \mathbb{R}^2}^2 + \frac{2\gamma}{\epsilon} \|f_k\|_{C^1(\bar{\Omega}, \mathbb{R}^2)}^2 N_1, \\ \langle \rho^d(u_\epsilon^d - f_k) \rangle_{\mathbb{R}^2, N}^2 \leq \epsilon |f_k|_{3, \Omega, \mathbb{R}^2}^2 + 2\gamma N_1 \|f_k\|_{C^1(\Omega, \mathbb{R}^2)}^2, \\ \langle \rho_1(\nabla u_\epsilon^d + (\nabla u_\epsilon^d)^T - \nabla f_k - \nabla f_k^T - \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R}), N_1}^2 \leq \frac{2\epsilon}{\gamma} |f_k|_{3, \Omega, \mathbb{R}^2}^2 + 4N_1 \|f_k\|_{C^1(\bar{\Omega}, \mathbb{R}^2)}^2. \end{cases}$$

866 As $A_0^d \subset A^d$ with A_0^d containing \mathcal{N} elements forming a P^1 -unisolvant set. So one has

$$867 \quad \sum_{a \in A_0^d} \langle u_\epsilon^d(a) - f_k(a) \rangle_{\mathbb{R}^2}^2 \leq \sum_{a \in A^d} \langle u_\epsilon^d(a) - f_k(a) \rangle_{\mathbb{R}^2}^2 \leq \epsilon |f_k|_{3, \Omega, \mathbb{R}^2}^2 + 2\gamma N_1 \|f_k\|_{C^1(\Omega, \mathbb{R}^2)}^2.$$

869 Moreover

$$\begin{aligned}
 870 \quad & \sum_{a \in A_0^d} \langle u_\epsilon^d(a) \rangle_{\mathbb{R}^2}^2 = \sum_{a \in A_0^d} \langle u_\epsilon^d(a) - f_k(a) + f_k(a) \rangle_{\mathbb{R}^2}^2, \\
 871 \quad & \leq 2 \sum_{a \in A_0^d} \langle u_\epsilon^d(a) - f_k(a) \rangle_{\mathbb{R}^2}^2 + \langle f_k(a) \rangle_{\mathbb{R}^2}^2, \\
 872 \quad & \leq 2\epsilon |f_k|_{3,\Omega,\mathbb{R}^2}^2 + 4N_1\gamma \|f_k\|_{C^1(\bar{\Omega},\mathbb{R}^2)}^2 + 2\mathcal{N} \|f_k\|_{C^0(\bar{\Omega},\mathbb{R}^2)}^2, \\
 873 \\
 874 \quad & \sum_{a \in A_1} \langle \nabla u_\epsilon^d(a) + (\nabla u_\epsilon^d(a))^T \rangle_{M_2(\mathbb{R})}^2 = \sum_{a \in A_1} \langle \nabla u_\epsilon^d(a) + \nabla u_\epsilon^d(a)^T - \nabla f_k(a) - \nabla f_k(a)^T \\
 875 \quad & - \nabla f_k(a)^T \nabla f_k(a) + \nabla f_k(a) + \nabla f_k(a)^T + \nabla f_k(a)^T \nabla f_k(a) \rangle_{M_2(\mathbb{R})}^2, \\
 876 \quad & \leq 2 \sum_{a \in A_1} \langle \nabla u_\epsilon^d(a) + \nabla u_\epsilon^d(a)^T - \nabla f_k(a) - \nabla f_k(a)^T - \nabla f_k(a)^T \nabla f_k(a) \rangle_{M_2(\mathbb{R})}^2 \\
 877 \quad & + \langle \nabla f_k(a) + \nabla f_k(a)^T + \nabla f_k(a)^T \nabla f_k(a) \rangle_{M_2(\mathbb{R})}^2, \\
 878 \quad & \leq \frac{2\epsilon}{\gamma} |f_k|_{3,\Omega,\mathbb{R}^2}^2 + 8N_1 \|f_k\|_{C^1(\bar{\Omega},\mathbb{R}^2)}^2 + 32N_1 \|f_k\|_{C^1(\bar{\Omega},\mathbb{R}^2)}^2.
 \end{aligned}$$

881 Finally, using the equivalence of norm from the previous lemmas, and the previous
882 inequalities, we obtain that

$$883 \quad \exists \nu > 0, \forall d \in D, d \leq \eta \Rightarrow \|u_\epsilon^d\|_{H^3(\Omega,\mathbb{R}^2)}^2 \leq \nu.$$

885 The sequence $(u_\epsilon^d)_{d \in D \cap [0,\eta]}$ is bounded in $H^3(\Omega, \mathbb{R}^2)$ independently of d so one can
886 extract a subsequence $(u_\epsilon^{d_l})_{l \in \mathbb{N}}$ with $\lim_{l \rightarrow +\infty} d_l = 0$ (since 0 is an accumulation point of
887 D) that weakly converges to an element of $H^3(\Omega, \mathbb{R}^2)$ denoted by $f_k^* : u_\epsilon^{d_l} \xrightarrow{l \rightarrow +\infty} f_k^*$ in
888 $H^3(\Omega, \mathbb{R}^2)$.

- 889 • **Second step :** In the second step, we prove that $f_k^* = f_k$. Let us assume that $f_k^* \neq f_k$
890 that is, there exists a non-empty open set w included in Ω and a positive real α such
891 that

$$892 \quad \forall x \in w, \langle f_k(x) - f_k^*(x) \rangle_{\mathbb{R}^2} > \alpha.$$

894 Let us now set $\xi = 1 + E \left[\frac{\epsilon |f_k|_{3,\Omega,\mathbb{R}^2}^2 + 2\gamma N_1 \|f_k\|_{C^1(\Omega,\mathbb{R}^2)}^2}{\alpha^2} \right]$, where $E[.]$ denotes the integer
895 part of the argument. Let $B_0 = \{p_{01}, p_{02}, \dots, p_{0\xi}\}$ be a subset of ξ distinct points
896 from w . One has :

$$897 \quad \forall i = 1, \dots, \xi, \exists (p_{0i}^d)_{d \in D}, \forall d \in D, p_{0i}^d \in A^d \text{ and } p_{0i}^d = \lim_{d \rightarrow 0} p_{0i}^d.$$

898 For any $d \in D$, let B_0^d be the set $\{p_{01}^d, \dots, p_{0\xi}^d\}$. As previously proved and taking into
899 account that $B_0^d \subset A^d$, we have

$$900 \quad \sum_{i=1}^{\xi} u_\epsilon^{d_i} (p_{0i}^{d_i} - f_k(p_{0i}^{d_i}))_{\mathbb{R}^2}^2 \leq \epsilon |f_k|_{3,\Omega,\mathbb{R}^2}^2 + 2\gamma N_1 \|f_k\|_{C^1(\bar{\Omega},\mathbb{R}^2)}^2.$$

903 Besides, $\forall i = 1, \dots, \xi$,

$$904 \quad \langle u_\epsilon^{d_l}(p_{0i}^{d_l}) - f_k^*(p_{0i}) \rangle_{\mathbb{R}^2} = \langle u_\epsilon^{d_l}(p_{0i}^{d_l}) - u_\epsilon^{d_l}(p_{0i}) + u_\epsilon^{d_l}(p_{0i}) - f_k^*(p_{0i}) \rangle_{\mathbb{R}^2}, \\ 905 \quad \leq \langle u_\epsilon^{d_l}(p_{0i}^{d_l}) - u_\epsilon^{d_l}(p_{0i}) \rangle_{\mathbb{R}^2} + \langle u_\epsilon^{d_l}(p_{0i}) - f_k^*(p_{0i}) \rangle_{\mathbb{R}^2}$$

907 Since $u_\epsilon^{d_l} \in H^3(\Omega, \mathbb{R}^2) \hookrightarrow \mathcal{C}^{0,\lambda}(\bar{\Omega}, \mathbb{R}^2)$, there exists $C_3 > 0$ such that

$$908 \quad \langle u_\epsilon^{d_l}(p_{0i}^{d_l}) - u_\epsilon^{d_l}(p_{0i}) \rangle_{\mathbb{R}^2} \leq C_3 \langle p_{0i}^{d_l} - p_{0i} \rangle_{\mathbb{R}^2}^\lambda.$$

910 But $\lim_{l \rightarrow +\infty} d_l = 0$, and $p_{0i} = \lim_{l \rightarrow +\infty} p_{0i}^{d_l}$, we thus deduce that

$$911 \quad \lim_{l \rightarrow +\infty} \langle u_\epsilon^{d_l}(p_{0i}^{d_l}) - u_\epsilon^{d_l}(p_{0i}) \rangle_{\mathbb{R}^2} = 0.$$

913 The Rellich-Kondrachov compact embedding theorem gives that $H^3(\Omega, \mathbb{R}^2) \\ 914 \hookrightarrow \mathcal{C}^{0,\lambda}(\bar{\Omega}, \mathbb{R}^2)$. Thus, $(u_\epsilon^{d_l})_{l \in \mathbb{N}}$ uniformly converges to f_k^* and then

$$915 \quad \lim_{l \rightarrow +\infty} \langle u_\epsilon^{d_l}(p_{0i}) - f_k^*(p_{0i}) \rangle_{\mathbb{R}^2} = 0.$$

917 Therefore, we can conclude that

$$918 \quad \lim_{l \rightarrow +\infty} \langle u_\epsilon^{d_l}(p_{0i}^{d_l}) - f_k^*(p_{0i}) \rangle_{\mathbb{R}^2} = 0, \text{ that is } \lim_{l \rightarrow +\infty} u_\epsilon^{d_l}(p_{0i}^{d_l}) = f_k^*(p_{0i}).$$

920 Letting l tend to infinity in the first inequality, we get

$$921 \quad \sum_{l=1}^{\xi} \langle f_k^*(p_{0i}) - f_k(p_{0i}) \rangle_{\mathbb{R}^2}^2 \leq \epsilon |f_k|_{3,\Omega,\mathbb{R}^2}^2 + 2\gamma N_1 \|f_k\|_{\mathcal{C}^1(\bar{\Omega}, \mathbb{R}^2)}^2,$$

923 and so

$$924 \quad \xi \alpha^2 \leq \epsilon |f_k|_{3,\Omega,\mathbb{R}^2}^2 + 2\gamma N_1 \|f_k\|_{\mathcal{C}^1(\bar{\Omega}, \mathbb{R}^2)}^2,$$

926 which is in contradiction with the choice of ξ . Then $f_k^* = f_k$.

- 927 • **Third step :** In the last step, we come back to the initial displacements in $W^{1,\infty}(\Omega, \mathbb{R}^2)$. ■
928 We have

$$929 \quad \lim_{k \rightarrow +\infty, l \rightarrow +\infty} \|u_\epsilon^{d_l} - u\|_{1,\infty} \leq \lim_{k \rightarrow +\infty} (\lim_{l \rightarrow +\infty} \|u_\epsilon^{d_l} - f_k\|_{1,\infty} + \|f_k - u\|_{1,\infty}), \\ 930 \quad \leq 0,$$

932 since $\|f_k - u\|_{1,\infty} \xrightarrow{k \rightarrow +\infty} 0$ by construction and for all $k \in \mathbb{N}$, $\lim_{l \rightarrow +\infty} \|u_\epsilon^{d_l} - f_k\|_{1,\infty}$ as
933 $H^3(\Omega, \mathbb{R}^2) \xrightarrow[c]{} W^{1,\infty}(\Omega, \mathbb{R}^2)$.
934

935 **SM11. Alternative convergence analysis for the third method.** Let us now consider the
936 following problem :

(SM11.1)

$$937 \quad \left\{ \begin{array}{l} \text{Search for } u_\epsilon^d \in H_0^3(\Omega) \text{ such that } \forall t \in H^3(\Omega, \mathbb{R}^2), \\ \langle \rho_0(u_\epsilon^d - f_k) \rangle_{\mathbb{R}^2, N_0}^2 + \frac{\gamma}{2} \langle \rho^d (\nabla u_\epsilon^d + (\nabla u_\epsilon^d)^T - \nabla f_k - \nabla f_k^T - \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R}), N}^2 + \epsilon |u_\epsilon^d|_{3, \Omega, \mathbb{R}^2}^2 . \\ \leq \langle \rho_0(v - f_k) \rangle_{\mathbb{R}^2, N_0}^2 + \frac{\gamma}{2} \langle \rho^d (\nabla v + \nabla v^T - \nabla f_k - \nabla f_k^T - \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R}), N}^2 + \epsilon |v|_{3, \Omega, \mathbb{R}^2}^2 \end{array} \right. .$$

938
939 In this case, the norm $\|\cdot\|_{A^d, 3, \Omega, \mathbb{R}^2}$ is defined by

$$940 \quad \|f\|_{A^d, 3, \Omega, \mathbb{R}^2} = [\langle \rho_0(f_k) \rangle_{\mathbb{R}^2, N_0}^2 + \langle \rho^d (\nabla f + \nabla f^T) \rangle_{M_2(\mathbb{R}), N}^2 + |f|_{3, \Omega, \mathbb{R}^2}^2]^{\frac{1}{2}}.$$

941 As shown in the previous lemma, this norm $\|\cdot\|_{A^d, 3, \Omega, \mathbb{R}^2}$ is equivalent to the norm $\|\cdot\|_{3, \Omega, \mathbb{R}^2}$ on
942 $H^3(\Omega, \mathbb{R}^2)$. We first derive the following lemma.

943 **Lemma SM11.1.** *Let $B_1 = \{b_{01}, \dots, b_{0N}\}$ be a fixed P^1 -unisolvent subset of $\bar{\Omega}$. By hypothesis,
944 0 $\in \bar{D}$ and $\limsup_{d \rightarrow 0} \delta(x, A^d) = 0$ holds so*

$$945 \quad \forall j = 1, \dots, \mathcal{N}, \exists (a_{0j}^d)_{d \in D}, \forall d \in D, a_{0j}^d \in A^d \text{ and } b_{0j} = \lim_{d \rightarrow 0} a_{0j}^d.$$

946 For any $d \in D$, let A_0^d be the set $\{a_{0j}^d, \dots, a_{0N}^d\}$ and let $\|\cdot\|_{A_0^d, 3, \Omega, \mathbb{R}^2}$ be the sum defined by
947 $\forall f \in H^3(\Omega, \mathbb{R}^2)$,

$$948 \quad \|f\|_{A_0^d, 3, \Omega, \mathbb{R}^2} = \left[\sum_{j=1}^{N_0} \langle f(a_j) \rangle_{\mathbb{R}^2}^2 + \sum_{j=1}^{\mathcal{N}} \langle \nabla f(a_{0j}^d) + \nabla f(a_{0j}^d)^T \rangle_{M_2(\mathbb{R})}^2 \right]^{\frac{1}{2}}.$$

949 Then, there exists $\eta > 0$ such that for any $d \leq \eta$, the norm $\|\cdot\|_{A_0^d, 3, \Omega, \mathbb{R}^2}$ is uniformly equivalent
950 over $D \cap [0, \eta]$ to the norm $\|\cdot\|_{3, \Omega, \mathbb{R}^2}$ in $H^3(\Omega, \mathbb{R}^2)$.

951 **Proof.** The Sobolev's embedding gives that

$$952 \quad \exists c_1 > 0, \forall d \in D, \forall f \in H^3(\Omega, \mathbb{R}^2), \|f\|_{A_0^d, 3, \Omega, \mathbb{R}^2} \leq C_1 \|f\|_{3, \Omega, \mathbb{R}^2},$$

953 by using similar arguments as previously. The constant C_1 is independent of d and only
954 depends on N_0 and \mathcal{N} .

955 Let us now find a constant C_2 independent of d such that the inequality

$$956 \quad \|f\|_{3, \Omega, \mathbb{R}^2} \leq C_2 \|f\|_{A_0^d, 3, \Omega, \mathbb{R}^2},$$

957 holds. For $f \in H^3(\Omega, \mathbb{R}^2)$,

$$\begin{aligned} 958 \quad & \frac{1}{2} \sum_{j=1}^{\mathcal{N}} \langle \nabla f(b_{0j}) + \nabla f(b_{0j})^T \rangle_{\mathbb{R}^2}^2 = \frac{1}{2} \sum_{j=1}^{\mathcal{N}} \langle \nabla f(b_{0j}) + \nabla f(b_{0j})^T - \nabla f(a_{0j}^d) - \nabla f(a_{0j}^d)^T \\ 959 \quad & + \nabla f(a_{0j}^d) + \nabla f(a_{0j}^d)^T \rangle_{M_2(\mathbb{R})}^2, \\ 960 \quad & \leq \sum_{j=1}^{\mathcal{N}} \langle \nabla f(b_{0j}) + \nabla f(b_{0j})^T - \nabla f(a_{0j}^d) - \nabla f(a_{0j}^d)^T \rangle_{M_2(\mathbb{R})}^2 + \langle \nabla f(a_{0j}^d) + \nabla f(a_{0j}^d)^T \rangle_{M_2(\mathbb{R})}^2. \end{aligned}$$

967 The open subset Ω having a Lipschitz-continuous boundary and the space $H^3(\Omega, \mathbb{R}^2)$ verifies
968 the Sobolev's Hölder embedding theorem, namely

969 $\exists \lambda \in]0, 1], H^3(\Omega, \mathbb{R}^2) \xrightarrow{c} \mathcal{C}^{1,\lambda}(\bar{\Omega}, \mathbb{R}^2).$
970

971 Thus $f \in \mathcal{C}^{1,\lambda}(\Omega, \mathbb{R}^2)$ and $\exists C > 0, \forall j = 1, \dots, \mathcal{N}, \forall d \in D,$

972 $\langle \nabla f(b_{0j}) + \nabla f(b_{0j})^T - \nabla f(a_{0j}^d) - \nabla f(a_{0j}^d)^T \rangle_{M_2(\mathbb{R})}^2 \leq 2\|f\|_{\mathcal{C}^{1,\lambda}(\Omega, \mathbb{R}^2)}^2 \langle b_{0j} - a_{0j}^d \rangle_{\mathbb{R}^2}^{2\lambda},$
973 $\leq 2C^2\|f\|_{3,\Omega, \mathbb{R}^2}^2 \langle b_{0j} - a_{0j}^d \rangle_{\mathbb{R}^2}^{2\lambda}.$

975 Besides, we also have from the assumptions that for all $j = 1, \dots, \mathcal{N},$

976 $\forall \beta_j > 0, \exists \eta_{\beta_j} > 0, \forall d \in D, (d \leq \eta_{\beta_j} \Rightarrow \langle a_{0j}^d - b_{0j} \rangle_{\mathbb{R}^2} \leq \beta_j).$

978 Then $\forall j = 1, \dots, \mathcal{N}$

979 $\forall \beta_j > 0, \exists \eta_{\beta_j} > 0, \forall d \in D, (d \leq \eta_{\beta_j} \Rightarrow \langle \nabla f(b_{0j}) + \nabla f(b_{0j})^T - \nabla f(a_{0j}^d) - \nabla f(a_{0j}^d)^T \rangle_{M_2(\mathbb{R})}^2$
980 $\leq 2C^2\|f\|_{3,\Omega, \mathbb{R}^2}^2 \beta_j^{2\lambda}.$

982 Let $\delta > 0$ and let us take $\beta_j = \beta, \forall j = 1 \dots, \mathcal{N}, \eta = \min\{\eta_{\beta_1}, \dots, \eta_{\beta_N}\}$, then

983 $\forall d \in D, (d \leq \eta \Rightarrow \sum_{j=1}^{\mathcal{N}} \langle \nabla f(b_{0j}) + \nabla f(b_{0j})^T - \nabla f(a_{0j}^d) - \nabla f(a_{0j}^d)^T \rangle_{M_2(\mathbb{R})}^2 \leq \mathcal{N}C^2\beta^{2\lambda}\|f\|_{3,\Omega, \mathbb{R}^2}^2).$ ■

985 Finally,

986 $\forall \beta > 0, \exists \eta > 0, \forall d \in D, \forall f \in H^3(\Omega, \mathbb{R}^2),$

987 $d \leq \eta \Rightarrow \sum_{j=1}^{\mathcal{N}} \langle \nabla f(b_{0j}) + \nabla f(b_{0j})^T - \nabla f(a_{0j}^d) - \nabla f(a_{0j}^d)^T \rangle_{M_2(\mathbb{R})}^2 \leq C^2\beta^{2\lambda}\mathcal{N}\|f\|_{3,\Omega, \mathbb{R}^2}^2.$

989 This implies that

990 $\forall \beta > 0, \exists \eta > 0, \forall d \in D, \forall f \in H^3(\Omega, \mathbb{R}^2), d \leq \eta$

991 $\Rightarrow \frac{1}{2} \sum_{j=1}^{\mathcal{N}} \langle \nabla f(b_{0j}) + \nabla f(b_{0j}^T) \rangle_{M_2(\mathbb{R})}^2 + \sum_{j=1}^{N_0} \langle f(a_i) \rangle_{\mathbb{R}^2}^2 + \|f\|_{3,\Omega, \mathbb{R}^2}^2 - C^2\beta^{2\lambda}\mathcal{N}\|f\|_{3,\Omega, \mathbb{R}^2}^2 \leq \|f\|_{A_0^d, 3, \Omega, \mathbb{R}^2}^2.$ ■

993 As previously shown, the mapping $f \in H^3(\Omega, \mathbb{R}^2) \mapsto \left[\sum_{j=1}^{N_0} \langle f(a_i) \rangle_{\mathbb{R}^2}^2 + \frac{1}{2} \sum_{j=1}^{\mathcal{N}} \langle \nabla f(b_{0j}) + \nabla f(b_{0j}^T) \rangle_{M_2(\mathbb{R})}^2 \right]$ ■

994 $+ \|f\|_{3,\Omega, \mathbb{R}^2}^2 \right]^{\frac{1}{2}}$ is a norm on $H^3(\Omega, \mathbb{R}^2)$ equivalent to the norm $\|\cdot\|_{3,\Omega, \mathbb{R}^2}$ so

995 $\forall \beta > 0, \exists \eta > 0, \forall d \in D, \forall f \in H^3(\Omega, \mathbb{R}^2),$

996 $d \leq \eta \Rightarrow (C'^2 - C^2\beta^{2\lambda}\mathcal{N})\|f\|_{3,\Omega, \mathbb{R}^2}^2 \leq \|f\|_{3,\Omega, \mathbb{R}^2}^2.$

998 By choosing β adequately, the norm equivalence is obtained. ■

999 Before considering a convergence result, let us first introduce the following problem

1000 (SM11.2)
$$\inf_{u \in H^3(\Omega, \mathbb{R}^2)} |u|_{3,\Omega, \mathbb{R}^2},$$

 1001 s.t. $\nabla u + \nabla u^T = \nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k$ everywhere on Ω ,
 1003 $\rho_0(u) = \rho_0(f_k),$

1004 and prove the existence of minimisers.

1005 **Theorem SM11.2 (Existence of minimisers).** *Let us assume that there exists $u_0 \in H^3(\Omega, \mathbb{R}^2)$
1006 such that*

1007
$$\frac{\partial u_{0,1}}{\partial x} = \frac{\partial f_{k,1}}{\partial x} + \left(\frac{\partial f_{k,1}}{\partial x}\right)^2 + \left(\frac{\partial f_{k,2}}{\partial x}\right)^2,$$

 1008
$$\frac{\partial u_{0,2}}{\partial y} = \frac{\partial f_{k,2}}{\partial y} + \left(\frac{\partial f_{k,2}}{\partial y}\right)^2 + \left(\frac{\partial f_{k,1}}{\partial y}\right)^2,$$

 1009
$$\frac{\partial u_{0,1}}{\partial y} + \frac{\partial u_{0,2}}{\partial x} = \frac{\partial f_{k,1}}{\partial y} + \frac{\partial f_{k,2}}{\partial x} + \frac{\partial f_{k,1}}{\partial x} \frac{\partial f_{k,1}}{\partial y} + \frac{\partial f_{k,2}}{\partial x} \frac{\partial f_{k,2}}{\partial y},$$

 1010

1011 *everywhere on Ω and for all $i = 1, \dots, N_0$, $u_0(a_i) = f_k(a_i)$. (This shouldn't be too restrictive
1012 if N_0 is small enough). Then there is at least one minimiser of the problem (SM11.2).*

1013 *Proof.* Let (u_j) be a minimising sequence of the problem such that $\forall j \in \mathbb{N}$,

1014 $u_j \in H^3(\Omega, \mathbb{R}^2),$
 1015 $\nabla u_j + \nabla u_j^T = \nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k$, everywhere on Ω ,
 1016 $\rho(u_j) = \rho(f_k).$

1018 For j large enough, we have the following coercivity inequality

1019 $|u_j|_{3,\Omega, \mathbb{R}^2} \leq |u_0|_{3,\Omega, \mathbb{R}^2}$, and so
 1020 $\langle \rho(\nabla u_j + \nabla u_j^T) \rangle_{M_2(\mathbb{R})}^2 + \langle \rho(u_j) \rangle_{\mathbb{R}^2}^2 + |u_j|_{3,\Omega, \mathbb{R}^2}^2 \leq \langle \rho(\nabla u_0 + \nabla u_0^T) \rangle_{M_2(\mathbb{R})}^2 + \langle \rho(u_0) \rangle_{\mathbb{R}^2}^2 + |u_0|_{3,\Omega, \mathbb{R}^2}^2.$ ■

1022 Thanks to the previous lemma that gives an equivalence of norms, we get that (u_j) is uniformly
1023 bounded according to j in $H^3(\Omega, \mathbb{R}^2)$. We thus can extract a sub-sequence still denoted by
1024 (u_j) such that

1025 $u_j \xrightarrow{j \rightarrow +\infty} \bar{u}$ in $H^3(\Omega, \mathbb{R}^2),$
 1026

1027 with $\bar{u} \in H^3(\Omega, \mathbb{R}^2)$. Since $H^3(\Omega, \mathbb{R}^2) \hookrightarrow_c \mathcal{C}^1(\bar{\Omega}, \mathbb{R}^2)$, we deduce that $u_k \rightarrow \bar{u}$ in $\mathcal{C}^1(\bar{\Omega}, \mathbb{R}^2)$, and
1028 therefore

1029 $u_j \xrightarrow{j \rightarrow +\infty} \bar{u}$ everywhere in Ω ,
 1030 $\nabla u_j \xrightarrow{j \rightarrow +\infty} \nabla \bar{u}$ everywhere in Ω ,
 1031 $\nabla u_j^T \xrightarrow{j \rightarrow +\infty} \nabla \bar{u}^T$ everywhere in Ω .
 1032

1033 Hence we have that

1034 $\rho(\bar{u}) = \lim_{j \rightarrow +\infty} \rho(u_j) = \lim_{j \rightarrow +\infty} \rho(f_k) = \rho(f_k),$

1035 $\nabla \bar{u}(x) + \nabla \bar{u}^T(x) = \lim_{j \rightarrow +\infty} \nabla u_j(x) + \nabla u_j(x)^T = \lim_{j \rightarrow +\infty} \nabla f_k(x) + \nabla f_k^T(x) + \nabla f_k^T(x) \nabla f_k(x)$
 1036 $= \nabla f_k(x) + \nabla f_k^T(x) + \nabla f_k^T(x) \nabla f_k(x), \text{ for every } x \in \Omega.$

1038 Finally, by the semi-continuity of $|\cdot|_{3,\Omega,\mathbb{R}^2}$, we get that

1039 $|\bar{u}|_{3,\Omega,\mathbb{R}^2} \leq \liminf_{j \rightarrow +\infty} |u_j|_{3,\Omega,\mathbb{R}^2},$

1041 and \bar{u} is a minimiser of the problem (SM11.2). ■

1042 We now establish the following convergence result

1043 **Theorem SM11.3 (Convergence).** *Let \hat{f}_k be a minimiser of problem (SM11.2). For any
 1044 $d \in D$, and $\epsilon \in]0, \epsilon_0]$, we denote by $u_{\epsilon,k}^d$ the unique solution of problem (SM11.1). Then under
 1045 the above assumptions, there exists a subsequence $(u_{\epsilon_l,k}^{d_l})$ with $\lim_{l \rightarrow +\infty} d_l = \lim_{l \rightarrow +\infty} \epsilon_l = 0$, such that*

1046 $u_{\epsilon_l,k}^{d_l} \xrightarrow{l \rightarrow +\infty} \hat{f}_k$

1048 in $H^3(\Omega, \mathbb{R}^2)$ with $\lim_{k \rightarrow +\infty} \lim_{l \rightarrow +\infty} \|\nabla u_{\epsilon_l,k}^{d_l} + (\nabla u_{\epsilon_l,k}^{d_l})^T - \nabla u - \nabla u^T - \nabla u^T \nabla u\|_{L^\infty(\Omega, M_2(\mathbb{R}))} = 0$.

1049 *Proof.* The proof is divided into four steps that we detail hereafter.

1050 • **First step :** We start by proving that the sequence $(u_{\epsilon,k}^d)_{d \in D \cap]0, \eta], \epsilon \in]0, \epsilon_0]}$ is bounded
 1051 in $H^3(\Omega, \mathbb{R}^2)$. In the minimisation problem (SM11.1), let us take $v = \hat{f}_k$. Then we
 1052 have

1053 $\langle \rho(u_{\epsilon,k}^d - f_k) \rangle_{\mathbb{R}^2, N_0}^2 + \frac{\gamma}{2} \langle \rho^d(\nabla u_{\epsilon,k}^d + (\nabla u_{\epsilon,k}^d)^T - \nabla f_k - \nabla f_k^T - \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R})}^2 + \epsilon |u_{\epsilon,k}^d|_{3,\Omega,\mathbb{R}}^2$
 1054 $\leq \epsilon |\hat{f}_k|_{3,\Omega,\mathbb{R}}^2,$ ■

1056 from which we deduce that

1057
$$\begin{cases} |u_{\epsilon,k}^d|_{3,\Omega,\mathbb{R}^2} \leq |\hat{f}_k|_{3,\Omega,\mathbb{R}^2}, \\ \frac{\gamma}{2} \langle \rho^d(\nabla u_{\epsilon,k}^d + (\nabla u_{\epsilon,k}^d)^T - \nabla f_k - \nabla f_k^T - \nabla f_k^T \nabla f_k) \rangle_{M_2(\mathbb{R})}^2 \leq \epsilon |\hat{f}_k|_{3,\Omega,\mathbb{R}^2}^2, \\ \langle \rho_0(u_{\epsilon,k}^d - f_k) \rangle_{\mathbb{R}^2}^2 \leq \epsilon |\hat{f}_k|_{3,\Omega,\mathbb{R}}^2 \leq \epsilon_0 |\hat{f}_k|_{3,\Omega,\mathbb{R}}^2. \end{cases}$$

1059 As $A_0^d \subset A^d$, one has

1060 $\sum_{a \in A_0^d} \langle \nabla u_{\epsilon,k}^d(a) + \nabla u_{\epsilon,k}^d(a)^T - \nabla \hat{f}_k(a) - \nabla \hat{f}_k(a)^T \rangle_{M_2(\mathbb{R})}^2 \leq \epsilon_0 |\hat{f}_k|_{3,\Omega,\mathbb{R}^2}^2.$

1062 Moreover

1063 $\sum_{a \in A_0^d} \langle \nabla u_{\epsilon,k}^d(a) + \nabla u_{\epsilon,k}^d(a)^T \rangle_{M_2(\mathbb{R})}^2 \leq 2 \sum_{a \in A_0^d} \langle \nabla u_{\epsilon,k}^d(a) + \nabla u_{\epsilon,k}^d(a)^T - \nabla \hat{f}_k(a) - \nabla \hat{f}_k(a)^T \rangle_{M_2(\mathbb{R})}^2$
 1064 $+ 2 \sum_{a \in A_0^d} \langle \nabla \hat{f}_k(a) + \nabla \hat{f}_k(a)^T \rangle_{M_2(\mathbb{R})}^2,$
 1066 $\leq 2\epsilon_0 |\hat{f}_k|_{3,\Omega,\mathbb{R}^2}^2 + 4\mathcal{N} \|\hat{f}_k\|_{C^1(\Omega, \mathbb{R}^2)}^2.$ ■

1067 Finally, using the equivalence of norm previously established, and the inequalities
1068 above, we obtain that

$$1069 \exists \nu > 0, \forall d \in D, \forall \epsilon \in]0, \epsilon_0], d \leq \eta \Rightarrow \|u_k^d\|_{3,\Omega,\mathbb{R}^2} \leq \nu.$$

1071 The sequence $(u_{\epsilon,k}^d)_{d \in D \cap [0,\eta], \epsilon \in]0,\epsilon_0]}$ is bounded in $H^3(\Omega, \mathbb{R}^2)$ so one can extract a sub-
1072 sequence $(u_{\epsilon_l,k}^{d_l})$ with $\lim_{l \rightarrow +\infty} d_l = \lim_{l \rightarrow +\infty} \epsilon_l = 0$ that weakly converges to f_k^* in $H^3(\Omega, \mathbb{R}^2)$.

1073 • **Second step:** We now prove that $f_k^* = \hat{f}_k$. Let us assume that $\nabla f_k^* + (\nabla f_k^*)^T \neq$
1074 $\nabla \hat{f}_k + \nabla \hat{f}_k^T$, that is, there exists a non-empty open set ω included in Ω and a positive
1075 real α such that

$$1076 \forall x \in \omega, \langle \nabla f_k^*(x) + (\nabla f_k^*)^T(x) - \nabla \hat{f}_k(x) - \nabla \hat{f}_k^T(x) \rangle_{M_2(\mathbb{R})}^2 > \alpha.$$

1078 Let us set $\xi = E \left[\frac{\epsilon |\hat{f}_k|_{3,\Omega,\mathbb{R}^2}^2}{\alpha^2} \right]$, where $E[.]$ denotes te integer part of the argument. Let
1079 $B_0 = \{p_{01}, \dots, p_{0\xi}\}$ be a subset of ξ distinct points from ω . One has

$$1080 \forall i = 1, \dots, \xi, \exists (p_{0i}^d)_{d \in D}, (\forall d \in D, p_{0i} \in A) \text{ and } p_{0i} = \lim_{d \rightarrow 0} p_{0i}^d.$$

1082 For any $d \in D$, let B_0^d be the set $\{p_{01}^d, \dots, p_{0\xi}^d\}$. As previously proved and taking into
1083 account that $B_0^d \subset A^d$, we have

$$1084 \sum_{i=1}^{\xi} \langle \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d) + \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d)^T - \nabla \hat{f}_k(p_{0i}^d) - \nabla \hat{f}_k(p_{0i}^d)^T \rangle_{M_2(\mathbb{R})}^2 \leq \epsilon |\hat{f}_k|_{3,\Omega,\mathbb{R}^2}^2.$$

1086 Besides, for all $i = 1, \dots, \xi$,

$$\begin{aligned} 1087 & \langle \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d) + \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d)^T - \nabla f_k^*(p_{0i}) - \nabla f_k^*(p_{0i})^T \rangle_{M_2(\mathbb{R})} \\ 1088 &= \langle \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d) + \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d)^T - \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}) - \nabla u_{\epsilon_l,k}^{d_l}(p_{0i})^T + \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}) + \nabla u_{\epsilon_l,k}^{d_l}(p_{0i})^T \\ 1089 &\quad - \nabla f_k^*(p_{0i}) - \nabla f_k^*(p_{0i})^T \rangle_{M_2(\mathbb{R})}, \\ 1090 &\leq \langle \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d) + \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d)^T - \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}) - \nabla u_{\epsilon_l,k}^{d_l}(p_{0i})^T \rangle_{M_2(\mathbb{R})} + \langle \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}) \\ 1091 &\quad + \nabla u_{\epsilon_l,k}^{d_l}(p_{0i})^T - \nabla f_k^*(p_{0i}) - \nabla f_k^*(p_{0i})^T \rangle_{M_2(\mathbb{R})}. \end{aligned}$$

1093 But $u_{\epsilon_l,k}^{d_l} \in H^3(\Omega, \mathbb{R}^2) \overset{c}{\hookrightarrow} C^{1,\lambda}(\Omega, \mathbb{R}^2)$ for $\lambda \in [0, 1]$ and so there exists $C_3 > 0$ such
1094 that

$$1095 \langle \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d) + \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d)^T - \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}) - \nabla u_{\epsilon_l,k}^{d_l}(p_{0i})^T \rangle_{M_2(\mathbb{R})} \leq 2C_3 \langle p_{0i}^d - p_{0i} \rangle_{\mathbb{R}^2}^\lambda.$$

1097 But $\lim_{l \rightarrow +\infty} d_l = 0$ and $p_{0i} = \lim_{l \rightarrow +\infty} p_{0i}^d$, and we deduce that

$$1098 \lim_{l \rightarrow +\infty} \langle \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d) + \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}^d)^T - \nabla u_{\epsilon_l,k}^{d_l}(p_{0i}) - \nabla u_{\epsilon_l,k}^{d_l}(p_{0i})^T \rangle_{M_2(\mathbb{R})} = 0.$$

The Rellich-Kondrachov compact embedding theorem gives that $H^3(\Omega, \mathbb{R}^2) \hookrightarrow_c \mathcal{C}^1(\Omega, \mathbb{R}^2)$, and thus

$$\lim_{l \rightarrow +\infty} \langle \nabla u_{\epsilon_l, k}^{d_l}(p_{0i}) + \nabla u_{\epsilon_l, k}^{d_l}(p_{0i})^T \nabla f_k^*(p_{0i}) - \nabla f_k^*(p_{0i})^T \rangle_{M_2(\mathbb{R})} = 0.$$

Therefore, we can conclude that

$$\lim_{l \rightarrow +\infty} \langle \nabla u_{\epsilon_l, k}^{d_l}(p_{0i}) + \nabla u_{\epsilon_l, k}^{d_l}(p_{0i})^T - \nabla f_k^*(p_{0i}) - \nabla f_k^*(p_{0i})^T \rangle_{M_2(\mathbb{R})} = 0,$$

and so $\lim_{l \rightarrow +\infty} \nabla u_{\epsilon_l, k}^{d_l}(p_{0i}) + \nabla u_{\epsilon_l, k}^{d_l}(p_{0i})^T = \nabla f_k^*(p_{0i}) + \nabla f_k^*(p_{0i})^T$. Letting l tend to infinity, it comes

$$\sum_{i=1}^{\xi} \langle \nabla f_k^*(p_{0i}) + \nabla f_k^*(p_{0i})^T - \nabla \hat{f}_k(p_{0i}) - \nabla \hat{f}_k(p_{0i})^T \rangle_{M_2(\mathbb{R})}^2 \leq \epsilon |\hat{f}_k|_{3, \Omega, \mathbb{R}^2}^2,$$

and so $\alpha^2 \xi \leq \epsilon |\hat{f}_k|_{3, \Omega, \mathbb{R}^2}^2$ which is in contradiction with the definition of ξ . Consequently, $\nabla f_k^* + (\nabla f_k^*)^T = \nabla \hat{f}_k + \nabla \hat{f}_k^T = \nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k$ everywhere on Ω . Also, since $H^3(\Omega, \mathbb{R}^2) \hookrightarrow_c \mathcal{C}^0(\bar{\Omega}, \mathbb{R}^2)$, we have that $\lim_{l \rightarrow +\infty} \langle \rho_0(u_{\epsilon_l, k}^{d_l} - f_k) \rangle_{\mathbb{R}^2, N_0}^2 = \langle \rho_0(f_k^* - f_k) \rangle_{\mathbb{R}^2, N_0}^2$, and by letting l tend to infinity in the last inequality we get that

$$\langle \rho_0(f_k^* - f_k) \rangle_{\mathbb{R}^2, N_0}^2 \leq \lim_{l \rightarrow +\infty} \epsilon_l |\hat{f}_k|_{3, \Omega, \mathbb{R}^2}^2 = 0.$$

Finally, by weak lower semicontinuity of the seminorm in $H^3(\Omega, \mathbb{R}^2)$, we get that

$$|f_k^*|_{3, \Omega, \mathbb{R}^2} \leq \liminf_{l \rightarrow +\infty} |u_{\epsilon_l, k}^{d_l}|_{3, \Omega, \mathbb{R}^2} \leq |\hat{f}_k|_{3, \Omega, \mathbb{R}^2}.$$

Thus f_k^* is a minimiser of problem (SM11.2). Without loss of generality, we say $f_k^* = \hat{f}_k$.

- **Third step:** We then have

$$\begin{aligned} & \| \nabla u_{\epsilon_l, k}^{d_l} + (\nabla u_{\epsilon_l, k}^{d_l})^T - \nabla u - \nabla u^T - \nabla u^T \nabla u \|_{L^\infty(\Omega, M_2(\mathbb{R}))} \\ & \leq \| \nabla u_{\epsilon_l, k}^{d_l} + (\nabla u_{\epsilon_l, k}^{d_l})^T - \nabla \hat{f}_k - \nabla \hat{f}_k^T \|_{L^\infty(\Omega, M_2(\mathbb{R}))} + \| \nabla \hat{f}_k + \nabla \hat{f}_k^T - \nabla u - \nabla u^T \\ & \quad - \nabla u^T \nabla u \|_{L^\infty(\Omega, M_2(\mathbb{R}))}, \\ & \leq \| \nabla u_{\epsilon_l, k}^{d_l} + (\nabla u_{\epsilon_l, k}^{d_l})^T - \nabla \hat{f}_k - \nabla \hat{f}_k^T \|_{L^\infty(\Omega, M_2(\mathbb{R}))} + \| \nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k - \nabla u \\ & \quad - \nabla u^T - \nabla u^T \nabla u \|_{L^\infty(\Omega, M_2(\mathbb{R}))}, \end{aligned}$$

with $\lim_{k \rightarrow +\infty} \lim_{l \rightarrow +\infty} \| \nabla u_{\epsilon_l, k}^{d_l} + (\nabla u_{\epsilon_l, k}^{d_l})^T - \nabla \hat{f}_k - \nabla \hat{f}_k^T \|_{L^\infty(\Omega, M_2(\mathbb{R}))} = 0$ from what precedes and the Sobolev embedding $H^3(\Omega, \mathbb{R}^2) \hookrightarrow_c W^{1, \infty}(\Omega, \mathbb{R}^2)$, and $\lim_{k \rightarrow +\infty} \lim_{l \rightarrow +\infty} \| \nabla f_k + \nabla f_k^T + \nabla f_k^T \nabla f_k - \nabla u - \nabla u^T - \nabla u^T \nabla u \|_{L^\infty(\Omega, M_2(\mathbb{R}))} = 0$, by construction of the sequence f_k . We thus have $\lim_{k \rightarrow +\infty} \lim_{l \rightarrow +\infty} \| \nabla u_{\epsilon_l, k}^{d_l} + (\nabla u_{\epsilon_l, k}^{d_l})^T - \nabla u - \nabla u^T - \nabla u^T \nabla u \|_{L^\infty(\Omega, M_2(\mathbb{R}))} = 0$.

- 1133 • **Fourth step** : We aim at proving that the sequence $(u_{\epsilon_l,k}^{d_l})$ strongly converges to \hat{f}_k
1134 in $H^3(\Omega, \mathbb{R}^2)$. The Rellich-Kondrachov compact embedding theorem gives that

1135 $\forall r, r' \in \mathbb{R}, r > r', H^r(\Omega, \mathbb{R}^2) \xrightarrow{c} H^{r'}(\Omega, \mathbb{R}^2).$

1137 In our case, it means that the sequence $(u_{\epsilon_l,k}^{d_l})$ that weakly converges to \hat{f}_k in $H^3(\Omega, \mathbb{R}^2)$,
1138 strongly converges to \hat{f}_k in $H^2(\Omega, \mathbb{R}^2)$. We thus just need to prove that

1139 $\lim_{l \rightarrow +\infty} |u_{\epsilon_l,k}^{d_l} - \hat{f}_k|_{3,\Omega,\mathbb{R}^2} = 0.$

1141 One has

1142 $|u_{\epsilon_l,k}^{d_l} - \hat{f}_k|_{3,\Omega,\mathbb{R}^2} = |u_{\epsilon_l,k}^{d_l}|_{3,\Omega,\mathbb{R}^2} + |\hat{f}_k|_{3,\Omega,\mathbb{R}^2} - 2(u_{\epsilon_l,k}^{d_l}, \hat{f}_k)_{3,\Omega,\mathbb{R}^2},$
1143 $\leq 2|\hat{f}_k|_{3,\Omega,\mathbb{R}^2} - 2(u_{\epsilon_l,k}^{d_l}, \hat{f}_k)_{3,\Omega,\mathbb{R}^2},$

1145 so $\lim_{l \rightarrow +\infty} |u_{\epsilon_l,k}^{d_l} - \hat{f}_k|_{3,\Omega,\mathbb{R}^2} = 0$, and finally $\lim_{l \rightarrow +\infty} \|u_{\epsilon_l,k}^{d_l} - \hat{f}_k\|_{3,\Omega,\mathbb{R}^2} = 0$ which concludes
1146 the proof. ■

1147 **SM12. Numerical Resolution of the Third Approach.** We now turn to the discretisation
1148 of the variational problem associated with [Theorem 4.4](#) in which u is a substitute for f_k . To
1149 do so, we use standard notations of the finite element theory similar to those in [?, ?]. Let \mathcal{H}
1150 be an open bounded subset of $]0, +\infty[$ admitting 0 as accumulation point. Let us recall that
1151 the elements of class $\mathcal{C}^{k'}$ can be used for the computation of discrete D^m -splines (in our case,
1152 $m = 3$) with $m \leq k' + 1$. As a consequence, $(k', m) = (2, 3)$ is a suitable combination. For
1153 all $n \in \mathbb{N}$ and for all subsets E of \mathbb{R}^2 , $Q_l(E)$ denotes the space of the restrictions of E of the
1154 polynomial functions over \mathbb{R}^2 of degree $\leq l$ with respect to each variable. $\forall h \in \mathcal{H}$, let $(V_h)^2$
1155 be the subspace of $H^3(\Omega, \mathbb{R}^2)$ of finite dimension with $(V_h)^2 \hookrightarrow \mathcal{C}^1(\bar{\Omega}, \mathbb{R}^2)$. The reference finite
1156 element is the Bogner-Fox-Schmit \mathcal{C}^2 rectangle denoted by (K, P_K, Σ_K) .

1157 Let $(v^q)_{q=1,2}$ be the components of $v \in H^3(\Omega, \mathbb{R}^2)$ and $w_i = \begin{pmatrix} w_{i,11} & w_{i,12} \\ w_{i,21} = w_{i,12} & w_{i,22} \end{pmatrix} :=$
1158 $\nabla u(a_i) + \nabla u(a_i)^T + \nabla u(a_i)^T \nabla u(a_i)$, $\forall i \in \{1, \dots, N\}$. Let also $(x_i^q)_{q=1,2}$ be the compo-
1159 nents of $u(a_i)$, $\forall i \in \{1, \dots, N\}$. Now let M_h be the dimension of V_h and $\{P_j^h\}_{j=1, \dots, M_h}$ be
1160 basis functions. If we denote by u_ϵ^h the solution of the variational problem associated with
1161 [Theorem 4.4](#) and approximated in $(V_h)^2$, we can thus decompose $u_\epsilon^h = (u_\epsilon^{h,q})_{q=1,2}$ into:

1162 $\forall q = 1, 2, \exists (\alpha_j^q)_{j=1, \dots, M_h} \in \mathbb{R}, u_\epsilon^{h,q} = \sum_{j=1}^{M_h} \alpha_j^q P_j^h.$

1163 Denoting by $A^h = \left(\frac{\partial P_j^h}{\partial x}(a_i) \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq M_h}}$, $B^h = \left(\frac{\partial P_j^h}{\partial y}(a_i) \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq M_h}} \in (M_{N \times M_h}(\mathbb{R}))^2$, $C^h =$
1164 $(P_j^h(a_i))_{\substack{1 \leq i \leq N \\ 1 \leq j \leq M_h}} \in M_{N \times M_h}(\mathbb{R})$ and $R^h = \left((P_j^h, P_i^h)_{3,\Omega,\mathbb{R}^2} \right)_{1 \leq i, j \leq M_h}$ and taking successively
1165

1166 in the variational problem $v = \begin{pmatrix} P_l^h \\ 0 \end{pmatrix}$, $l = 1, \dots, M_h$ and then $v = \begin{pmatrix} 0 \\ P_l^h \end{pmatrix}$, $l = 1, \dots, M_h$, the
1167 problem amounts to solving the following linear system

$$1168 \quad M_h \begin{pmatrix} & M_h \\ \left(\begin{array}{c|c} 2\gamma(A^h)^T A^h + \gamma(B^h)^T B^h & \gamma(B^h)^T A^h \\ \hline & \gamma(A^h)^T B^h + 2\gamma(B^h)^T B^h \end{array} \right) & \begin{pmatrix} \alpha^1 \\ \hline \alpha^2 \end{pmatrix} = \begin{pmatrix} \gamma(A^h)^T w_{11} + \gamma(B^h)^T w_{12} \\ \hline \gamma(A^h)^T w_{12} + \gamma(B^h)^T w_{22} \end{pmatrix}. \\ M_h & \end{pmatrix}$$

1169
1170