



HAL
open science

Linear relationship between effective radius and precipitation water content near the top of convective clouds: measurement results from ACRIDICON–CHUVA campaign

Ramon Campos Braga, Daniel Rosenfeld, Ovid Krüger, Barbara Ervens, Bruna Holanda, Manfred Wendisch, Trismono Krisna, Ulrich Pöschl, Meinrat Andreae, Christiane Voigt, et al.

► To cite this version:

Ramon Campos Braga, Daniel Rosenfeld, Ovid Krüger, Barbara Ervens, Bruna Holanda, et al.. Linear relationship between effective radius and precipitation water content near the top of convective clouds: measurement results from ACRIDICON–CHUVA campaign. *Atmospheric Chemistry and Physics*, 2021, 21 (18), pp.14079-14088. 10.5194/acp-21-14079-2021 . hal-03483606

HAL Id: hal-03483606

<https://uca.hal.science/hal-03483606v1>

Submitted on 18 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Linear relationship between effective radius and precipitation water content near the top of convective clouds: measurement results from ACRIDICON–CHUVA campaign

Ramon Campos Braga¹, Daniel Rosenfeld², Ovid O. Krüger¹, Barbara Ervens³, Bruna A. Holanda¹, Manfred Wendisch⁴, Trismono Krisna⁴, Ulrich Pöschl¹, Meinrat O. Andreae^{1,5,6}, Christiane Voigt^{7,8}, and Mira L. Pöhlker^{1,a}

¹Multiphase Chemistry Department, Max Planck Institute for Chemistry, 55128 Mainz, Germany

²Institute of Earth Sciences, The Hebrew University of Jerusalem, 9190401 Jerusalem, Israel

³Université Clermont Auvergne, CNRS, SIGMA Clermont, Institut de Chimie de Clermont-Ferrand, 63000 Clermont-Ferrand, France

⁴Leipziger Institut für Meteorologie (LIM), Universität Leipzig, Stephanstr. 3, 04103 Leipzig, Germany

⁵Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA 92037, USA

⁶Department of Geology and Geophysics, King Saud University, 11457 Riyadh, Saudi Arabia

⁷Institute of Atmospheric Physics, German Aerospace Center (DLR), 82234 Oberpfaffenhofen, Germany

⁸Institut für Physik der Atmosphäre, Johannes Gutenberg-Universität, 55099 Mainz, Germany

^anow at: Department of Experimental Aerosol and Cloud Microphysics, Leibniz Institute for Tropospheric Research, 04318 Leipzig, Germany

Correspondence: Ramon Campos Braga (r.braga@mpic.de) and Mira L. Pöhlker (poehlker@tropos.de)

Received: 6 April 2021 – Discussion started: 15 April 2021

Revised: 26 July 2021 – Accepted: 28 July 2021 – Published: 22 September 2021

Abstract. Quantifying the precipitation within clouds is a crucial challenge to improve our current understanding of the Earth's hydrological cycle. We have investigated the relationship between the effective radius of droplets and ice particles (r_e) and precipitation water content (PWC) measured by cloud probes near the top of growing convective cumuli. The data for this study were collected during the ACRIDICON–CHUVA campaign on the HALO research aircraft in clean and polluted conditions over the Amazon Basin and over the western tropical Atlantic in September 2014. Our results indicate a threshold of $r_e \sim 13 \mu\text{m}$ for warm rain initiation in convective clouds, which is in agreement with previous studies. In clouds over the Atlantic Ocean, warm rain starts at smaller r_e , likely linked to the enhancement of coalescence of drops formed on giant cloud condensation nuclei. In cloud passes where precipitation starts as ice hydrometeors, the threshold of r_e is also shifted to values smaller than $13 \mu\text{m}$ when coalescence processes are suppressed and precipitating particles are formed by accretion. We found a statisti-

cally significant linear relationship between PWC and r_e for measurements at cloud tops, with a correlation coefficient of ~ 0.94 . The tight relationship between r_e and PWC was established only when particles with sizes large enough to precipitate (drizzle and raindrops) are included in calculating r_e . Our results emphasize for the first time that r_e is a key parameter to determine both initiation and amount of precipitation at the top of convective clouds.

1 Introduction

Convective cloud formation and precipitation processes have different characteristics depending on the atmospheric thermodynamic conditions and aerosol particle concentration (Reutter et al., 2009; Rosenfeld et al., 2008; Tao et al., 2012). In clean air masses, low concentrations of cloud condensation nuclei (CCN) lead to clouds with relatively fewer droplets ($\sim 50\text{--}200 \text{cm}^{-3}$) at cloud base but with larger sizes

(Twomey, 1974; Andreae et al., 2004; Rosenfeld et al., 2008; Braga et al., 2017a; Sorooshian et al., 2019). These droplets initially grow fast by condensation and subsequently coalesce rapidly into raindrops. In polluted air masses, high concentrations of CCN produce clouds with high concentrations of small drops at cloud base, which can exceed 1000 cm^{-3} . The small and numerous drops grow slowly by condensation due to the high competition for water vapor. In such a case, the coalescence of cloud drops into raindrops is suppressed, and thus, raindrop formation takes place from the melting of ice particles (Andreae et al., 2004; Braga et al., 2017b; Khain et al., 2008; Rosenfeld et al., 2008; Berg et al., 2008).

Over the tropics, convective clouds and mesoscale convective systems account for most of precipitation and severe weather (Liu et al., 2007; Roca et al., 2014; Zipser et al., 2006). In the Amazon Basin, the formation and development of precipitation-forming processes of convective clouds occur at different levels of atmospheric pollution (Andreae et al., 2004; Pöhlker et al., 2016). Previous studies (such as Roberts et al., 2001; Martin et al., 2016) have shown that during the wet season (February–May), low concentrations of CCN particles, mainly consisting of forest biogenic aerosols, are found in the Amazon Basin ($\sim 200\text{--}300\text{ cm}^{-3}$ for 1% of supersaturation), leading to the formation of shallow convective clouds with low ice water content and lightning activity (Albrecht et al., 2011; Williams et al., 2002). These characteristics of the Amazonian clouds and CCN concentrations during the wet season led several authors to refer to this region as a “green ocean” to highlight its similarity with maritime-like regions (e.g., Pöhlker et al., 2016; Roberts et al., 2001; Martin et al., 2016). On the other hand, during the dry season (August–October), the background concentrations of CCN over the Amazon can reach values ~ 10 times higher than those of the green ocean (Artaxo et al., 2002, 2013; Pöhlker et al., 2018; Roberts et al., 2003). The increase of particle concentrations results from forest, savanna, and agricultural fires that release large amounts of biomass-burning aerosols over the pristine rain forest (Andreae et al., 1988). Such conditions inhibit the formation of shallow precipitating clouds and invigorate the ice processes within convective clouds and their lightning activity (Albrecht et al., 2011; Williams et al., 2002; Rosenfeld et al., 2008).

Braga et al. (2017b) have described the general characteristics of growing convective cumulus formed over the Amazon Basin and Atlantic Ocean based on in situ measurements. The measurements were performed during the ACRIDICON–CHUVA (Aerosol, Cloud, Precipitation, and Radiation Interactions and Dynamics of Convective Cloud Systems–Cloud Processes of the Main Precipitation Systems in Brazil: A Contribution to Cloud Resolving Modeling and to the Global Precipitation measurements) campaign in the Amazonian dry season in September 2014 (Wendisch et al., 2016). During the campaign, cloud-profiling flights were performed in regions of different pollution levels and thermo-

dynamic conditions. Braga et al. (2017b) showed that the heights of cloud base are higher over the continental Amazon due to the smaller relative humidity in comparison with the maritime region. Convective clouds formed over the Atlantic Ocean near the Brazilian coast have smaller cloud droplet concentrations at cloud base due to the smaller concentration of aerosol and updraft velocities below cloud base. For convective clouds formed over forested and deforested regions, larger aerosol concentration and updrafts were observed below cloud base, leading to larger droplets activated at cloud base. The precipitating particles were formed mostly by coalescence of drops at temperatures above 0°C over ocean. Over the forest, lightly polluted air masses were found and the precipitation initiation (liquid raindrops) was observed near 0°C . For very polluted air masses, found over the deforestation arc region, the collision and coalescence processes were totally suppressed and the formation of precipitating particles took place at higher altitudes as ice hydrometeors. In these cases, precipitating particles were formed mostly by accretion processes at temperatures below 0°C , when the growth of ice hydrometeors took place from collision with supercooled drops that freeze completely or partially upon contact.

The relationship between particle sizes and precipitation is associated with the coalescence rate which increases in direct proportion to the cloud droplet effective radius (r_{cc}^5) (Freud and Rosenfeld, 2012). Previous studies have found r_{cc} between 13 and $14\text{ }\mu\text{m}$ as a suitable threshold for precipitation initiation (Freud and Rosenfeld, 2012; Rosenfeld and Gutman, 1994; Braga et al., 2017b). The relation between rain initiation and r_{cc} is associated with the increase of both the drop-swept volume and collision efficiency. The collision efficiency of drops increases as a function of their sizes (Khain and Pinsky, 2018). For raindrops, this value is close to unity and is several times larger than that for small drops ($r < 10\text{ }\mu\text{m}$) (Pinsky et al., 2012). The collision and coalescence processes of liquid drops and the accretions processes at supercooled temperatures have a strong effect on the broadening of the particle size distribution and thus particle sizes. In this study, we have investigated measurements of the effective radius (r_e) of cloud particles and the rain and ice precipitation water content (PWC) using data from cloud probes, measured at the cloud tops of growing convective cumulus during the ACRIDICON–CHUVA campaign. We focus our analysis on flights in which precipitation was found in the cloud tops of convective clouds. Our findings shown in the next sections describe the tight relationship between r_e and PWC for in situ measurements in cloud tops in different pollution states and temperature levels. We show that r_e determines both the initiation and amount of precipitation at the top of convective clouds.

2 Methods

2.1 Data and instrumentation

2.1.1 Research flights

The data used in this study are droplet and ice particle concentrations measured in convective clouds by cloud probes mounted on the HALO aircraft during the ACRIDICON–CHUVA campaign (Wendisch et al., 2016). The HALO aircraft was equipped with a meteorological sensor system (Basic HALO Measurement And Sensor System – BAHAMAS) located at the nose of the aircraft (Wendisch et al., 2016). The description of typical meteorological measurements can be found in Mallaun et al. (2015). The HALO flights took place over the Amazon region under various conditions of aerosol concentrations and land cover. Figure 1 shows the flight tracks of cloud profiling flights where precipitation was observed in convective clouds (Braga et al., 2017b). The region of measurements is indicated by circles for each flight. Convective clouds formed in clean air masses were found above the Atlantic Ocean during flight AC19 (in blue, Fig. 1). Flights AC09 and AC18 took place in lightly polluted conditions over the tropical rain forest (in green, Fig. 1). Clouds forming in deforested regions in very polluted (biomass-burning) environments were measured during flights AC07 and AC13 (in red and orange, Fig. 1).

2.1.2 Cloud particle measurements

Cloud particle number concentrations and size distributions were measured by the Cloud Combination Probe (CCP) mounted on board HALO. The CCP combines two detectors, the Cloud Droplet Probe (CDP) and the grayscale Cloud Imaging Probe (CIPgs). The CDP is an open-path instrument that detects forward-scattered laser light from cloud particles as they pass through the detection area (Lance et al., 2010). The CIP records 2-D shadow-cast images of cloud elements. The identification of water drops and ice hydrometeors was performed by Braga et al. (2017b) from the occurrence of visually spherical and non-spherical shapes of the shadows. The combination of CCP–CDP and CCP–CIPgs information provides the ability to measure particles within clouds for nearly the same air sample volume.

Cloud particle size distributions (DSDs) between 3 and 960 μm in diameter were measured at a temporal resolution of 1 s by the CCP–CDP and CCP–CIPgs (Brennguier et al., 2013; Weigel et al., 2016). Each DSD spectrum represents 1 s of flight path (covering between 63 and 112 m of horizontal distance at the aircraft speed). Details about the cloud probe measurement characteristics during ACRIDICON–CHUVA campaign are described in Wendisch et al. (2016), Weigel et al. (2016), and Braga et al. (2017a, b). In this study, a cloud pass is assumed when the total water content (TWC) exceeds 0.05 g m^{-3} and the number concentration of drops




Figure 1. HALO flight tracks during the ACRIDICON–CHUVA experiment. The flight number is indicated at the top by colors. Colored circles indicate the region of cloud profiling in each flight. Convective clouds formed in clean air masses were found above the Atlantic Ocean during flight AC19 (shown in blue). Clouds in lightly pollution conditions were found during flights AC09 and AC18 over the tropical rain forest (shown in green). Polluted clouds forming in deforested regions were measured during flights AC07 and AC13 (shown in red and orange, respectively). The average aerosol particle concentrations measured near cloud bases during flights AC07, AC09, AC13, AC18, and AC19 were 2498 , 821 , 4093 , 744 , and 465 cm^{-3} , respectively (Cecchini et al., 2017).

(N_d) exceeds 20 cm^{-3} . This criterion was applied to avoid cloud passes well mixed with subsaturated environment air ($\text{RH} < 100 \%$) and counts of haze particles, typically found at cloud edges and dissipating convective clouds during the ACRIDICON–CHUVA campaign (Braga et al., 2017b). The N_d and TWC are defined as

$$N_d = \int_{1.5 \mu\text{m}}^{480 \mu\text{m}} N(r) dr \quad (1)$$

and

$$\text{TWC} = \frac{4\pi}{3} \rho \int_{1.5 \mu\text{m}}^{480 \mu\text{m}} r^3 N(r) dr, \quad (2)$$

where N is the particle number concentration (cm^{-3}), ρ is the particle density, and r the particle radius (μm).

2.2 Analysis of cloud properties

Previous studies (e.g., Freud and Rosenfeld, 2012; Braga et al., 2017b) have calculated r_c using data of particle number concentration with radii between 1.5 and 25 μm (r_{ec}), which

does not include precipitating particles. Here, the relationship between cloud particle sizes and PWC is investigated by calculating r_e taking into account the concentration of particles with precipitating sizes ($1.5 \mu\text{m} < r \leq 480 \mu\text{m}$). Precipitating particles are considered those with terminal fall speeds large enough ($\gtrsim 0.5 \text{ m s}^{-1}$) to survive evaporative dissipation over a distance of the order of several hundred meters (Beard, 1976). Droplets smaller than drizzle particles fall slowly enough from most clouds that they evaporate before reaching the ground. The size range of the PWC calculation includes particles with drizzle ($25 \mu\text{m} \leq r \leq 125 \mu\text{m}$) and raindrop ($125 \mu\text{m} < r \leq 480 \mu\text{m}$) sizes. The drizzle water content (DWC) and PWC are calculated using the size range of drizzle and raindrop in Eq. (2). Precipitating particles in this size range ($r > 25 \mu\text{m}$) were often imaged by cloud probes within convective cumulus during the ACRIDICON–CHUVA campaign (Braga et al., 2017a).

Here, we performed our analysis in the following general steps.

- The relationship between the measured r_e and PWC near the top of convective clouds is calculated based on CCP measurements (described in Sect. 3.1).
- The precipitation probability as a function of the measured r_e and DWC near the top of convective clouds is detailed in Sect. 3.2.
- The vertical development of cloud particle growth near the top of growing convective cumuli is described for clean and polluted conditions in Sect. 3.3.
- The extent of agreement between r_e and PWC measured near cloud tops is discussed in Sects. 4 and 5.

To this end, the following cloud properties were taken into account during our analysis.

Effective radius ($r_e - [\mu\text{m}]$) is calculated as follows:

$$r_e = \frac{\int_{1.5 \mu\text{m}}^{480 \mu\text{m}} r^3 N(r) dr}{\int_{1.5 \mu\text{m}}^{480 \mu\text{m}} r^2 N(r) dr}. \quad (3)$$

Cloud particle effective radius ($r_{ec} - [\mu\text{m}]$) is calculated as follows:

$$r_{ec} = \frac{\int_{1.5 \mu\text{m}}^{25 \mu\text{m}} r^3 N(r) dr}{\int_{1.5 \mu\text{m}}^{25 \mu\text{m}} r^2 N(r) dr}. \quad (4)$$

Mean radius ($r_M - [\mu\text{m}]$) is calculated as follows:

$$r_M = \frac{1}{N} \int_{1.5 \mu\text{m}}^{480 \mu\text{m}} r N(r) dr. \quad (5)$$

Mean volume radius ($r_V - [\mu\text{m}]$) is calculated as follows:

$$r_V = \left(\frac{\int_{1.5 \mu\text{m}}^{480 \mu\text{m}} r^3 N(r) dr}{\int_{1.5 \mu\text{m}}^{480 \mu\text{m}} N(r) dr} \right)^{\frac{1}{3}}. \quad (6)$$

Modal radius ($r_{\text{MOD}} - [\mu\text{m}]$) is the radius in which

$$\frac{\partial N(r)}{\partial r} \Big|_{1.5 \mu\text{m}}^{480 \mu\text{m}} = 0. \quad (7)$$

Cloud mass ratio (CMR) is calculated as follows:

$$\text{CMR} = \frac{\int_{1.5 \mu\text{m}}^{25 \mu\text{m}} r^3 N(r) dr}{\int_{1.5 \mu\text{m}}^{480 \mu\text{m}} r^3 N(r) dr}. \quad (8)$$

Precipitation mass ratio (PMR) is calculated as follows:

$$\text{PMR} = \frac{\int_{25 \mu\text{m}}^{480 \mu\text{m}} r^3 N(r) dr}{\int_{1.5 \mu\text{m}}^{480 \mu\text{m}} r^3 N(r) dr}. \quad (9)$$

The uncertainties of the calculated values of r_e , r_{ec} , r_V , r_{MOD} , CMR, and PMR are $\sim 10\%$ (Braga et al., 2017a, b). The uncertainties of calculated DWC and PWC are $\sim 30\%$. Furthermore, Braga et al. (2017b) showed that water drops were observed near cloud tops for air temperatures (T) warmer than -9°C over the Amazon Basin. For $T \leq -9^\circ\text{C}$, ice initiation was found. Therefore, for cloud particles measured at $T > -9^\circ\text{C}$ the density of water (1 g cm^{-3}) is used in calculations of cloud properties. For $T \leq -9^\circ\text{C}$, Braga et al. (2017b) showed that mostly graupel and frozen drops were imaged by the CIPgs, and thus we assume in our calculations that the density of frozen particles is 0.9 g cm^{-3} (\sim the density of pure ice). Results assuming smaller density for ice particles are also shown in the Supplement. In addition, we assume a spherical shape for water and ice particles in our calculations. The density of ice particles within clouds is associated with the microphysical mechanism of their growth. An ice particle that originated from a frozen drop or ice crystal due to accretion processes to an irregular or roundish particle has bulk density of $0.8 \text{ g cm}^{-3} < \rho < 0.99 \text{ g cm}^{-3}$ (Pruppacher and Klett, 1997). Furthermore, the density of rimed ice particles can be strongly influenced by the cloud drops frozen onto the ice crystal. These factors can result in graupel particles with densities ranging between 0.05 and 0.9 g cm^{-3} . Ice particles formed by deposition of water vapor and collision of snow crystals (e.g., snowflakes, ice crystals, needles, columns, and sheets) typically have low densities ($\sim 0.05 \text{ g cm}^{-3} < \rho < 0.5 \text{ g cm}^{-3}$). Therefore, based on the type of particles imaged by the CIPgs during our measurements we assume that the uncertainty in the calculated r_e and PWC is small in comparison to the measurement uncertainty (i.e., $\sim 10\%$ and $\sim 30\%$ for r_e and PWC, respectively).

3 Results

3.1 Comparison of measured r_e and PWC near the top of convective clouds

Figure 2 shows the measured r_e and PWC near the top of convective clouds. The precipitation was found in liquid




Figure 2. Effective radius of cloud particles (r_e) vs. precipitation water content (PWC) measured near cloud tops of convective clouds over the Atlantic Ocean for temperatures (T) warmer than -9°C (in blue), and over the continent for $T > -9^\circ\text{C}$ (in green) and $T \leq -9^\circ\text{C}$ (in red). The black line indicates the fit of PWC as a function of r_e (shown on the top of the graphic). The r_e threshold of $13\ \mu\text{m}$ applied for the fit is based on the value of r_e at which light precipitation starts (drizzle water content $> 0.01\ \text{g m}^{-3}$). The coefficient of determination (R^2) from the fit function of this analysis is shown on the top.

and solid phases for temperatures ranging between -26 and 10°C (see Fig. S1 in the Supplement for T - r_e profiles). The relationship between r_e and PWC can be well expressed by a linear function ($R^2 \sim 0.89$) for liquid and frozen precipitation. The high correlation between r_e and PWC was not found for r_e when considering only the cloud drop size range ($r < 25\ \mu\text{m}$) (see Fig. S2). When precipitating particles are neglected in the calculation of r_e , the large increase of precipitation mass is not captured, and thus, r_{ec} values do not exceed $17\ \mu\text{m}$ in our analysis. Nevertheless, these characteristics do not prevent r_{ec} from identifying the threshold of precipitation initiation, as shown in previous studies (Freud and Rosenfeld, 2012; Rosenfeld and Gutman, 1994; Braga et al., 2017b). A possible explanation for the similar linear relationships for liquid and frozen precipitation is that the formation of ice particles was initiated mostly by freezing raindrops during flights AC09 and AC18, cases in which warm rain formation was not completely suppressed (Braga et al., 2017b). In addition, the assumption of an ice density of $0.9\ \text{g cm}^{-3}$ for frozen particles while calculating PWC can also lead to deviations in the values of the adjusted equation of r_e - PWC. Nevertheless, similar results were found when assuming frozen particles with lower density ($0.45\ \text{g cm}^{-3}$) when calculating PWC (see Fig. S3).

3.2 Precipitation probability as a function of the measured r_e

Figure 3a shows the precipitation probability as a function of r_e near cloud tops of convective clouds. The probability of precipitation (PP) is the fraction of in-cloud measurements (at 1 Hz) that exceed a given DWC threshold (e.g., for $\text{DWC} > 0.01\ \text{g m}^{-3}$). This was calculated as a function of r_e to identify the threshold of precipitation initiation. The DWC includes only particles with a terminal fall speed of $\sim 1\ \text{m s}^{-1}$ or less, which maximizes the chance that the drizzle was formed in situ and had not fallen a large distance from above (Freud and Rosenfeld, 2012; Braga et al., 2017b). The figure shows that precipitation initiation is expected to occur at $r_e > 13\ \mu\text{m}$, but some very light precipitation can occur already between 10 and $12\ \mu\text{m}$.

For $r_e < 13\ \mu\text{m}$, a few passes with light precipitation were found (see Fig. 3b). For warm temperatures, these measurements were performed over the Atlantic Ocean during flight AC19. Above the ocean, the presence of giant CCN can lead to warm rain initiation for r_e below $13\ \mu\text{m}$ (Freud and Rosenfeld, 2012; Konwar et al., 2012). Precipitating particles were measured for $r_e < 13\ \mu\text{m}$ in cloud passes with cold temperatures, in which graupel particles (probably with low density) were imaged by the CCP (see Fig. S4). This type of particles was imaged during flights AC13 and AC07, in clouds in which the coalescence process was completely suppressed, and thus precipitating particles are formed mainly by accretion. For $r_e > 13\ \mu\text{m}$, PWC increases rapidly as a function of r_e , which is probably associated with an increase of drop collision (or collision kernel) in cumulus clouds. Similar results are found for measurements with cloud water content larger than 25 % of the adiabatic water content (see Fig. S5), in which convectively diluted or dissipating clouds are excluded. This r_e threshold is consistent with the result found by Braga et al. (2017b) for r_{ec} .

3.3 The relationship between r_e and cloud mass

The thermal instability in the boundary layer promotes the formation of convective clouds consisting of regions with updrafts and downdrafts. Tropical convective cumuli typically develop in updrafts, and during their vertical development cloud droplets are converted into precipitation by coalescence or accretion processes. Near cloud tops the amount of water or ice mass (M) within the cloud parcel of convective clouds can be described by

$$M = M_c + M_p, \quad (10)$$

where M_c is the mass of particles with cloud droplet sizes $1.5\ \mu\text{m} < r < 25\ \mu\text{m}$ and M_p is the mass of particles with precipitating sizes ($r \geq 25\ \mu\text{m}$).

Figure 4a shows the particle size distribution (PSD) measured within relatively clean clouds and different resulting




Figure 3. (a) Precipitation probability as a function of r_e for different drizzle water content (DWC) thresholds (black: $\text{DWC} > 0.01 \text{ g m}^{-3}$; blue: $\text{DWC} > 0.02 \text{ g m}^{-3}$; green: $\text{DWC} > 0.03 \text{ g m}^{-3}$; yellow: $\text{DWC} > 0.05 \text{ g m}^{-3}$; red: $\text{DWC} > 0.1 \text{ g m}^{-3}$) measured within convective cloud tops over the Amazon Basin and Atlantic Ocean. The dashed line indicates the number of cases for each r_e size interval (right axis); each case represents a 1 s in-cloud measurement. (b) Effective radius (r_e) as a function of DWC measured within convective cloud tops over the Atlantic Ocean for temperatures (T) warmer than -9°C (in blue), and over the continent for $T > -9^\circ\text{C}$ (in green) and $T \leq -9^\circ\text{C}$ (in red).

r_e during flight AC09, while Fig. 4b and c show measured PSDs in marine clean clouds during flight AC19 and very polluted convective clouds during flight AC13, respectively. The figure shows that for the cleaner cases, droplets grow by coalescence to large drops and form precipitation at warmer temperatures. For the polluted case, the cloud droplets do not coalesce and the precipitation-size particles in ice phase are formed by accretion. More numerous small particles are found in polluted clouds in comparison with the clean clouds. Furthermore, for clean and polluted clouds, the number con-

centration of precipitating particles, and thus M_p , increases as a function of r_e .

Figure 5 shows the cloud mass ratio (CMR) and the precipitation mass ratio (PMR) in convective clouds for cloud passes with $r_e > 13 \mu\text{m}$ (\sim precipitation initiation threshold). This figure shows the precipitation ratio increasing as a function of r_e , while the cloud mass ratio is decreasing at the same time. There is a clear anti-correlation between CMR and PMR. This inverse relationship was found because no precipitation from higher cloud regions disturbed the measurements, and thus, the formation of precipitating particles is associated with the growth of smaller particles at cloud tops.

4 Discussion

The findings shown in this study highlight r_e as a crucial quantity to define the microphysical stage of convective cloud development. We show that precipitation near the cloud tops of convective clouds can be identified and estimated with high accuracy based on in situ measurements of r_e . For $r_e > 13 \mu\text{m}$ the mass of cloud drops and precipitation can be retrieved as well. Furthermore, our analysis shows that neglecting precipitating particles ($r > 25 \mu\text{m}$) in the calculation of r_e , as performed in previous studies (Freud and Rosenfeld, 2012; Braga et al., 2017b), has obscured the tight r_e –PWC relationship shown in this study. A similar tight relationship between the size of hydrometeors and rain rates was found initially by Marshall and Palmer (1948). These authors showed that the rain rate and hence PWC has a strong correlation with the raindrop diameter (D) and concentration. The radar reflectivity of particles depends on D^6 and is commonly used to estimate rain rates. Here, we show for the first time the close relationship between r_e and PWC measured at cloud tops of convective clouds. This close relationship was found due to the inclusion of particles with precipitating sizes up to $\sim 1 \text{ mm}$ in diameter when deriving the r_e –PWC relationship. Our analysis also suggests that similar linear relationships may be found for cloud particles with diameters up to $250 \mu\text{m}$ (see Fig. S6). In addition, we found a larger sensitivity of r_e to precipitating particles in comparison with typical quantities used to characterize size distributions of particles (e.g., mean radius, mean volume radius, and modal radius) (see Fig. S7). This sensitivity of r_e to precipitating particles was more evident for cloud passes with precipitation formed by coalescence processes (see Fig. S8).

The relationships between in situ r_e and cloud properties were found in cloud tops of growing convective cumuli. Nevertheless, similar relations between these measurements can be expected to be found in cloud tops of other types of clouds, in which the same precipitation-forming processes take place (i.e., coagulation and accretion). The applicability of our findings, e.g., for satellite measurements, depends on the sen-




Figure 4. (a) Number size distribution of particles in clean convective clouds measured during flight AC09 for different r_e and temperatures (T). The values of r_e , the number concentration of particles (N_d), and T for each case are shown on the right side of the panels. Panels (b) and (c) are similar to (a) for marine clouds measured during flight AC19 and for polluted clouds measured during flight AC13, respectively. Particles with precipitating sizes (raindrops and ice hydrometeors) are indicated by colors.




Figure 5. Effective radius of cloud particles (r_e) vs. cloud mass ratio (CMR) indicated by blue dots and precipitation mass ratio (PMR) indicated by red dots for cloud passes where $r_e > 13 \mu\text{m}$. The blue line indicates the best fit for measurements of CMR as a function of r_e (the equation is indicated in blue at the top of the graph). The R^2 from this fit function is 0.97. The red line indicates the best fit for measurements of PMR as a function of r_e (the equation is indicated in red at the top of the graphic). The R^2 from the fit function of $r_e - \text{PMR}$ is 0.93. The number of cloud passes in this analysis is 254.

sitivity of the retrieved r_e to precipitating particles at the top of clouds. Previous studies (King et al., 2013; Krisna et al., 2018; Noble and Hudson, 2015; Painemal and Zuidema, 2011) have shown good agreement between in situ measurements of r_e and co-located r_e retrieved from the Moderate Resolution Imaging Spectroradiometer (MODIS) aboard the Aqua and Terra satellites. Furthermore, the retrieved r_e from MODIS has also shown good agreement with the retrieved r_e from the new generation of Chinese geostationary meteorological satellites FY-4A (Chen et al., 2020). The r_e retrieved based on satellite passive infrared remote sensing represents a vertically weighted value, where the cloud top layers are weighted the most. Investigating the relationship between the r_e retrieved by satellites with in situ measurements of PWC is an important task to confirm the relationship identified in the present study and to further establish r_e as a parameter to quantify PWC within clouds.

5 Conclusions

This study investigated the relationship between the effective radius, r_e , of droplets and ice particles and PWC measured near the top of growing convective clouds. Data collected during the dry season over the Amazon Basin and over the western tropical Atlantic with the CCP probe on board the HALO aircraft were used in the analysis. The measurements

were performed on clean and polluted air masses with cloud tops with temperatures between ~ 10 and ~ -26 °C. Our results show that rain starts at the top of most of the convective clouds over the Amazon Basin during the dry season when the measured r_e exceeds about 13 μm . For marine clouds, warm rain started when r_e was between 10 and 12 μm , probably due to the presence of giant CCN in marine air masses. In polluted air masses, in which warm rain was completely suppressed, precipitation starts at smaller r_e (~ 10 μm), and the observed precipitation particles were ice hydrometeors. We show for the first time that there is a clear linear relationship ($R \sim 0.94$) between r_e and PWC at the tops of convective clouds. Our results also highlight that at cloud tops, the mass of cloud and precipitating particles can be estimated based on the value of r_e after rain starts. These remarkable results were found because the aircraft preferably measured growing convective cloud towers near their tops, where no precipitation from higher cloud regions disturbed the in situ precipitation-forming processes. Our findings from cloud top measurements under different thermodynamic and pollution conditions over the Amazon Basin (dry and polluted air masses) and Atlantic Ocean (wet and clean air masses) suggest that the r_e –PWC relationship can be extended for modeling and remote sensing applications. However, further analysis of the r_e –PWC relationship including different types of precipitating clouds (at cloud tops and below), environmental conditions, and others precipitating-forming processes (e.g., aggregation of ice hydrometeors) is needed to assess the limits of the applicability of this study.

Data availability. The data used in this study can be found at <https://halo-db.pa.op.dlr.de/mission/5> (last access: 30 April 2021, Deutsches Zentrum für Luft- und Raumfahrt, 2014).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-21-14079-2021-supplement>.

Author contributions. RCB, DR, OOK, BE, and MLP led the analyses and the manuscript preparation. The measurements of cloud properties were conducted and analyzed by RCB, CV, DR, and MLP. The measurements were led by MOA, MW, and UP. The results were reviewed by DR, BE, BAH, MW, TK, UP, and MOA.

Competing interests. Some authors are members of the editorial board of *Atmospheric Chemistry and Physics*. The peer-review process was guided by an independent editor, and the authors have also no other competing interests to declare.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. We thank the ACRIDICON–CHUVA team. The ACRIDICON–CHUVA campaign was supported by the Max Planck Society (MPG), the German Science Foundation (DFG Priority Program SPP 1294), the German Aerospace Center (DLR), and a wide range of other institutional partners.

Financial support. This research has been supported by the German Science Foundation (DFG Priority Program SPP 1294). Barbara Ervens was financially supported by the French National Research Agency (ANR) (grant no. ANR-17-MPGA-0013).

The article processing charges for this open-access publication were covered by the Max Planck Society.

Review statement. This paper was edited by Farahnaz Khosrawi and reviewed by two anonymous referees.

References

- Albrecht, R. I., Morales, C. A., and Silva Dias, M. A. F.: Electrification of precipitating systems over the Amazon: Physical processes of thunderstorm development, *J. Geophys. Res.*, 116, D08209, <https://doi.org/10.1029/2010JD014756>, 2011.
- Andreae, M. O., Browell, E. V., Garstang, M., Gregory, G. L., Harriss, R. C., Hill, G. F., Jacob, D. J., Pereira, M. C., Sachse, G. W., Setzer, A. W., Dias, P. L. S., Talbot, R. W., Torres, A. L., and Wofsy, S. C.: Biomass-burning emissions and associated haze layers over Amazonia, *J. Geophys. Res.*, 93, 1509–1527, <https://doi.org/10.1029/JD093iD02p01509>, 1988.
- Andreae, M. O., Rosenfeld, D., Artaxo, P., Costa, A. A., Frank, G. P., Longo, K. M., and Silva-Dias, M. A. F.: Smoking rain clouds over the Amazon, *Science*, 303, 1337–1342, <https://doi.org/10.1126/science.1092779>, 2004.
- Artaxo, P., Martins, J. V., Yamasoe, M. A., Procópio, A. S., Pauliquevis, T. M., Andreae, M. O., Guyon, P., Gatti, L. V., and Leal, A. M. C.: Physical and chemical properties of aerosols in the wet and dry seasons in Rondônia, Amazonia, *J. Geophys. Res.*, 107, LBA 49-1–LBA 49-14, <https://doi.org/10.1029/2001JD000666>, 2002.
- Artaxo, P., Rizzo, L. V., Brito, J. F., Barbosa, H. M. J., Arana, A., Sena, E. T., Cirino, G. G., Bastos, W., Martin, S. T., and Andreae, M. O.: Atmospheric aerosols in Amazonia and land use change: from natural biogenic to biomass burning conditions, *Faraday Discuss.*, 165, 203–235, <https://doi.org/10.1039/C3FD00052D>, 2013.
- Beard, K.: Terminal Velocity and Shape of Cloud and Precipitation Drops Aloft, *J. Atmos. Sci.*, 33, 851–864, [https://doi.org/10.1175/1520-0469\(1976\)033<0851:TVASOC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033<0851:TVASOC>2.0.CO;2), 1976.
- Berg, W., L'Ecuyer, T., and van den Heever, S.: Evidence for the impact of aerosols on the onset and microphysical properties of rainfall from a combination of satellite observations and cloud-resolving model simulations, *J. Geophys. Res.*, 113, D14S23, <https://doi.org/10.1029/2007JD009649>, 2008.
- Braga, R. C., Rosenfeld, D., Weigel, R., Jurkat, T., Andreae, M. O., Wendisch, M., Pöhlker, M. L., Klimach, T., Pöschl, U.,

- Pöhlker, C., Voigt, C., Mahnke, C., Borrmann, S., Albrecht, R. I., Mollenker, S., Vila, D. A., Machado, L. A. T., and Artaxo, P.: Comparing parameterized versus measured microphysical properties of tropical convective cloud bases during the ACRIDICON-CHUVA campaign, *Atmos. Chem. Phys.*, 17, 7365–7386, <https://doi.org/10.5194/acp-17-7365-2017>, 2017a.
- Braga, R. C., Rosenfeld, D., Weigel, R., Jurkat, T., Andreae, M. O., Wendisch, M., Pöschl, U., Voigt, C., Mahnke, C., Borrmann, S., Albrecht, R. I., Mollenker, S., Vila, D. A., Machado, L. A. T., and Grulich, L.: Further evidence for CCN aerosol concentrations determining the height of warm rain and ice initiation in convective clouds over the Amazon basin, *Atmos. Chem. Phys.*, 17, 14433–14456, <https://doi.org/10.5194/acp-17-14433-2017>, 2017b.
- Brenguier, J.-L., Bachalo, W. D., Chuang, P.Y., Esposito, B. M., Fugal, J., Garrett, T., Gayet, J.-F., Gerber, H., Heymsfield, A., Kokhanovsky, A., Korolev, A., Lawson, R. P., Rogers, D. C., Shaw, R. A., Strapp, W., and Wendisch, M.: In Situ Measurements of Cloud and Precipitation Particles, in: *Airborne Measurements for Environmental Research*, edited by: Kokhanovsky, A., Wendisch, M., and Brenguier, J.-L., 225–301, <https://doi.org/10.1002/9783527653218.ch5>, 2013.
- Cecchini, M. A., Machado, L. A. T., Andreae, M. O., Martin, S. T., Albrecht, R. I., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Fütterer, D., Jurkat, T., Mahnke, C., Minikin, A., Mollenker, S., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Voigt, C., Weinzierl, A., and Wendisch, M.: Sensitivities of Amazonian clouds to aerosols and updraft speed, *Atmos. Chem. Phys.*, 17, 10037–10050, <https://doi.org/10.5194/acp-17-10037-2017>, 2017.
- Chen, Y., Chen, G., Cui, C., Zhang, A., Wan, R., Zhou, S., Wang, D., and Fu, Y.: Retrieval of the vertical evolution of the cloud effective radius from the Chinese FY-4 (Feng Yun 4) next-generation geostationary satellites, *Atmos. Chem. Phys.*, 20, 1131–1145, <https://doi.org/10.5194/acp-20-1131-2020>, 2020.
- Deutsches Zentrum für Luft- und Raumfahrt: Mission: ACRIDICONCHUVA, HALO [data set], available at: <https://halo-db.pa.op.dlr.de/mission/5> (last access: 30 April 2021), 2014.
- Freud, E. and Rosenfeld, D.: Linear relation between convective cloud drop number concentration and depth for rain initiation, *J. Geophys. Res.*, 117, D02207, <https://doi.org/10.1029/2011JD016457>, 2012.
- Khain, A. P. and Pinsky, M.: *Physical Processes in Clouds and Cloud Modeling*, Cambridge University Press, Cambridge, 222–229, <https://doi.org/10.1017/9781139049481>, ISBN:9781139049481, 2018.
- Khain, A. P., BenMoshe, N., and Pokrovsky, A.: Factors Determining the Impact of Aerosols on Surface Precipitation from Clouds: An Attempt at Classification, *J. Atmos. Sci.*, 65, 1721–1748, <https://doi.org/10.1175/2007JAS2515.1>, 2008.
- King, N. J., Bower, K. N., Crosier, J., and Crawford, I.: Evaluating MODIS cloud retrievals with in situ observations from VOCALS-REx, *Atmos. Chem. Phys.*, 13, 191–209, <https://doi.org/10.5194/acp-13-191-2013>, 2013.
- Konwar, M., Maheskumar, R. S., Kulkarni, J. R., Freud, E., Goswami, B. N., and Rosenfeld, D.: Aerosol control on depth of warm rain in convective clouds, *J. Geophys. Res.*, 117, D13204, <https://doi.org/10.1029/2012JD017585>, 2012.
- Krisna, T. C., Wendisch, M., Ehrlich, A., Jäkel, E., Werner, F., Weigel, R., Borrmann, S., Mahnke, C., Pöschl, U., Andreae, M. O., Voigt, C., and Machado, L. A. T.: Comparing airborne and satellite retrievals of cloud optical thickness and particle effective radius using a spectral radiance ratio technique: two case studies for cirrus and deep convective clouds, *Atmos. Chem. Phys.*, 18, 4439–4462, <https://doi.org/10.5194/acp-18-4439-2018>, 2018.
- Lance, S., Brock, C. A., Rogers, D., and Gordon, J. A.: Water droplet calibration of the Cloud Droplet Probe (CDP) and in-flight performance in liquid, ice and mixed-phase clouds during ARCPAC, *Atmos. Meas. Tech.*, 3, 1683–1706, <https://doi.org/10.5194/amt-3-1683-2010>, 2010.
- Liu, C., Zipser, E. J., and Nesbitt, S. W.: Global Distribution of Tropical Deep Convection: Different Perspectives from TRMM Infrared and Radar Data, *J. Climate*, 20, 489–503, <https://doi.org/10.1175/JCLI4023.1>, 2007.
- Mallaun, C., Giez, A., and Baumann, R.: Calibration of 3-D wind measurements on a single-engine research aircraft, *Atmos. Meas. Tech.*, 8, 3177–3196, <https://doi.org/10.5194/amt-8-3177-2015>, 2015.
- Marshall, J. S. and Palmer, W. M. K.: The Distribution of Raindrops With Size, *J. Atmos. Sci.*, 5, 165–166, [https://doi.org/10.1175/1520-0469\(1948\)005<0165:TDORWS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1948)005<0165:TDORWS>2.0.CO;2), 1948.
- Martin, S. T., Artaxo, P., Machado, L., Manzi, A. O., Souza, R. A. F., Schumacher, C., Wang, J., Biscaro, T., Brito, J., Calheiros, A., Jardine, K., Medeiros, A., Portela, B., de Sá, S. S., Adachi, K., Aiken, A. C., Albrecht, R., Alexander, L., Andreae, M. O., Barbosa, H. M. J., Buseck, P., Chand, D., Comstock, J. M., Day, D. A., Dubey, M., Fan, J., Fast, J., Fisch, G., Fortner, E., Giangrande, S., Gilles, M., Goldstein, A. H., Guenther, A., Hubbe, J., Jensen, M., Jimenez, J. L., Keutsch, F. N., Kim, S., Kuang, C., Laskin, A., McKinney, K., Mei, F., Miller, M., Nascimento, R., Pauliquevis, T., Pekour, M., Peres, J., Petäjä, T., Pöhlker, C., Pöschl, U., Rizzo, L., Schmid, B., Shilling, J. E., Dias, M. A. S., Smith, J. N., Tomlinson, J. M., Tóta, J., and Wendisch, M.: The Green Ocean Amazon Experiment (GoAmazon2014/5) Observes Pollution Affecting Gases, Aerosols, Clouds, and Rainfall over the Rain Forest, *B. Am. Meteorol. Soc.*, 98, 981–997, <https://doi.org/10.1175/BAMS-D-15-00221.1>, 2016.
- Noble, S. R. and Hudson, J. G.: MODIS comparisons with northeastern Pacific in situ stratocumulus microphysics, *J. Geophys. Res.-Atmos.*, 120, 8332–8344, <https://doi.org/10.1002/2014JD022785>, 2015.
- Painemal, D. and Zuidema, P.: Assessment of MODIS cloud effective radius and optical thickness retrievals over the Southeast Pacific with VOCALS-REx in situ measurements, *J. Geophys. Res.*, 116, D24206, <https://doi.org/10.1029/2011JD016155>, 2011.
- Pinsky, M., Mazin, M., Khain, A., and Korolev, A.: Analytical estimation of droplet concentration at cloud base, *J. Geophys. Res.*, 117, D18211, <https://doi.org/10.1029/2012JD017753>, 2012.
- Pöhlker, M. L., Pöhlker, C., Ditas, F., Klimach, T., Hrabec de Angelis, I., Araújo, A., Brito, J., Carbone, S., Cheng, Y., Chi, X., Ditz, R., Gunthe, S. S., Kesselmeier, J., Könemann, T., Lavrič, J. V., Martin, S. T., Mikhailov, E., Moran-Zuloaga, D., Rose, D., Satorno, J., Su, H., Thalman, R., Walter, D., Wang, J., Wolff, S., Barbosa, H. M. J., Artaxo, P., Andreae, M. O., and Pöschl, U.: Long-term observations of cloud condensation nuclei in the Amazon rain forest – Part 1: Aerosol size distribution, hygroscopicity, and new model parametrizations for CCN prediction, *Atmos. Chem. Phys.*, 16, 15709–15740, <https://doi.org/10.5194/acp-16-15709-2016>, 2016.

- Pöhlker, M. L., Ditas, F., Saturno, J., Klimach, T., Hrabě de Angelis, I., Araújo, A. C., Brito, J., Carbone, S., Cheng, Y., Chi, X., Ditz, R., Gunthe, S. S., Holanda, B. A., Kandler, K., Kesselmeier, J., Könemann, T., Krüger, O. O., Lavrič, J. V., Martin, S. T., Mikhailov, E., Moran-Zuloaga, D., Rizzo, L. V., Rose, D., Su, H., Thalman, R., Walter, D., Wang, J., Wolff, S., Barbosa, H. M. J., Artaxo, P., Andreae, M. O., Pöschl, U., and Pöhlker, C.: Long-term observations of cloud condensation nuclei over the Amazon rain forest – Part 2: Variability and characteristics of biomass burning, long-range transport, and pristine rain forest aerosols, *Atmos. Chem. Phys.*, 18, 10289–10331, <https://doi.org/10.5194/acp-18-10289-2018>, 2018.
- Pruppacher, H. and Klett, J. D.: *Microphysics of Clouds and Precipitation*, Kluwer Academic Publishers, 2nd edn., 38–58, ISBN:0-7923-4211-9, 1997.
- Reutter, P., Su, H., Trentmann, J., Simmel, M., Rose, D., Gunthe, S. S., Wernli, H., Andreae, M. O., and Pöschl, U.: Aerosol- and updraft-limited regimes of cloud droplet formation: influence of particle number, size and hygroscopicity on the activation of cloud condensation nuclei (CCN), *Atmos. Chem. Phys.*, 9, 7067–7080, <https://doi.org/10.5194/acp-9-7067-2009>, 2009.
- Roberts, G. C., Andreae, M. O., Zhou, J., and Artaxo, P.: Cloud condensation nuclei in the Amazon Basin: “marine” conditions over a continent?, *Geophys. Res. Lett.*, 28, 2807–2810, <https://doi.org/10.1029/2000GL012585>, 2001.
- Roberts, G. C., Nenes, A., Seinfeld, J. H., and Andreae, M. O.: Impact of biomass burning on cloud properties in the Amazon Basin, *J. Geophys. Res.*, 108, 4062, <https://doi.org/10.1029/2001JD000985>, 2003.
- Roca, R., Aublanc, J., Chambon, P., Fiolleau, T., and Viltard, N.: Robust Observational Quantification of the Contribution of Mesoscale Convective Systems to Rainfall in the Tropics, *J. Climate*, 27, 4952–4958, <https://doi.org/10.1175/JCLI-D-13-00628.1>, 2014.
- Rosenfeld, D. and Gutman, G.: Retrieving microphysical properties near the tops of potential rain clouds by multispectral analysis of AVHRR data, *Atmos. Res.*, 34, 259–283, [https://doi.org/10.1016/0169-8095\(94\)90096-5](https://doi.org/10.1016/0169-8095(94)90096-5), 1994.
- Rosenfeld, D., Lohmann, U., Raga, G. B., O’Dowd, C. D., Kulmala, M., Fuzzi, S., Reissell, A., and Andreae, M. O.: Flood or Drought: How Do Aerosols Affect Precipitation?, *Science*, 321, 1309–1313, <https://doi.org/10.1126/science.1160606>, 2008.
- Sorooshian, A., Anderson, B., Bauer, S. E., Braun, R. A., Cairns, B., Crosbie, E., Dadashazar, H., Diskin, G., Ferrare, R., Flagan, R. C., Hair, J., Hostetler, C., Jonsson, H. H., Kleb, M. M., Liu, H., MacDonald, A. B., McComiskey, A., Moore, R., Painemal, D., Russell, L. M., Seinfeld, J. H., Shook, M., Smith, W. L., Thornhill, K., Tselioudis, G., Wang, H., Zeng, X., Zhang, B., Ziemba, L., and Zuidema, P.: Aerosol–Cloud–Meteorology Interaction Airborne Field Investigations: Using Lessons Learned from the U.S. West Coast in the Design of ACTIVATE off the U.S. East Coast, *B. Am. Meteorol. Soc.*, 100, 1511–1528, <https://doi.org/10.1175/BAMS-D-18-0100.1>, 2019.
- Tao, W. K., Chen, J. P., Li, Z., Wang, C., and Zhang, C.: Impact of aerosols on convective clouds and precipitation, *Rev. Geophys.*, 50, RG2001, <https://doi.org/10.1029/2011RG000369>, 2012.
- Twomey, S.: Pollution and the planetary albedo, *Atmospheric Environment* (1967), 8, 1251–1256, [https://doi.org/10.1016/0004-6981\(74\)90004-3](https://doi.org/10.1016/0004-6981(74)90004-3), 1974.
- Weigel, R., Spichtinger, P., Mahnke, C., Klingebiel, M., Afchine, A., Petzold, A., Krämer, M., Costa, A., Molleker, S., Reutter, P., Szakáll, M., Port, M., Grulich, L., Jurkat, T., Minikin, A., and Borrmann, S.: Thermodynamic correction of particle concentrations measured by underwing probes on fast-flying aircraft, *Atmos. Meas. Tech.*, 9, 5135–5162, <https://doi.org/10.5194/amt-9-5135-2016>, 2016.
- Wendisch, M., Poschl, U., Andreae, M. O., MacHado, L. A. T., Albrecht, R., Schlager, H., Rosenfeld, D., Martin, S. T., Abdelmonem, A., Afchine, A., Araújo, A. C., Artaxo, P., Aufmhoff, H., Barbosa, H. M. J., Borrmann, S., Braga, R., Buchholz, B., Cecchini, M. A., Costa, A., Curtius, J., Dollner, M., Dorf, M., Dreiling, V., Ebert, V., Ehrlich, A., Ewald, F., Fisch, G., Fix, A., Frank, F., Futterer, D., Heckl, C., Heidelberg, F., Huneke, T., Jakel, E., Jarvinen, E., Jurkat, T., Kanter, S., Kastner, U., Kentner, M., Kesselmeier, J., Klimach, T., Knecht, M., Kohl, R., Kolling, T., Kramer, M., Kruger, M., Krisna, T. C., Lavric, J. V., Longo, K., Mahnke, C., Manzi, A. O., Mayer, B., Mertes, S., Minikin, A., Molleker, S., Munch, S., Nillius, B., Pfeilsticker, K., Pöhlker, C., Roiger, A., Rose, D., Rosenow, D., Sauer, D., Schnaiter, M., Schneider, J., Schulz, C., De Souza, R. A. F., Spanu, A., Stock, P., Vila, D., Voigt, C., Walsler, A., Walter, D., Weigel, R., Weinzierl, B., Werner, F., Yamasoe, M. A., Ziereis, H., Zinner, T., and Zoger, M.: Acridicon-chuva campaign: Studying tropical deep convective clouds and precipitation over amazonia using the New German research aircraft HALO, *B. Am. Meteorol. Soc.*, 97, 1885–1908, <https://doi.org/10.1175/BAMS-D-14-00255.1>, 2016.
- Williams, E., Rosenfeld, D., Madden, N., Gerlach, J., Gears, N., Atkinson, L., Dunnemann, N., Frostrom, G., Antonio, M., Biazon, B., Camargo, R., Franca, H., Gomes, A., Lima, M., Machado, R., Manhaes, S., Nachtigall, L., Piva, H., Quintiliano, W., Machado, L., Artaxo, P., Roberts, G., Renno, N., Blakeslee, R., Bailey, J., Boccippio, D., Betts, A., Wolff, D., Roy, B., Halverson, J., Rickenbach, T., Fuentes, J., and Avelino, E.: Contrasting convective regimes over the Amazon: Implications for cloud electrification, *J. Geophys. Res.*, 107, LBA 50-1–LBA 50-19, <https://doi.org/10.1029/2001JD000380>, 2002.
- Zipser, E. J., Cecil, D. J., Liu, C., Nesbitt, S. W., and Yorty, D. P.: Where Are the Most Intense Thunderstorms on Earth?, *B. Am. Meteorol. Soc.*, 87, 1057–1072, <https://doi.org/10.1175/BAMS-87-8-1057>, 2006.