



HAL
open science

Sulphur dioxide emission rates from Yasur volcano, Vanuatu archipelago

Philipson Bani, M. Lardy

► **To cite this version:**

Philipson Bani, M. Lardy. Sulphur dioxide emission rates from Yasur volcano, Vanuatu archipelago. Geophysical Research Letters, 2007, 34 (20), pp.L20309. 10.1029/2007GL030411 . hal-03881731

HAL Id: hal-03881731

<https://uca.hal.science/hal-03881731>

Submitted on 2 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Sulphur dioxide emission rates from Yasur volcano, Vanuatu archipelago

P. Bani^{1,2} and M. Lardy^{1,2}

Received 18 April 2007; revised 21 August 2007; accepted 19 September 2007; published 23 October 2007.

[1] Yasur volcano has long been suspected as one of the volcanic emission sources responsible for a significant contribution of sulphur dioxide to the atmosphere. But due to its isolation in the south Pacific region, very little is known about its emission rates. We report here sulphur dioxide flux measurements, obtained on Yasur, using ultraviolet spectroscopy, from April 2004 to November 2005. We found an average flux of 7.9 kg.s^{-1} of SO_2 , which constitutes some 1–2% and 1.5–2.5% of estimated global time-integrated volcanic emissions to the atmosphere and troposphere, respectively. **Citation:** Bani, P., and M. Lardy (2007), Sulphur dioxide emission rates from Yasur volcano, Vanuatu archipelago, *Geophys. Res. Lett.*, *34*, L20309, doi:10.1029/2007GL030411.

1. Introduction

[2] SO_2 emission rates are important indicators of volcanic activity and are routinely measured at a number of volcanoes in support of hazard assessment. These emissions can also have important impacts on the atmosphere and climate at various temporal and spatial scales. For instance, volcanic sulphur emissions have been implicated in tropospheric cooling, stratospheric heating, stratospheric ozone depletion, and alteration in stratospheric global circulation patterns, due to radiative and chemical interactions [Robock and Oppenheimer, 2003]. Volcanic SO_2 clouds are measured from above by spaceborne instruments, and from below by traversing, or scanning the plume from fixed positions, with correlation spectrometers (COSPECs) [Stoiber *et al.*, 1983], or more recently with miniature ultraviolet spectrometers, by differential optical absorption spectroscopy (DOAS) [Galle *et al.*, 2003; McGonigle *et al.*, 2002]. The product of these flux estimates and mass ratios of other plume species to SO_2 (i.e., X/SO_2) yields to the fluxes of other components (e.g., CO_2 , HCl, HF).

[3] In the compilations of global volcanic SO_2 output [Andres and Kasgnoc, 1998], Yasur volcano in Vanuatu archipelago appears in the top ten of a list of forty-nine continuous volcanic SO_2 emitters. However, until this work, the flux estimates for Yasur were based on comparisons of plume size with other volcanoes [Nairn, 1988; Lardy and Marty, 1990]. The only field SO_2 flux measurements were performed in July 1999 with a correlation spectrometer but constrained to weather condition [Gauthier, 1999]. Thus

Yasur gas discharge is arguably one of the least well constrained of all those reported in global flux budgets. We present here a series of SO_2 flux measurements results, obtained between April 2004 and November 2005 on Yasur volcano using a DOAS instrument.

2. Geologic Setting

[4] The island Tanna lies approximately 150 km east of New Hebrides trench and 150 km above the Benioff zone [Louat *et al.*, 1998]. It covers 550 km^2 and is built up by Pliocene to recent subaerial and subaqueous basaltic and andesitic volcanic units and reef limestone. The volcanic units belong to three major groups (Figure 1): the upper Pliocene to Pleistocene Green Hill group to the north and east, the Pleistocene Tukosmeru group to the south, and the upper Pleistocene to Recent Siwi group in the easternmost part within a 24 km^2 caldera [Carney and Macfarlane, 1979; Coulon and Maury, 1981; Robin *et al.*, 1994]. The Green Hill Group and Tukosmeru group are predominantly basaltic or basaltic andesite in composition while Siwi group are basaltic andesite to andesite in composition. An uplift of more than 150 m has occurred in the eastern portion of Siwi group forming the Yenkahe Horst [Chen *et al.*, 1995]. Yasur, a small scoria cone volcano (361 m above sea level) which occupies the western part of the Yenkahe Horts, is characterised by a sustained degassing with a continuous strombolian to vulcanian type activity.

3. Methodology

[5] Fluxes were obtained by traversing on a 4WD vehicle underneath the plume with a USB2000 miniature ultraviolet spectrometer which spanned the spectral interval 280–400 nm with a spectral resolution of 0.5 nm. Exposure time for individual spectra was 200 ms, and we co-added 8 spectra to enhance the signal-to-noise ratio. This device was coupled to a vertically pointing telescope of 8 mrad field of view with an optic fibre bundle. A USB cable connected the spectrometer to a laptop computer, providing power and data transfer. Software control of the USB2000 was achieved using Jscript executed in DOASIS software [Kraus, 2001] to save and analyse spectra, providing real time concentration readings. The position of each UV spectrum was determined from a continuously recording GPS unit. These spectrometers have also been used for measurements of volcanic BrO [Bobrowski *et al.*, 2003, 2007; Oppenheimer *et al.*, 2006], H_2S [O'Dwyer *et al.*, 2003] and NO_2 [Oppenheimer *et al.*, 2005]. Details of the DOAS routine used in concentration evaluation are given by Platt [1994] and/or Galle *et al.* [2003]. Fluxes were obtained from the retrieved SO_2 columns and the GPS track

¹Laboratoire des Géosciences et Physique de la Matière Condensée, Université de la Nouvelle Calédonie, Noumea, New Caledonia.

²Institut de Recherche pour le Développement, Noumea, New Caledonia.




Figure 1. (left) Map of Vanuatu, showing Tanna Island and the New Hebrides Trench. (right) The geology of Tanna, redrawn after *Carney and Macfarlane* [1979]. (a) Map of eastern Tanna, showing the Yenkahe horst, the Yasur crater, the ash plain where most of the traverses were obtained and the positions of spectrometers in the plume speed measurements. Note that on the 10th January 2005 the plume dispersion was to the south due to the presence of cyclone Kerry located 1800 km southwest of Vanuatu at that time.

by multiplying the distance traversed perpendicular to the plume transport direction corresponding to each spectrum by that spectrum's column amount, summing these products across the whole plume and multiplying the total by the estimated plume speed. For Yasur the traverses were performed on the ash plain (Figure 1) and plume speeds were obtained either from a hand held anemometer on top of Yasur, or from simultaneously recording overhead SO₂ time series from underneath the plume, using 2 and 3 spectrometers (Table 1). These devices are situated between 0.5 and 1.0 km from the source and between 80–150 m from each other (Figure 1). Peaks and troughs obtained from the series, corresponding to overhead transit of plume structure, have been cross-correlated, using methods described by

McGonigle et al. [2005] and *Williams-Jones et al.* [2006], to find the plume speed.

4. Results and Discussion

[6] From April 2004 to November 2005 a total of 86 traverses were accomplished under the plume less than 2 km

Table 1. SO₂ Fluxes From Yasur Volcano Measured During 2004–2005

Measurement Period	Date, day/month/year	Number of Traverses	Measurement Period Mean Flux, kg.s ⁻¹	Method Used for Plume Speed
P1	02–04/04/04	26	5.3 ± 1.0	Anemo.
P2	11–14/07/04	33	12.3 ± 2.2	Anemo.
P3	19/09/04	5	4.9 ± 0.9	Anemo.
P4	29/10/04	6	5.3 ± 0.9	Anemo.
P5	10/01/05	4	11.2 ± 2.1	2 spectro
P6	18/03/05	1	2.5 ± 0.5	Anemo.
P7	02/07/05	5	6.1 ± 1.1	2 spectro.
P8	01–02/11/05	6	13.5 ± 2.5	3 spectro.




Figure 2. Representative plot of a cross sectional profile of SO₂ concentration for one traverse obtained under the Yasur plume in July 2005. The three peaks correspond to individual plumes emitted from the three active vents in the crater.




Figure 3. Correlation between seismic counts and SO₂ fluxes; volcanic events with seismograph displacements greater than 12.5 μm were recorded and plotted here from the period between January 2004 and December 2005. The SO₂ flux for each measurement period (P) is represented by circles.

downwind of Yasur volcano. The average fluxes from these measurements for each measurement period (P1 to P8) are shown in Table 1. Figure 2 shows a representative plot of concentration versus distance-traversed perpendicular to the plume transport direction. The profiles of the column amounts recorded on Yasur present frequently three maxima, and indicate the dispersion and overlap of individual plumes emitted from the three active vents within the crater. The overall average flux obtained for the traverses is

7.9 kg.s⁻¹, with individual fluxes ranging from 2.5 to 17.2 kg.s⁻¹. Apparent SO₂ fluxes from volcanoes can, and do, vary considerably, even on short timescales, so no direct agreement between these estimates made at different times is expected. However, the general flux fluctuation from P1 through P8 tend to be consistent with the volcanic behaviour of Yasur recorded by a permanent seismic monitoring station located 2 km from the volcano (Figure 3) where seismograph displacements greater than 12.5 μm




Figure 4. Correlation plot between seismic events and SO₂ emission rates. Emissions rate which correspond to P5 and P7 were omitted because of high gas discharge which occurred with no significant seismic events and lack of seismic data respectively.

gives a representative idea of the volcano's daily activity [Tabbagh *et al.*, 2002]. An exception in this general correlation appears in P5 where high gas discharge occurred with no sustained explosions which probably due to higher pressure in the magmatic feeding system as evidenced by the high level of magmatic column. The lowest gas discharge, measured in March 2005 (216 t.d⁻¹), and the maximum discharge rate, measured in July 2004 (1063 t.d⁻¹) and November 2005 (1166 t.d⁻¹), are correlated to the seismic activity of Yasur (Figure 4). This parallel variation of the SO₂ fluxes and the seismic activity records implies that the SO₂ emission rates from Yasur do vary with the volcanic activity. Similar correlations between volcanic seismicity and SO₂ have been observed on Etna [Leonardi *et al.*, 2000] or Soufriere Hill [Watson *et al.*, 2000]. Thus during low activity phases Yasur SO₂ emission rates can be reduced to around 200 t.d⁻¹, while during high activity with virulent explosions, Yasur releases more than a thousand tons per day. High activity phases on Yasur usually last more than a month (Institut de Recherche pour le Développement, unpublished data, 1993–2005), and thus release regularly an equivalent of more than 30 kt of SO₂ into the atmosphere during one simple high activity phase.

[7] Our results confirm that Yasur is one of the largest point sources of SO₂ on Earth and constitutes 1–2% of the global time-integrated volcanic emissions to the whole atmosphere (425–665 kg.s⁻¹), and 1.5–2.5% of the total volcanogenic emission to the troposphere (308–540 kg.s⁻¹) [e.g., Andres and Kasgnoc, 1998; Halmer *et al.*, 2002].

5. Conclusions

[8] We have reported on SO₂ flux measurements made with a miniature ultraviolet spectrometer for Yasur volcano in Vanuatu archipelago. We obtained fluxes between 2.5 and 17.2 kg.s⁻¹ with a mean flux value of 7.9 kg.s⁻¹: equivalent to 1–2% of the time-integrated total global volcanic SO₂ emission. Such volcanic emissions merit greater attention in future, in order to provide better spatial and temporal constraints on global volcanic sulphur loading into the atmosphere.

[9] **Acknowledgments.** We gratefully acknowledge the support of the “Institut de Recherche pour le Développement” (New Caledonia), Cambridge volcanology group (UK), the Department of Geology Mines and Waters Resources (Vanuatu) and the “Laboratoire de Géosciences et de Physique de la Matière Condensée” (University of New Caledonia).

References

- Andres, R. J., and A. D. Kasgnoc (1998), A time-average inventory of subaerial volcanic sulphur emissions, *J. Geophys. Res.*, *103*, 25,251–25,261.
- Bobrowski, N., G. Hönninger, B. Galle, and U. Platt (2003), Detection of bromine monoxide in a volcanic plume, *Nature*, *423*, 273–276, doi:10.1038/nature01625.
- Bobrowski, N., R. von Glasow, A. Aiuppa, S. Inguaggiato, I. Louban, O. W. Ibrahim, and U. Platt (2007), Reactive halogen chemistry in volcanic plumes, *J. Geophys. Res.*, *112*, D06311, doi:10.1029/2006JD007206.
- Carney, J. N., and A. Macfarlane (1979), Geology of Tanna, Aneityum, Futuna and Aniwa, report, pp. 5–29, New Hebrides Gov. Geol. Surv., Port Vila, Vanuatu.
- Chen, J. K., F. W. Talor, R. L. Edwards, H. Cheng, and G. S. Burr (1995), Recent emerged reef terraces of the Yenkahe resurgent block, Tanna, Vanuatu: Implications for volcanic, landslide and tsunami hazards, *J. Geol.*, *103*, 577–590.

- Coulon, C., and R. C. Maury (1981), Petrology of tholeiitic lavas from cesse in island arc magmatism, *Bull. Volcanol. Geotherm. Res.*, *13*, 147–171.
- Galle, B., C. Oppenheimer, A. Geyer, A. J. S. McGonigle, M. Edmonds, and L. A. Horrock (2003), A miniaturised UV spectrometer for remote sensing of SO₂ fluxes: A new tool for volcano surveillance, *J. Volcanol. Geotherm. Res.*, *119*, 241–254, doi:10.1016/S0377-0273(02)00356-6.
- Gauthier, P. J. (1999), Géochimie des fluides et des gaz émis par le système volcano-hydrothermal du volcan Yasur, circ:le de Tanna, République du Vanuatu, report, 56 pp., Los Alamos Natl. Lab., Los Alamos, N. M.
- Halmer, M. M., H.-U. Schmincke, and H.-F. Graf (2002), The annual volcanic gas input into the atmosphere, in particular into the stratosphere: A global data set for the last 100 years, *J. Volcanol. Geotherm. Res.*, *115*, 511–528, doi:10.1016/S0377-0273(01)00318-3.
- Kraus, S. (2001), The DOASIS software, paper presented at the 1st International Differential Optical Absorption Spectroscopy Workshop, AC-CENT/AT2, Heidelberg, Germany, 13–14 Sept.
- Lardy, M., and B. Marty (1990), Incandescent tephra ejection; small lava lakes; strong SO₂ emission, *Bull. Global Volcanism Network*, *15*, 11.
- Leonardi, S., S. Gresta, and F. Mulargia (2000), Cross-correlation between volcanic tremor and SO₂ flux data from Mount Etna volcano, 1987–1992, *Phys. Chem. Earth Part A*, *25*(9–11), 737–740.
- Louat, R., M. Hamburger, and M. Monzier (1998), Shallow and intermediate depth seismicity in the New Hebrides Arc: Constraints on subduction process, in *Geology and Offshore Resources of Pacific Island Arcs-Vanuatu Region*, *Earth Sci. Ser.*, vol. 8, edited by H. G. Green and F. L. Wong, pp. 329–356, Circum-Pac. Coun. for Energy and Min. Resour., Menlo Park, Calif.
- McGonigle, A. J. S., C. Oppenheimer, B. Galle, T. A. Mather, and D. M. Pyle (2002), Walking traverse and scanning DOAS measurements of volcanic gas emission rates, *Geophys. Res. Lett.*, *29*(20), 1985, doi:10.1029/2002GL015827.
- McGonigle, A. J. S., D. R. Hilton, T. P. Fischer, and C. Oppenheimer (2005), Plume velocity determination for volcanic SO₂ flux measurements, *Geophys. Res. Lett.*, *32*, L11302, doi:10.1029/2005GL022470.
- Nairn, I. (1988), Centuries-long Strombolian eruption continues from four vents. Yasur volcano, *Sci. Event Alert Network*, *13*, 12.
- O'Dwyer, M., M. J. Padgett, A. J. S. McGonigle, C. Oppenheimer, and S. Inguaggiato (2003), Real-time measurement of volcanic H₂S and SO₂ concentrations by UV spectroscopy, *Geophys. Res. Lett.*, *30*(12), 1652, doi:10.1029/2003GL017246.
- Oppenheimer, C., P. R. Kyle, V. I. Tsanev, A. J. S. McGonigle, T. A. Mather, and D. Sweeney (2005), Mt Erebus, the largest point source of NO₂ in Antarctica, *Atmos. Environ.*, *39*, 6000–6006, doi:10.1016/j.atmosenv.2005.06.036.
- Oppenheimer, C., I. V. Tsanev, C. F. Braban, R. A. Cox, J. W. Adams, A. Aiuppa, N. Bobrowski, P. Delmelle, J. Barclay, and J. S. McGonigle (2006), BrO formation in volcanic plumes, *Geochim. Cosmochim. Acta*, *70*, 2935–2941.
- Platt, U. (1994), Differential Optical Absorption Spectroscopy (DOAS), in *Air Monitoring by Spectroscopic Techniques*, *Chem. Anal. Ser.*, vol. 127, edited by M. W. Sigrist, pp. 27–84, John Wiley, New York.
- Robin, C., J.-P. Essein, and M. Monzier (1994), Ignimbrites of basaltic andesite and andesite compositions from Tanna, New Hebrides Arc, *Bull. Volcanol.*, *56*, 170–183.
- Robock, A., and C. Oppenheimer (Eds.) (2003), *Volcanism and Earth's Atmosphere*, *Geophys. Monogr. Ser.*, vol. 139, 360 pp., AGU, Washington, D.C.
- Stoiber, R. F., L. L. Malinconico Jr., and S. N. Williams (1983), Use of the correlation spectrometer at volcanoes, in *Forecasting Volcanic Events*, edited by H. Tazieff and J. C. Sabroux, pp. 425–444, Elsevier Sci., New York.
- Tabbagh, J., M. Lardy, S. Wallez, and D. Charley (2002), Mild eruption phase ends and leads to vigorous phase in December 2001; seismic data, *Bull. Global Volcanism Network*, *27*, 01.
- Watson, I. M., et al. (2000), The relation between degassing and ground deformation at Soufriere Hills Volcano, Montserrat, *J. Volcanol. Geotherm. Res.*, *98*, 117–126.
- Williams-Jones, G., K. Horton, H. Garbeil, P. J. Mougins-Mark, A. J. L. Harris, A. J. Sutton, and T. Elias (2006), Accurately measuring volcanic plume velocities with UV spectrometers, *Bull. Volcanol.*, *68*, 323–327.

P. Bani, Laboratoire des Géosciences et physique de la matière condensée, Université de la Nouvelle Calédonie, BP 4477, F-98848 Nouméa, Nouvelle Calédonie. (philpson.bani@noumea.ird.nc)

M. Lardy, Institut de Recherche pour le Développement, BP A5, F-98848 Nouméa Cedex, Nouvelle Calédonie.