



HAL
open science

Exploring alternative salting methods to reduce sodium content in blueveined cheeses

Imène Ferroukhi, Cécile Bord, René Lavigne, Christophe Chassard, Julie Mardon

► To cite this version:

Imène Ferroukhi, Cécile Bord, René Lavigne, Christophe Chassard, Julie Mardon. Exploring alternative salting methods to reduce sodium content in blueveined cheeses. *International Dairy Journal*, 2023, 138, 10.1016/j.idairyj.2022.105555 . hal-03963923v2

HAL Id: hal-03963923

<https://uca.hal.science/hal-03963923v2>

Submitted on 15 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



Contents lists available at ScienceDirect

International Dairy Journal

journal homepage: www.elsevier.com/locate/idairyj

Exploring alternative salting methods to reduce sodium content in blue-veined cheeses



Imène Ferroukhi, Cécile Bord, René Lavigne, Christophe Chassard, Julie Mardon*

Université Clermont Auvergne, INRAE, VetAgro Sup, UMR 0545 Fromage, 63370 Lempdes, France

ARTICLE INFO

Article history:

Received 13 August 2022

Received in revised form

14 November 2022

Accepted 18 November 2022

Available online 30 November 2022

ABSTRACT

Reducing sodium content in food is a major public health challenge, especially for cheeses. Different salt reduction methods in surface dry-salted blue cheese and their effect on biochemical, physicochemical, microbiological, rheological and sensory characteristics of cheese were investigated. The cheeses were salted with coarse or fine salt using various methods. All methods decreased sodium content in cheese core (−11% to −67%) and induced higher water activity. Sprinkling (−45% NaCl content) and calcium lactate substitution (−10% NaCl content) were technically feasible. The sanitary quality and cheese microbiota were not impacted except for a reduction of yeasts and moulds. Higher indices of richness and diversity were found on salt-reduced cheeses with *Streptococcus*, *Lactococcus* and *Leuconostoc mesenteroides* dominating. Texture and appearance of the cheeses were also affected. The research identified the limits of salt reduction in blue-veined cheeses; sprinkling and partial substitution with calcium lactate should be further studied.

© 2022 Elsevier Ltd. All rights reserved.

1. Introduction

The sodium content of processed foods is considered a public health concern. There is strong evidence for a causal relationship between salt intake and blood pressure, leading to cardiovascular disease (Doyle & Glass, 2010). The World Health Organisation has suggested reducing sodium intake by 30% to obtain the WHO guideline of 2 g day^{−1} (i.e., 5 g of salt day^{−1}) by 2025 (WHO, 2021). It was estimated that 75% of sodium intake comes from processed foods (Brown, Tzoulaki, Candeias, & Elliott, 2009).

In France, bread, meat products, soups and cheese are the main contributors to sodium chloride (NaCl) intake in the population (ANSES, 2012). Blue-veined cheeses appear as the saltiest variety of cheeses with salt content up to 4% (ANSES, 2012). Despite its nutritional value (Ferroukhi et al., 2022), one serving of blue cheese may contribute up to 40% of the recommended dietary Na intake (Guinee & O'Kennedy, 2007). Thus, reducing NaCl content of blue cheese is necessary to help improve health outcomes. However, the reduction of salt is difficult to achieve because of its major functions in cheese: it modulates the physicochemical and biochemical properties of the cheese during ripening, maintains a low a_w , which

controls the development of micro-organisms, and is involved in the sensory characteristics of finished products (Delahunty & Drake, 2004; Guinee, 2004).

Many varieties of blue cheese are salted with an excess of salt by repeated surface application of dry NaCl while others are brine-salted (Guinee, 2004). Surface dry-salting process is complex and poorly studied. Dry salt is dissolved on the surface and diffuses slowly into the matrix, creating a counterflow of moisture from the centre to the outside (Guinee, 2004). Salt diffusion depends on the intrinsic properties of the cheese, the quantity of salt added, the degree of salt rehydration on the surface and the salt granularity (Guinee & Fox, 2017). Consequently, studying salt reduction strategies in such salted products implies investigation of these parameters.

One of the salt reduction strategies is based on the simple reduction of the level of added NaCl (or direct decrease of NaCl) as already performed in soft and semi-hard cheeses, Feta and Cheddar cheese (Aly, 1995; Dugat-Bony et al., 2019; Møller, Rattray, Høier, & Ardö, 2012). Dugat-Bony et al. (2019) reported that salt reduction in soft cheese induced more sensory changes than in semi-hard cheese. In contrast, salt reduction increased the growth of spoilage agent in semi-hard cheese. Nevertheless, there is still a lack of information on reducing the salt content of other cheeses such as dry-salted cheeses. The direct reduction of salt quantities has so far been applied in cheese technologies where salting is done

* Corresponding author.

E-mail address: julie.mardon@vetagro-sup.fr (J. Mardon).

in coagulum and the common quantity of salt used is known. This is different from dry surface salting of blue-veined cheeses where an uncontrolled quantity of dry salt is rubbed and retained on the cheese surface.

The second strategy focuses on the total or partial replacement of NaCl by other salts. In Cheddar, mozzarella and kefalograviera cheeses it has been shown that potassium chloride (KCl) can be used successfully to achieve a large reduction in sodium without adverse effect on the quality of cheeses (Chavhan, Kanawjia, Khetra, & Puri, 2015; Grummer, Bobowski, Karalus, Vickers, & Schoenfuss, 2013; Grummer, Karalus, Zhang, Vickers, & Schoenfuss, 2012; Katsiari, Voutsinas, Alichanidis, & Roussis, 1998). In contrast, other authors have reported that substitution of salt by KCl increases the bitterness of Cheddar cheese and disrupt the biochemical reactions of the cheese such as lipolysis and proteolysis (Fitzgerald & Buckley, 1985; Lindsay, Hargett, & Bush, 1982; Rulikowska et al., 2013). However, because of hypercalcaemia risks, KCl-substituted products should be consumed with caution by consumers with renal disease (Berthet, 2009). The substitution of NaCl by MgCl₂ or CaCl₂ has been reported imparting off flavours (sour, bitter, metallic, and soapy) and decreases Cheddar cheese acceptability (Fitzgerald & Buckley, 1985; Grummer et al., 2012). In addition, from a nutritional standpoint MgCl₂ and CaCl₂ are acid-forming compounds (Remer, 2001). Little research has been done on the salt substitution in blue-veined cheeses. One recent study used organic calcium salts to reduce the acid-forming potential of blue-veined cheeses (Gore, Mardon, Cécile, & Lebecque, 2019). This research revealed that a partial NaCl replacement by calcium lactate induced a reduction of Na content by 19% in Fourme d'Ambert. Except for the Gore et al. (2019) study, calcium lactate has been used so far in calcium fortification of milk (39%), cottage cheese and yoghurt (36%) (Reykdal & Lee, 1993; Singh & Muthukumarappan, 2008; Yonis, Elzamzamy, & Elmorsi, 2013). Therefore, this substitute could be of nutritional interest.

In this context, it is mandatory to further explore salt reduction strategies in surface dry-salted products to provide relevant alternative process and allow blue cheeses to better fulfil nutritional guidelines. Thus, the objective of this study was to investigate the relevance of alternative salting methods to reduce the salt level in a surface dry salted blue-veined cheese. Various salting processes and salt granularity were tested through two kinds of reduction strategies: simple reduction of NaCl quantities and partial substitution of NaCl by calcium lactate. The different effects of these salt reductions on physicochemical, biochemical, microbiological, rheological and sensory parameters of finished products were monitored.

2. Materials and methods

2.1. Cheese making

Cheesemaking of a blue-veined cheese 'Bleu d'Auvergne' was carried out according to the PDO specifications and under industrial conditions (Supplementary material Fig. S1) (INAO, 2021). Raw cow milk (about 1000 L) was pasteurised (72 °C, 30 s) and standardised for fat content and calcium chloride. Milk then was matured (36 °C, 45 min) and added with a classical starter culture (*Lactococcus lactis* subsp. *cremoris*, *L. lactis* subsp. *lactis*, *Streptococcus thermophilus*) (Danisco, Denmark), heterofermentative bacteria (*Leuconostoc mesenteroides* subsp. *mesenteroides*) (Lallemand, France) and *Penicillium roqueforti* (SAS LIP, France). A quantity (30 mL 100 L⁻¹) of liquid rennet extract (Caglificio Clerici, Italy) was also introduced shortly thereafter at 34.6 °C. After that, the coagulum was cut vertically and horizontally and then stirred to form 'the cap' of the coagulum grains (3 cm³). The syneresis was carried out in moulds to obtain drained curds. After the moulding, the cheese curds were

dry-salted according to the various salting treatments tested. Six days after the manufacture, the cheeses were pricked. This step aims to create air channels in the core of the cheese to enable the *P. roqueforti* to grow and colonise it. The ripening was carried out at 8 °C and 95% relative humidity (RH) over a period of 28 days.

2.2. Salting treatments and sampling

Surface dry-salting of Bleu d'Auvergne are described in Table 1. To obtain anhydrous salts, all the salts were previously dried (102 °C, 24 h). Two control treatments (Cc and Cf) varying in the salt granularity [coarse salt (0.9–3.15 mm) or fine salt (0.22 mm)] were applied according to the traditional method by rubbing the cheese surface with excess of salt. Then, five methods of net reduction of NaCl quantities were tested by changing salt granularity and application procedure (STc, SRc, STf, SRf and SP). STc and STf consisted of rolling cheese on a layer of coarse or fine salt spread on a tray. SRc and SRf were to remove the excess of salt from the surface by rubbing. The SP method was to sprinkle 100 g of fine salt over the whole surface of the cheese. One method of partial substitution of NaCl with calcium lactate (S30) was carried out. The substitution (S30) was carried out using fine salt replaced with 30% of calcium lactate (0.162 mm) (Merck KGaA, Germany), to have a homogeneous grain size mixture. The mixture of NaCl-calcium lactate was homogenised in a mixer and placed in sealed containers until applications.

Eight blocks of cheese (2 ± 0.5 kg) were sampled at 28 days of ripening. The rind was cut to 5 mm using a cutting wire and the

Table 1
Salting methods.

Code name	Approach	Salt	Method description
Coarse salt–salt reduction			
Cg	Control	NaCl	Salting with standard method of rubbing the cheese surface with excess of coarse salt.
STc	Salting on a tray	NaCl	A layer of coarse salt was spread on a horizontal tray. The cheese was salted by rolling the around and both sides in coarse salt without rubbing or pressing.
SRc	Surface rubbing	NaCl	After salting in an excess of coarse salt, the surface of the cheese was scrubbed to remove the excess of salt.
Fine salt–salt reduction			
Cf	Control	NaCl	Salting with standard method of rubbing the cheese surface with excess of fine salt.
STf	Salting on a tray	NaCl	A layer of fine salt was spread on a horizontal tray. The cheese was salted by rolling the around and both sides in fine salt without rubbing or pressing.
SRf	Surface rubbing	NaCl	After salting in an excess of fine salt, the surface of the cheese was scrubbed to remove the excess of salt.
SP	Sprinkling	NaCl	A fixed quantity of fine salt (100 g) was sprinkled all over the cheese surface.
Fine salt–salt substitution			
S30	Partial NaCl substitution	70% NaCl + 30% calcium lactate	Salting with the standard method in an excess of salt mixture.

core was cut into pieces and ground in a blender (Moulinex, France). To limit the variability due to the heterogeneity of blue cheese matrix, the core samples were homogenised by mechanical agitation and frozen at $-20\text{ }^{\circ}\text{C}$ in sealed bags until analysed. For each salting modality, one ripened cheese was held in cold storage at $1\text{ }^{\circ}\text{C}$ prior to sensory analysis. A commercial cheese (Std) was included to compare the produced cheeses with a reference, particularly for sensory evaluation. This cheese was only studied for physicochemical and sensory analysis.

2.3. Physicochemical and biochemical analyses

The physicochemical composition of the cheeses was analysed in compliance with ISO standards. Cheese pH was measured using a penetration pH electrode CG 840 (Schott, Mainz, Germany). Dry matter (DM) was analysed by desiccation. In compliance with the standard NF ISO 5534:2004, a weighed test portion mixed with sand was dried by heating it for 24 h at $102\text{ }^{\circ}\text{C}$. The dried test portion was then weighed to determine the mass loss. Fat content was determined using the HEISS method according to the procedure described in standard NF V 04-287:2019. In a cheese butyrometer, the proteins were first dissolved with a combination of acetic acid and perchloric acid and then separated from the fat by centrifugation. The fat content was read directly on the butyrometric scale with correction equation. Water activity (a_w) was measured in $\approx 5\text{ g}$ grated cheese samples at $20\text{ }^{\circ}\text{C}$, as per the manufacturer's instructions (HYGROLAB C1, Rotronic, Bassersdorf, Schweiz). Ratio of fat in dry matter (FDM) was calculated. Total nitrogen matter (TNM) was determined by the Kjeldahl method in compliance with the standard NF ISO 8968-1:2014. Total nitrogen content was converted to crude protein by a factor of 6.38. Calcium (Ca) and sodium (Na) contents were measured by inductively coupled plasma optical emission spectroscopy according to NF EN 16943:2017. The samples were first digested with nitric acid and hydrochloric acid. After nebulisation, the aerosol was directed to a high frequency induced argon plasma, in which the elements were atomised and excited for irradiation. Sodium content was converted into salt percent. All measurements were performed in triplicate.

2.4. Microbiological analyses

Cheese samples (25 g) were homogenised in a stomacher for 2 min with 250 mL peptone water (BagMixer CC, France). Serial dilutions (10^{-7}) and the sowing were made with spiral method (EasySpiral Dilute, interscience, France). Total microflora was counted according to NF EN ISO 4833-2:2013. *Leuconostocs* were enumerated on Mayeux, Sandine and Elliker agar (MSE), *Lactococcus* on M17 agar and *Lactobacillus* on MRS agar according to NF EN 15787. Yeasts and moulds were counted with NF V 08-059. *Enterobacteriaceae* were also screened on VRBG agar. Cell counts on media were performed in triplicate.

To investigate microbial communities in blue cheese, a meta-genomic analysis was performed on cheeses after 28 days of ripening. DNA was extracted from the cheeses (*Penicillium*-free zone) using the FastDNA® SPIN Kit for Soil (MP Biomedicals, Illkirch, France). DNA was quantified using Qubit Fluorometric Quantitation (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA) method. Amplification and sequencing were carried out in the V3–V4 region of the 16S rDNA gene. Amplicons were sequenced with Illumina technology in $2 \times 250\text{ bp}$. Amplicon data from high-throughput sequencing was analysed using the rANOMALY pipeline (Theil & Rifa, 2021), which relies on the dada2 R package to produce amplicon sequence variants (ASVs) as taxonomic units. A decontamination step was carried out based on prevalence of

contaminant ASVs, as identified in the blank samples, and on DNA concentration, as described by the decontam package included in the rANOMALY pipeline. Taxonomic assignment of bacterial sequences was based on two databases, i.e. DAIRYdb v2.0 and SILVA 138, keeping the assignment with the highest confidence or the deepest taxonomic rank (Meola et al., 2019; Quast et al., 2013). The filtered ASVs count table was used to perform statistical analyses.

2.5. Sensory analyses

A panel composed of 8 assessors from the sensory analysis laboratory of higher education and research institute VetAgro Sup was selected according to standard ISO 8586-1:2012 guidelines. These panellists had previous experience in the evaluation of dairy products. To identify the differences between cheeses and to classify the product groups in a sensory space, a flash profile was conducted. Flash profiling is a fast descriptive method to study the degree of similarity between samples. It is based on the selection of a combination of free choice terms and a comparative ranking evaluation (Dairou & Sieffermann, 2002; Delarue & Lawlor, 2014) and provides a relative sensory positioning (mapping) of the samples.

The evaluation was conducted in two sessions. A slice of cheese (50 g) was presented on plates coded with random 3-digit numbers. All cheeses were served at $20 \pm 1\text{ }^{\circ}\text{C}$, presented monadically and distributed according to a Williams Latin Square designed to take into account the first order effect and carry-over based on the simultaneous presentation of the whole samples set. During the first session (2.5 h), subjects selected their own terms to describe and evaluate 9 blue-cheeses simultaneously (8 salted with various processes presented in Table 1 and 1 commercial blue cheese as a standard). Between each sample, panellists were asked to rinse their mouth successively with unsalted crackers and tepid water to remove fatty residual. At the end of this session, a discussion with panellists allowed finalisation of the list of descriptors generated by each panellist. During the second session, all samples were presented again to the assessors who were asked to rank each product on a ranking scale (rank 1 = least intense, rank 9 = most intense). For the final session, assessors evaluated the cheeses again and ranked them, using the generated list of descriptors. All evaluation sessions were led in computerised booth according to ISO standard 8589:2010. Data was collected using Tastel software® (version 2011; ABT Informatique, Rouvroy-sur-Marne, France).

2.6. Texture measurement

The texture parameters hardness (N), adhesiveness (N s), cohesiveness and gumminess were determined using a rheometer (Kinexus pro+, rSpace for Kinexus 1.61 Software, Malvern Instruments, Malvern, UK), by carrying out the texture profile analysis (TPA) method described by (Nishinari, Kohyama, Kumagai, Funami, & Bourne, 2013). Three cylindrical samples (2 cm of diameter) were randomly collected from each cheese. Each sample was held at $20\text{ }^{\circ}\text{C}$ for 1 h before analysis and was maintained at $20\text{ }^{\circ}\text{C}$ during measurement with the Peltier heating element. Samples were compressed to 20% of their original height, using two compression cycles at a constant crosshead velocity of 50 mm s^{-2} . The texture parameters were calculated according to Henneberry, Wilkinson, Kilcawley, Kelly, and Guinee (2015).

2.7. Statistical analyses

Statistical analysis of all data was performed using XLSTAT software 2020 (Addinsoft, Paris, France). Results are reported as a means \pm standard deviation. Shapiro–Wilk test ($P < 0.05$) was used

to check the normality of the data. A non-parametric test (Kruskal–Wallis) and a post-hoc comparison (Conover–Iman procedure) were applied to all data (Except for sensory data) as it did not have a normal distribution. Differences between mean values were considered significant at $P < 0.05$. Regarding sensory data, generalised procrustes analysis (GPA) was used for the configuration of the consensus between the judges' sensory maps. GPA calculates a consensus from data matrices of an experiment in sensory profiling and allows comparing the proximity between the attributes that are generated by panellists. For the flash profile, one data matrix corresponded to each judge.

3. Results and discussion

As salt intake is a major public health problem, it is essential to implement actions to reduce salt in foods (Bansal & Mishra, 2020). Blue-veined cheeses represent an important source of salt due to their specific salting methods usually performed in an excess of salt. For the first time, this work investigates the relevance of different surface dry salting methods to reduce salt content in blue-veined cheeses. In other words, this research was done to screen a high number of salting methods to identify the most relevant alternatives that will be considered in blue-veined cheese technology.

3.1. Effect of reducing salt content on physicochemical and biochemical composition

The physicochemical and biochemical composition of blue-veined cheese cores salted by different methods is presented in Table 2. Applied salt reduction methods have impacted all parameters of the ripened cheeses ($P < 0.05$), except for mean fat and total nitrogen content of 28 and 19 g 100 g⁻¹, respectively (data not shown).

The pH obtained for all the cheeses in this study was comparable with the pH found in Bleu d'Auvergne (Bord, Guerinon, & Lebecque, 2016; Duval et al., 2016; Ferroukhi et al., 2022). pH values did not greatly change in the coarse salted cheeses. In contrast, Cf and S30 fine salted cheeses had a higher pH than others with pH of 6.08 and 6.28, respectively. Among the tested salting processes, a significant difference was observed in the a_w values. Cf cheese showed the lowest a_w (0.954) compared with the salt-reduced and the coarse-salted cheeses. A high a_w was also observed in salt-reduced Cheddar cheese (McCarthy, Wilkinson, Kelly, & Guinee, 2015; Rulikowska et al., 2013). The dry matter of the studied cheeses did not vary with salt reduction or granularity. However, it was observed that SRc cheese had a low dry matter content (51.89%),

related to high a_w . The commercial cheese (Std) showed a higher dry matter than other cheeses and a low a_w , probably related to a high salt content.

Regarding salt content and salt/moisture, there was no significant difference between Cc (1.70% NaCl) and Cf (1.68% NaCl) control cheeses salted with two different salt grain sizes. The opposite observations have been made in Fourme d'Ambert cheese, where cheeses salted with fine salt had a higher salt content than cheeses salted with coarse salt (Gore et al., 2019). This reflects that the change in salt granularity did not induce a change in sodium content in the cheese core, i.e., coarse and fine salt penetrated the cheese in the same way. Nevertheless, there was more variability in cheeses salted with coarse salt than in cheeses salted with fine salt. This suggests the difficulty of controlling dry surface salting with coarse salt. The commercial cheese (Std) contained a high salt content (3.5% in salt and 6.4% in salt/moisture) compared with other cheeses. This elevated salt value is probably due to a double dry-salting process which is a common practice in Bleu d'Auvergne cheese making, explaining the difference reported in salt content. To assess the impact of new applied salting methods and limit the variability, cheeses from this research were salted only once.

In coarse salted cheeses, tray salting (STc) and rubbing surface salting (SRc) processes induced a high reduction of sodium contents (28 and 67%, respectively) compared with control (Cc). Rubbing off the excess salt on the surface produced a significant reduction in salt, particularly with coarse salt SRc, indicating that coarse salt is greatly removed if it is rubbed off. However, it has been observed that rubbing the wet surface of the cheese may deteriorate the rind and heterogeneously remove a significant salt quantity. For cheeses salted with fine salt, STf and SRf cheeses were also reduced by 21 and 17% salt compared with control cheese Cf. The tray salting method (STc and STf), which aimed to control deposition of salt on each side of cheese, gave a close reduction of salt of both granularities. The disadvantage of this technique (STc or STf) was the time required to salt the full cheese surface. The objective was to find solutions for the cheese industry, this method could control and reduce the amount of salt but may be difficult for a large-scale production. The sprinkling treatment also led to a significant salt reduction (–45% of salt) compared with control (Cf).

Calcium lactate substitution reduced sodium content by 10% ($P < 0.0001$). A significant decrease (–19%) in sodium has been reported by Gore et al. (2019) in Fourme d'Ambert cheese upon the substitution of salt with 75% calcium lactate. For the same substitution ratio of salt, other authors obtained a higher reduction using KCl in Coalho and Akkawi cheeses (Costa et al., 2018; Kamleh et al., 2015). In dry-salted Feta cheese, the substitution of NaCl by KCl (3:1 and 1:1) did not affect the salt content (Katsiari, Voutsinas,

Table 2
Gross composition of cheese cores salted by different methods.^a

Cheese	Ca (mg 100 g ⁻¹)	Na (mg 100 g ⁻¹)	NaCl (%)	Salt/moisture (%)	pH	Dry matter (%)	a_w
Cc	410.5 ± 0.1 ^b	683.7 ± 22.2 ^f	1.7 ± 0.1 ^f	3.3 ± 0.1 ^f	5.7 ± 0.0 ^{cd}	52.1 ± 0.0 ^a	0.969 ± 0.001 ^c
STc	439.9 ± 0.7 ^c	491.3 ± 11.1 ^c	1.2 ± 0.0 ^c	2.4 ± 0.0 ^c	5.5 ± 0.1 ^b	52.0 ± 0.2 ^a	0.967 ± 0.002 ^c
SRc	381.8 ± 1.0 ^a	223.5 ± 16.9 ^a	0.6 ± 0.0 ^a	1.1 ± 0.1 ^a	5.7 ± 0.0 ^c	51.9 ± 0.6 ^a	0.981 ± 0.001 ^d
Cf	477.7 ± 3.7 ^d	675.0 ± 12.4 ^f	1.7 ± 0.0 ^f	3.2 ± 0.1 ^f	6.1 ± 0.0 ^g	53.4 ± 1.3 ^a	0.954 ± 0.001 ^b
STf	517.5 ± 1.5 ^e	535.9 ± 5.5 ^d	1.3 ± 0.0 ^d	2.5 ± 0.1 ^{cd}	5.4 ± 0.0 ^a	53.1 ± 0.6 ^a	0.966 ± 0.003 ^c
SRf	484.5 ± 2.3 ^d	559.9 ± 5.3 ^d	1.4 ± 0.0 ^d	2.7 ± 0.0 ^d	5.8 ± 0.0 ^{de}	52.5 ± 0.0 ^a	0.968 ± 0.003 ^c
SP	467.1 ± 2.5 ^{cd}	369.5 ± 1.6 ^b	0.9 ± 0.0 ^b	1.8 ± 0.0 ^b	5.9 ± 0.0 ^f	52.4 ± 0.1 ^a	0.964 ± 0.006 ^c
S30	606.6 ± 3.5 ^f	609.5 ± 25.8 ^e	1.5 ± 0.1 ^e	2.9 ± 0.1 ^e	6.3 ± 0.0 ^h	52.7 ± 0.3 ^a	0.980 ± 0.001 ^d
Std	523.5 ± 28.5 ^e	1415.0 ± 15.0 ^g	3.5 ± 0.0 ^g	6.4 ± 0.0 ^g	5.8 ± 0.0 ^{ef}	55.1 ± 0.6 ^b	0.867 ± 0.007 ^a
P-value	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	0.0001	<0.0001

^a Abbreviations are: Cc and Cf, controls with coarse and fine salt, respectively; STc and STf, salting on a tray with coarse and fine salt, respectively; SRc and SRf, surface rubbing with coarse and fine salt, respectively; SP, sprinkling with fine salt; S30, 30% of fine salt replaced with calcium lactate; Std, commercial cheese salted with coarse salt by industrial operator (see Table 1 for salting details). Values are the mean ± standard deviation; means in a row sharing common superscript letters are similar as tested by Conover–Iman test ($P < 0.05$). NaCl was estimated by sodium measurement.

Alichanidis, & Roussis, 1997). We hypothesised that the low salt reduction resulting from calcium lactate substitution in this study is probably due to the heterogeneity of the cheese. The second salting process, as commonly used in the salting of blue cheese, could probably increase the salt reduction level. Nevertheless, only one study previously tested salt replacement in dry salting on the surface of blue-veined cheeses (Gore et al., 2019).

Salt reductions influenced Ca content of cheese cores ($P < 0.0001$). Cheeses salted with fine salt had a relatively higher amount of calcium than cheeses salted with coarse salt. Blue-veined cheeses are a category with a complex matrix resulting in the variability of composition parameters such as salt (Ferroukhi et al., 2022; Gkatzionis, Yunita, Linforth, Dickinson, & Dodd, 2014). Difference in Ca content might be related to the heterogeneity of the matrix that causes variable mineral migration during ripening.

The calcium lactate-substituted cheese had the highest calcium content (606.6 mg kg^{-1}) and this was 21 and 13% higher than the control Cf and the standard Std cheese. Considering the Ca level brought by S30 treatment, a 40 g serving of cheese would attain 25.5% of recommended daily calcium intake (ANSES, 2021). This salt has already been studied to fortify Cottage cheese and yoghurt with calcium (Reykdal & Lee, 1993; Wongkhalaung & Boonyaratanakornkit, 2000; Yonis et al., 2013). This may highlight the nutritional interest of this salt in cheese making.

The level of calcium in the cheese depends on the Ca content of the milk but also on mineral exchange after salting. It was reported that the salt on the surface enters inside the cheese, causing a counterflow of whey which exists towards the surface loaded with minerals such as calcium (Le Graet & Brulé, 1988; Le Graet, Lepienne, Brule, & Ducruet, 1983). However, salt diffusion into the cheese matrix (and conversely Na and Cl migration) is limited by some cheese characteristics, such as free water, viscosity of the aqueous phase, tortuosity and porosity (Guinee, 2004; Guinee & Fox, 1983). This might be the reason for the difference in calcium content in obtained cheeses.

3.2. Effect of reducing salt content on the microbial communities

Table 3 shows the counts of total microorganisms, lactic acid bacteria (LAB), yeasts and moulds for cheeses salted with different salting methods. The microbial composition of cheeses was significantly impacted by the salting processes ($P < 0.05$) (Table 3).

Total counts were not different in the salt-reduced cheeses. A comparable effect was observed in a dry salted cheese (São João cheese) reduced in salt content (Soares, Fernando, Mendes, & Martins, 2015). As opposed to other studies of salt reduction in Cheddar (Rulikowska et al., 2013; Schroeder, Bodyfelt, Wyatt, & McDaniel, 1988), there were no marked effects of salt treatment among all experimental Bleu d'Auvergne cheeses in the total bacterial count. Aerobic mesophilic bacteria should mainly correspond

to lactic acid bacteria (LAB) added after milk pasteurisation, as expected levels of total aerobic mesophilic bacteria were found in experimental and control cheeses. Nevertheless, the *Lactococcus* counts were lower in the Cf and STf cheeses. This might be due to test or sample variability during the analysis. In this study, *Enterobacteriaceae* were not found (below the required count) suggesting that the salt reductions implemented had no impact on the sanitary aspect of the cheeses. A similar result has been reported in salt-reduced cheese (Soares et al., 2015).

Yeasts and moulds (YM) were also affected by salting methods. In the coarse-salted cheeses, the salt-reduced cheeses contained a lower YM counts than control cheeses which had $9.00 \text{ log cfu g}^{-1}$. Cheeses salted with fine salt showed a lower YM counts than coarse salted cheeses except for S30 which had the highest level of YM ($9.26 \text{ log cfu g}^{-1}$). A large proportion of cheeses contained a lower YM count than found in Bleu d'Auvergne cheese (Duval et al., 2016). Moreover, opposite observations were reported on Halloumi and Cheddar cheeses substituted with up to 50% KCl, which had no impact on yeast and mould counts (Kamleh et al., 2012; Reddy & Marth, 1995). The development of the distinctive flavours of blue cheese is dominated by metabolism of moulds during ripening. Salt in the moisture and opening of the matrix by *Leuconostocs*, which produces CO_2 through the citrate metabolism, lead to the growth of *P. roqueforti* in the cheese (Marth & Steele, 2001). The low salt levels obtained in this study could be responsible for the low growth of yeasts and moulds.

A total of 354620 16S rRNA (V3–V4 regions) sequencing reads were generated from the 8 cheese samples, covering different salting methods. The reads were assigned to 36 ASVs for which taxonomic assignment was possible down to species level (or group of species sharing almost identical sequences) in most cases. These ASVs belonged to 21 different bacterial species (Fig. 1) and four different bacterial phyla (*Firmicutes*, *Actinobacteriota*, *Proteobacteria* and *Bacteroidetes*). In all cheeses analysed, *Streptococcus*, *Lactococcus* and *Leuconostoc mesenteroides* species were dominant with a raw abundance sum of 123,966, 127,126 and 27,923 reads, respectively. In a previous study, a dominance of these starter culture species was observed in Bleu d'Auvergne cheese (Ferroukhi et al., 2022) and other blue-cheeses (Caron et al., 2021; Flórez & Mayo, 2006). This bacterial dominance could explain why the bacterial abundance of non-starter LAB strains was extremely low, as these bacteria occur at higher levels only in later stages of ripening (Blaya, Barzideh, & LaPointe, 2018).

Species of *Brevibacterium* and *Brachybacterium* genus were also present with a relative abundance compared with the others of 0.1 and 3%, respectively (Supplementary material Table S1). *Enterobacteriaceae* species were detected in all samples with a relative abundance of less than 0.5%. In line with the microbial count, the low percentage of *Enterobacteriaceae* indicates that the different salting methods did not negatively impact the sanitary quality of the products. *Lactococcus* and *Leuconostoc* species showed a higher

Table 3
Microbial populations in cheese cores at 28 days of ripening.^a

Microorganism	Cc	STc	SRc	Cf	STf	SRf	SP	S30	P-value
<i>Lactococcus</i>	9.00 ± 0.09 ^{cd}	8.84 ± 0.02 ^{bc}	9.23 ± 0.02 ^d	6.74 ± 0.07 ^a	6.97 ± 0.04 ^{ab}	8.97 ± 0.04 ^{cd}	8.96 ± 0.06 ^{cd}	8.49 ± 0.06 ^{ab}	0.00
<i>Lactobacillus</i>	8.84 ± 0.02 ^{bc}	8.93 ± 0.03 ^{cd}	9.04 ± 0.05 ^d	8.75 ± 0.04 ^{ab}	8.86 ± 0.05 ^c	8.85 ± 0.02 ^{bc}	8.65 ± 0.05 ^a	8.66 ± 0.11 ^a	0.00
<i>Leuconostoc</i>	8.35 ± 0.13 ^{ab}	8.25 ± 0.04 ^{ab}	8.30 ± 0.08 ^b	7.98 ± 0.16 ^{ab}	7.97 ± 0.14 ^{ab}	7.98 ± 0.07 ^a	8.07 ± 0.17 ^{ab}	8.09 ± 0.10 ^{ab}	0.03
Yeasts and moulds	9.00 ± 0.05 ^{de}	7.16 ± 0.03 ^{ab}	7.23 ± 0.05 ^{bc}	7.27 ± 0.03 ^{cd}	7.24 ± 0.04 ^{bc}	7.28 ± 0.01 ^{cde}	6.83 ± 0.08 ^a	9.26 ± 0.02 ^e	0.00
Total microbiota	8.58 ± 0.12 ^a	8.79 ± 0.05 ^{abc}	8.75 ± 0.04 ^{ab}	8.95 ± 0.05 ^{bcd}	9.04 ± 0.05 ^d	9.01 ± 0.08 ^{cd}	8.95 ± 0.08 ^{bcd}	8.90 ± 0.12 ^{abcd}	0.01

^a Abbreviations are: Cc and Cf, controls with coarse and fine salt, respectively; STc and STf, salting on a tray with coarse and fine salt, respectively; SRc and SRf, surface rubbing with coarse and fine salt, respectively; SP, sprinkling with fine salt; S30, 30% of fine salt replaced with calcium lactate (see Table 1 for salting details). Values (log cfu g^{-1}) are the mean ± standard deviation; means in a row sharing common superscript letters are similar as tested by Conover–Iman test ($P < 0.05$). *Enterobacteriaceae* were not detected.




Fig. 1. Microbial composition of produced cheeses: A, total bacterial composition in each cheese; B, bacterial composition of the subdominant population (without *Streptococcus* species, *Lactococcus* species and *Leuconostoc mesenteroides*) in each cheese. The vertical bars indicate the percentage of relative abundance of the corresponding bacterial sequences at species level.

percentage in the salt-reduced cheeses, especially in the SP and S30 cheeses. This is supported by the results obtained with microbial enumeration. In contrast, *Streptococcus* was lower in the salt-reduced cheeses. Another *Leuconostoc* species was detected only in SP cheese with 0.011% relative abundance.

In [Supplementary material Table S2](#), community richness and diversity were estimated for each cheese using four alpha diversity parameters (Observed, Chao1, Shannon and Simpson). The values of richness indicators were higher in the salt-reduced cheeses. SP and S30 cheeses had Observed and Chao1 indices of 27 and 28, respectively, for both indicators ($P < 0.05$). The same observations were made on bacterial diversity with Shannon and Simpson's

index ([Supplementary material Table S2](#)). The control cheeses had the same richness and diversity values. Species dissimilarity between cheeses (Beta-diversity) was assessed using the Bray–Curtis method. The results showed that species present in the cheeses were similar ($P > 0.05$) and the salt reduction did not affect the type of species present in the cheese.

3.3. Effect of reducing salt content on rheological and sensory parameters

[Table 4](#) shows the means for the TPA parameters studied in Bleu d'Auvergne cheese at 28 days of ripening. Hardness, cohesiveness

Table 4
Texture profile analysis parameters for cheeses at 28 days of ripening.^a

Cheese	Hardness (N)	Cohesiveness	Gumminess (N)	Adhesiveness (N s)
Cc	1.41 ± 0.07 ^a	0.59 ± 0.15 ^a	0.82 ± 0.17 ^a	-0.07 ± 0.00 ^{bc}
STc	1.13 ± 0.06 ^a	0.47 ± 0.00 ^a	0.53 ± 0.03 ^a	-0.54 ± 0.00 ^a
SRc	0.67 ± 0.42 ^a	0.76 ± 0.17 ^a	0.47 ± 0.21 ^a	-0.472 ± 0.01 ^a
Cf	1.49 ± 0.27 ^a	0.66 ± 0.14 ^a	1.00 ± 0.39 ^a	-0.01 ± 0.00 ^c
STf	0.78 ± 0.24 ^a	0.45 ± 0.02 ^a	0.35 ± 0.09 ^a	-0.50 ± 0.01 ^a
SRf	1.44 ± 0.40 ^a	0.43 ± 0.12 ^a	0.59 ± 0.00 ^a	-0.25 ± 0.01 ^{abc}
SP	1.03 ± 0.00 ^a	0.58 ± 0.03 ^a	0.59 ± 0.04 ^a	-0.22 ± 0.01 ^{abc}
S30	1.38 ± 0.40 ^a	0.42 ± 0.23 ^a	0.57 ± 0.30 ^a	-0.37 ± 0.24 ^{ab}
P-value	0.05	0.22	0.17	0.00

^a Abbreviations are: Cc and Cf, controls with coarse and fine salt, respectively; STc and STf, salting on a tray with coarse and fine salt, respectively; SRc and SRf, surface rubbing with coarse and fine salt, respectively; SP, sprinkling with fine salt; S30, 30% of fine salt replaced with calcium lactate (see Table 1 for salting details). Values are the mean ± standard deviation; means in a row sharing common superscript letters are similar as tested by Conover–Iman test ($P < 0.05$).

and gumminess parameters were not significantly affected by the salting methods. This finding is consistent with other salt reduction investigations in Cheddar and feta cheeses (Fitzgerald & Buckley, 1985; Katsiari et al., 1997). However, SRc and STf cheeses tended to have a lower hardness value. The variability in the matrix of blue-veined cheeses could account for the non-significant effect on the hardness of cheeses. Adhesiveness was significantly lower in control cheeses Cc and Cf than in salt-reduced cheeses. STc, SRc and STf cheeses had the highest adhesiveness (-0.539, -0.472 and -0.501, respectively) SP and S30 had a lower adhesiveness value than the other salt-reduced cheeses. Similarly, an increase in adhesiveness due to salt reduction has been reported by Henneberry et al. (2015) in mozzarella cheese. These authors explained that the increase in adhesiveness during ripening is related to the increased hydration of caseins which should reduce the cohesive (attractive) forces with the calcium phosphate para-casein network.

The sensory analysis was carried out using a flash profile to assess the sensory differences between different salt reductions applied to blue cheeses compared with control cheeses and a standard commercial cheese. The panellists used different terms to characterise the cheeses according to appearance, texture, aroma, and taste. The number of attributes varied between 8 and 12, with an average of 10 attributes for each participant and a total of 74 terms of differentiation (Supplementary material Fig. S2). Most of the sensory attributes obtained in this study were also generated in other blue-veined cheeses (Gkatzionis et al., 2013) and Camembert cheese (Galli, Baptista, Cavalheiro, & Gigante, 2019) analysed by flash profile. The consensus index (Rc) was 0.639 (63.9%) with a high residual percentage (>100%). Std cheese had the lowest residual percentage (83.4%), this could suggest that the commercial cheese had a complex descriptors from other cheeses and a less homogeneous consensus among the panel for this product. This was different from the consensus and higher total variance reported by (Gkatzionis et al., 2013). The low consensus among the testers demonstrated in the test may be due to the difficulty in ranking of the product caused by the high number of cheeses and the variability of their matrix.

The generalised procrustes analysis (GPA) of the Flash profile data provided the relative positioning of the samples based on generated attributes. The plots defined by the first two factors of the GPA analysis (Fig. 2) explained a medium percentage of the total variance (50.48%; 29.10% and 21.38% for F1 and F2 respectively). The distribution of products in the Factorial Map was heterogeneous (Fig. 2A). Generally, good discrimination between cheeses was observed. The first axis discriminated the coarse control cheese Cc and commercial cheese Std with sprinkled cheese SP on one side

and the fine salt control cheese Cf, the substituted cheese S30 and rubbed cheese SRc with other reduced salt cheeses (STc, SRf and STf) on the other side.

Looking at the attribute map (Fig. 2B), among the 74 descriptors generated, the attributes of creamy, marbling and odour intensity were found by many judges and were grouped on the biplot. Bitterness, salty and firmness also appeared in some panellists.

The results showed that texture was impacted by salt reductions. The control cheese Cc and the standard commercial cheese Std were characterised by a firm texture whereas the salt reduced cheeses STc, STf and SRf were characterised by a creamy texture. This result was in agreement with the hardness and adhesiveness index previously found in the texture of these cheeses. A number of studies have demonstrated the impact of salt on the structure and texture of cheeses (Pastorino, Hansen, & McMahon, 2003; Schroeder et al., 1988). Regarding the marbling and odour intensity descriptors, it seemed that this character was more present in SP and Std cheeses. This indicated that marbling quantity was impacted by salt reduction (in agreement with microbiological results), but also related to the heterogeneity of cheese slices distributed to panellists. A within-product variability, due to blue-veined heterogeneity distribution, was already observed during the sensory analysis of this type of cheese (Bord et al., 2016).

Bitterness was found in the S30 and both the Cc and Cf cheeses controls, but this was not representative of many judges. Bitterness in cheese was often related to salt levels. Mistry and Kasperon (1998) showed a decrease in bitterness with increasing salt in Cheddar cheese. However, the bitterness found in our study was not necessarily associated with salt (-10% of sodium ($\text{g } 100 \text{ g}^{-1}$) in S30) but instead, with a higher pH in these cheeses, especially in S30 (6.28). Lee and Warthesen (1996) reported a high pH in the most bitter cheeses. Furthermore, a significant level of calcium (such as in S30 cheese) contributes to bitterness (Engel, Nicklaus, Septier, Salles, & Le Quéré, 2000). Contrary to other salt substitutions with MgCl_2 or CaCl_2 used in the salting of Cheddar cheese (Grummer et al., 2012), the impact of the calcium lactate salt substitution on off-flavour was not found in this study.

During this investigation, salty taste was generated only by five panellists and did not characterise the commercial cheese Std. It has been reported that food structures impacts the sensory perception of salt, through changes in salt release rate and availability (Busch, Yong, & Goh, 2013; Tournier, Grass, Septier, Bertrand, & Salles, 2014). An important relationship between food texture and masticatory behaviour of subjects was also established (Kohyama, Hatakeyama, Dan, & Sasaki, 2005). For example, the quantity of salt released from a soft cheese is higher than that released from a hard cheese (Phan et al., 2008).

Distribution of other sensory attributes was relatively variable and heterogeneous between judges, suggesting that salting methods have impacted sensory quality of products (notably texture and marbling development) and this effect was differentially perceived between panellists. In summary, in this study, the salting methods had a clear effect on cheese texture with a higher adhesiveness index and a creamy appearance but had little effect on other sensory aspects.

4. Conclusion

We investigated for the first time the relevance of different alternative salting processes to reduce the salt level in a surface dry-salted blue cheese. All the salting methods applied significantly reduced the salt content without altering the sanitary quality of the cheeses. However, several tested methods showed their technical




Fig. 2. Map of the two principal axes of a Generalised Procrustes analysis (GPA) conducted on the flash profile data: A, cheese sample configuration; B, variables plot representing the distribution of subjects (S) according to the generated descriptors.

non feasibility, especially the application of coarse salt, which increases variability in blue cheeses.

Calcium lactate appeared to be a good salt substitute with a salt reduction. Calcium lactate has proven to be a substitute for salt with a 10% reduction in salt and a 21% increase in calcium. Given the variability and heterogeneity of blue-veined cheeses, it would be

interesting to validate the effect of calcium lactate as a salt substitute on the quality of these cheeses.

Regarding the sensory characteristics, a good discrimination between cheese samples and the identification of the most salient sensory attributes of all samples was achieved. Texture, appearance and certain sensory characteristics of the cheese, such as odour,

were highly influenced compared with the salty perception. The bitterness was not representative and there was no off-flavour in this result.

This exploratory study clearly demonstrated the technical feasibility of various salt reduction methods and their impact on the quality of blue cheese. Sprinkling and calcium lactate substitution methods are a practicable technique that could be improved and applied at large scale. Finally, this research has screened a large number of salting methods to identify the most relevant alternatives that can be further investigated in the salting of blue cheeses.

Declaration of competing interest

None.

Acknowledgements

We thank S. Alvarez, K. Fayolle and D. Guerinon (UMR 545 Fromage, Vetagro Sup), for their valuable technical assistance. We also thank B. Desserre for biomolecular analysis help and S. Theil for the bioinformatics support.

This work was supported by FEDER (European Regional Development Funds) in the framework of the call for proposals Pack Ambition Recherche 2018 in the Auvergne-Rhône-Alpes region (France). It was also co-financed by the Syndicat Interprofessionnel Régional du Bleu d'Auvergne (SIRBA) et de la Fourme d'Ambert (SIFAM). We also thank the Pôle Fromager AOP Massif Central for its assistance.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.idairyj.2022.105555>.

References

- Aly, M. E. (1995). An attempt for producing low-sodium Feta-type cheese. *Food Chemistry*, 52, 295–299.
- ANSES. (2012). *Travail relatif au suivi des teneurs en sel des principaux vecteurs entre 2003 et 2011 et simulation des impacts sur les apports en sel de la population française* (p. 23) (Avis No. 2012-SA-0052). <https://www.anses.fr/fr/content/avis-de-l%E2%80%99anses-relatif-au-suivi-des-teneurs-en-sel-des-principaux-vecteurs-entre-2003-et-0>
- ANSES. (2021). *Les références nutritionnelles en vitamines et minéraux*. <https://www.anses.fr/fr/content/les-r%C3%A9f%C3%A9rences-nutritionnelles-en-vitamines-et-min%C3%A9raux>.
- Bansal, V., & Mishra, S. K. (2020). Reduced-sodium cheeses: Implications of reducing sodium chloride on cheese quality and safety. *Comprehensive Reviews in Food Science and Food Safety*, 19, 733–758.
- Berthet, A. (2009). *Nutrition et insuffisance rénale chronique*. <https://dumas.ccsd.cnrs.fr/dumas-01165273>.
- Blaya, J., Barzideh, Z., & LaPointe, G. (2018). Symposium review: Interaction of starter cultures and nonstarter lactic acid bacteria in the cheese environment. *Journal of Dairy Science*, 101, 3611–3629.
- Bord, C., Guerinon, D., & Lebecque, A. (2016). Impact of heating on sensory properties of French Protected Designation of Origin (PDO) blue cheeses. Relationships with physicochemical parameters. *Food Science and Technology International*, 22, 377–388.
- Brown, I. J., Tzoulaki, I., Candeias, V., & Elliott, P. (2009). Salt intakes around the world: Implications for public health. *International Journal of Epidemiology*, 38, 791–813.
- Busch, J. L. H. C., Yong, F. Y. S., & Goh, S. M. (2013). Sodium reduction: Optimizing product composition and structure towards increasing saltiness perception. *Trends in Food Science & Technology*, 29, 21–34.
- Caron, T., Piver, M. L., Péron, A.-C., Lieben, P., Lavigne, R., Brunel, S., et al. (2021). Strong effect of *Penicillium roqueforti* populations on volatile and metabolic compounds responsible for aromas, flavor and texture in blue cheeses. *International Journal of Food Microbiology*, 354, Article 109174.
- Chavhan, G. B., Kanawjia, S. K., Khetra, Y., & Puri, R. (2015). Effect of potassium-based emulsifying salts on sensory, textural, and functional attributes of low-sodium processed Mozzarella cheese. *Dairy Science & Technology*, 95, 265–278.
- Costa, R. G. B., Alves, R. C., Cruz, A. G. da, Sobral, D., Teodoro, V. A. M., Costa Junior, L. C. G., et al. (2018). Manufacture of reduced-sodium Coalho cheese by partial replacement of NaCl with KCl. *International Dairy Journal*, 87, 37–43.
- Dairou, V., & Sieffermann, J.-M. (2002). A comparison of 14 jams characterized by conventional profile and a quick original method, the flash profile. *Journal of Food Science*, 67, 826–834.
- Delahunty, C. M., & Drake, M. A. (2004). Sensory character of cheese and its evaluation. In P. F. Fox, P. L. H. McSweeney, T. M. Cogan, & T. P. Guinee (Eds.), *General aspects: Vol. 1. Cheese: Chemistry, physics and microbiology* (pp. 455–487). Amsterdam, the Netherlands: Elsevier.
- Delarue, J., & Lawlor, B. (2014). *Rapid sensory profiling techniques: Applications in New product development and consumer research*. Amsterdam, the Netherlands: Elsevier.
- Doyle, M. E., & Glass, K. A. (2010). Sodium reduction and its effect on food safety, food quality, and human health. *Comprehensive Reviews in Food Science and Food Safety*, 9, 44–56.
- Dugat-Bony, E., Bonnarne, P., Fraud, S., Catellote, J., Sarthou, A.-S., Loux, V., et al. (2019). Effect of sodium chloride reduction or partial substitution with potassium chloride on the microbiological, biochemical and sensory characteristics of semi-hard and soft cheeses. *Food Research International*, 125, Article 108643.
- Duval, P., Chatelard-Chauvin, C., Gayard, C., Rifa, E., Bouchard, P., Hulin, S., et al. (2016). Microbial dynamics in industrial blue veined cheeses in different packaging. *International Dairy Journal*, 56, 198–207.
- Engel, E., Nicklaus, S., Septier, C., Salles, C., & Le Quéré, J. L. (2000). Taste active compounds in a goat cheese water-soluble extract. 2. Determination of the relative impact of water-soluble extract components on its taste using omission tests. *Journal of Agricultural and Food Chemistry*, 48, 4260–4267.
- Ferroukhi, I., Bord, C., Alvarez, S., Fayolle, K., Theil, S., Lavigne, R., et al. (2022). Functional changes in Bleu d'Auvergne cheese during ripening. *Food Chemistry*, 397, Article 133850.
- Fitzgerald, E., & Buckley, J. (1985). Effect of total and partial substitution of sodium chloride on the quality of Cheddar cheese. *Journal of Dairy Science*, 68, 3127–3134.
- Flórez, A. B., & Mayo, B. (2006). Microbial diversity and succession during the manufacture and ripening of traditional, Spanish, blue-veined Cabrales cheese, as determined by PCR-DGGE. *International Journal of Food Microbiology*, 110, 165–171.
- Galli, B. D., Baptista, D. P., Cavalheiro, F. G., & Gigante, M. L. (2019). *Lactobacillus rhamnosus* GG improves the sensorial profile of Camembert-type cheese: An approach through flash-profile and CATA. *LWT - Food Science and Technology*, 107, 72–78.
- Gkatzionis, K., Hewson, L., Hollowood, T., Hort, J., Dodd, C. E. R., & Linforth, R. S. T. (2013). Effect of *Yarrowia lipolytica* on blue cheese odour development: Flash profile sensory evaluation of microbiological models and cheeses. *International Dairy Journal*, 30, 8–13.
- Gkatzionis, K., Yunita, D., Linforth, R. S. T., Dickinson, M., & Dodd, C. E. R. (2014). Diversity and activities of yeasts from different parts of a Stilton cheese. *International Journal of Food Microbiology*, 177, 109–116.
- Gore, E., Mardon, J., Cécile, B., & Lebecque, A. (2019). Calcium lactate as an attractive compound to partly replace salt in blue-veined cheese. *Journal of Dairy Science*, 102, 1–13.
- Grummer, J., Bobowski, N., Karal, M., Vickers, Z., & Schoenfuss, T. (2013). Use of potassium chloride and flavor enhancers in low sodium Cheddar cheese. *Journal of Dairy Science*, 96, 1401–1418.
- Grummer, J., Karal, M., Zhang, K., Vickers, Z., & Schoenfuss, T. C. (2012). Manufacture of reduced-sodium Cheddar-style cheese with mineral salt replacers. *Journal of Dairy Science*, 95, 2830–2839.
- Guinee, T. P. (2004). Salting and the role of salt in cheese. *International Journal of Dairy Technology*, 57, 99–109.
- Guinee, T. P., & Fox, P. F. (1983). Sodium chloride and moisture changes in Romano-type cheese during salting. *Journal of Dairy Research*, 50, 511–518.
- Guinee, T. P., & Fox, P. F. (2017). In P. L. H. McSweeney, P. F. Fox, P. D. Cotter, & D. W. Everett (Eds.) (4th ed., *CheeseSalt in cheese: Physical, chemical and biological aspects* (pp. 317–375). San Diego, CA, USA: Academic Press.
- Guinee, T. P., & O'Kennedy, B. T. (2007). Reducing salt in cheese and dairy spreads. In D. Kilcast, & F. Angus (Eds.), *Reducing salt in foods* (pp. 316–357). Cambridge, UK: Woodhead Publishing.
- Henneberry, S., Wilkinson, M. G., Kilcawley, K. N., Kelly, P. M., & Guinee, T. P. (2015). Interactive effects of salt and fat reduction on composition, rheology and functional properties of mozzarella-style cheese. *Dairy Science & Technology*, 95, 613–638.
- INAO. (2021). Institut national de l'origine et de la qualité. In *Cahier des charges de l'appellation d'origine « Bleu d'Auvergne »*. <https://www.inao.gov.fr/produit/4562>.
- Kamleh, R., Olabi, A., Toufeili, I., Daroub, H., Younis, T., & Ajib, R. (2015). The effect of partial substitution of NaCl with KCl on the physicochemical, microbiological and sensory properties of Akkawi cheese. *Journal of the Science of Food and Agriculture*, 95, 1940–1948.
- Kamleh, R., Olabi, A., Toufeili, I., Najm, N. E. O., Younis, T., & Ajib, R. (2012). The effect of substitution of sodium chloride with potassium chloride on the physicochemical, microbiological, and sensory properties of Halloumi cheese. *Journal of Dairy Science*, 95, 1140–1151.
- Katsiari, M. C., Voutsinas, L. P., Alichanidis, E., & Roussis, I. G. (1997). Reduction of sodium content in Feta cheese by partial substitution of NaCl by KCl. *International Dairy Journal*, 7, 465–472.
- Katsiari, M. C., Voutsinas, L. P., Alichanidis, E., & Roussis, I. G. (1998). Manufacture of Kefalograviera cheese with less sodium by partial replacement of NaCl with KCl. *Food Chemistry*, 61, 63–70.

- Kohyama, K., Hatakeyama, E., Dan, H., & Sasaki, T. (2005). Effects of sample thickness on bite force for raw carrots and fish gels. *Journal of Texture Studies*, 36, 157–173.
- Le Graet, Y., & Brulé, G. (1988). Migration des macro et oligo-éléments dans un fromage à pâte molle de type Camembert. *Le Lait*, 68, 219–234.
- Le Graet, Y. L., Lepienne, A., Brulé, G., & Ducruet, P. (1983). Migration du calcium et des phosphates inorganiques dans les fromages à pâte molle de type Camembert au cours de l'affinage. *Le Lait*, 63, 317–332.
- Lee, K.-P. D., & Warthesen, J. J. (1996). Preparative methods of isolating bitter peptides from Cheddar cheese. *Journal of Agricultural and Food Chemistry*, 44, 1058–1063.
- Lindsay, R. C., Hargett, S. M., & Bush, C. S. (1982). Effect of sodium/potassium (1:1) chloride and low sodium chloride concentrations on quality of Cheddar cheese. *Journal of Dairy Science*, 65, 360–370.
- Marth, E. H., & Steele, J. (2001). *Applied dairy microbiology* (2nd ed.). Boca Raton, FL, USA: CRC Press.
- McCarthy, C. M., Wilkinson, M. G., Kelly, P. M., & Guinee, T. P. (2015). Effect of salt and fat reduction on the composition, lactose metabolism, water activity and microbiology of Cheddar cheese. *Dairy Science & Technology*, 95, 587–611.
- Meola, M., Rifa, E., Shani, N., Delbès, C., Berthoud, H., & Chassard, C. (2019). DALRYdb: A manually curated reference database for improved taxonomy annotation of 16S rRNA gene sequences from dairy products. *BMC Genomics*, 20, Article 560.
- Mistry, V. V., & Kasperson, K. M. (1998). Influence of salt on the quality of reduced fat Cheddar cheese. *Journal of Dairy Science*, 81, 1214–1221.
- Møller, K. K., Rattray, F. P., Høier, E., & Ardö, Y. (2012). Erratum to: Manufacture and biochemical characteristics during ripening of Cheddar cheese with variable NaCl and equal moisture content. *Dairy Science & Technology*, 92, 541–568.
- Nishinari, K., Kohyama, K., Kumagai, H., Funami, T., & Bourne, M. C. (2013). Parameters of texture profile analysis. *Food Science and Technology Research*, 19, 519–521.
- Pastorino, A. J., Hansen, C. L., & McMahon, D. J. (2003). Effect of salt on structure-function relationships of cheese. *Journal of Dairy Science*, 86, 60–69.
- Phan, V. A., Yven, C., Lawrence, G., Chabanet, C., Reparet, J. M., & Salles, C. (2008). In vivo sodium release related to salty perception during eating model cheeses of different textures. *International Dairy Journal*, 18, 956–963.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2013). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research*, 41, D590–D596.
- Reddy, K. A., & Marth, E. H. (1995). Microflora of Cheddar cheese made with sodium chloride, potassium chloride, or mixtures of sodium and potassium chloride. *Journal of Food Protection*, 58, 54–61.
- Remer, T. (2001). Influence of nutrition on acid-base balance—Metabolic aspects. *European Journal of Nutrition*, 40, 214–220.
- Reykdal, O., & Lee, K. (1993). Validation of chemical measures of calcium with bioassay of calcium-fortified cottage cheese. *Food Chemistry*, 47, 195–200.
- Rulikowska, A., Kilcawley, K. N., Doolan, I. A., Alonso-Gomez, M., Nongonierma, A. B., Hannon, J. A., et al. (2013). The impact of reduced sodium chloride content on Cheddar cheese quality. *International Dairy Journal*, 28, 45–55.
- Schroeder, C. L., Bodyfelt, F. W., Wyatt, C. J., & McDaniel, M. R. (1988). Reduction of sodium chloride in Cheddar cheese: Effect on sensory, microbiological, and chemical properties. *Journal of Dairy Science*, 71, 2010–2020.
- Singh, G., & Muthukumarappan, K. (2008). Influence of calcium fortification on sensory, physical and rheological characteristics of fruit yogurt. *LWT – Food Science and Technology*, 41, 1145–1152.
- Soares, C., Fernando, A. L., Mendes, B., & Martins, A. P. L. (2015). The effect of lowering salt on the physicochemical, microbiological and sensory properties of São João cheese of Pico Island. *International Journal of Dairy Technology*, 68, 409–419.
- Theil, S., & Rifa, E. (2021). rANOMALY: Amplicon workflow for Microbial community Analysis. *F1000Research*, 10, Article 7.
- Tournier, C., Grass, M., Septier, C., Bertrand, D., & Salles, C. (2014). The impact of mastication, salivation and food bolus formation on salt release during bread consumption. *Food & Function*, 5, 2969–2980.
- WHO. (2021). *New WHO benchmarks help countries reduce salt intake and save lives*. <https://www.who.int/news/item/05-05-2021-new-who-benchmarks-help-countries-reduce-salt-intake-and-save-lives>.
- Wongkhalaung, C., & Boonyaratanakornkit, M. (2000). Development of a yogurt-type product from saccharified rice. *Agriculture and Natural Resources*, 34, Article 1.
- Yonis, A. A. M., Elzamzamy, F. M., & Elmorsi, S. A. (2013). Fortification of banana stirred yogurt with calcium. *Journal of Food and Dairy Sciences*, 4, 183–192.