



**HAL**  
open science

## Inhibitory Potential of $\alpha$ -Amylase, $\alpha$ -Glucosidase, and Pancreatic Lipase by a Formulation of Five Plant Extracts: TOTUM-63

Quentin Haguët, Florian Le Joubioux, Vivien Chavanelle, Hugo Groult, Nathan Schoonjans, Cédric Langhi, Arnaud Michaux, Yolanda F Otero, Nathalie Boisseau, Sébastien L Peltier, et al.

### ► To cite this version:

Quentin Haguët, Florian Le Joubioux, Vivien Chavanelle, Hugo Groult, Nathan Schoonjans, et al.. Inhibitory Potential of  $\alpha$ -Amylase,  $\alpha$ -Glucosidase, and Pancreatic Lipase by a Formulation of Five Plant Extracts: TOTUM-63. *International Journal of Molecular Sciences*, 2023, 24 (4), pp.3652. 10.3390/ijms24043652 . hal-03983988

**HAL Id: hal-03983988**

<https://uca.hal.science/hal-03983988v1>

Submitted on 11 Feb 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Article

# Inhibitory Potential of $\alpha$ -Amylase, $\alpha$ -Glucosidase, and Pancreatic Lipase by a Formulation of Five Plant Extracts: TOTUM-63

Quentin Haguet <sup>1,†</sup>, Florian Le Joubioux <sup>2,\*</sup>, Vivien Chavanelle <sup>3</sup>, Hugo Groult <sup>1</sup>, Nathan Schoonjans <sup>2</sup>, Cédric Langhi <sup>3</sup>, Arnaud Michaux <sup>3</sup>, Yolanda F. Otero <sup>3</sup>, Nathalie Boisseau <sup>4</sup>, Sébastien L. Peltier <sup>2</sup>, Pascal Sirvent <sup>3</sup> and Thierry Maugard <sup>1,\*</sup>

<sup>1</sup> UMR 7266 CNRS-ULR, LIENSs, Equipe BCBS, La Rochelle Université, Avenue Michel Crépeau, 17042 La Rochelle, France

<sup>2</sup> Valbiotis, R&D Center, 23 Avenue Albert Einstein, 17000 La Rochelle, France

<sup>3</sup> Valbiotis, R&D Center, 20-22 Rue Henri et Gilberte Goudier, 63200 Riom, France

<sup>4</sup> AME2P, STAPS, Université Clermont Auvergne, 5 Impasse Amélie Murat, 63001 Clermont-Ferrand, France

\* Correspondence: florian.lejoubioux@valbiotis.com (F.L.J.); thierry.maugard@univ-lr.fr (T.M.)

† These authors contributed equally to this work.

**Abstract:** Controlling post-prandial hyperglycemia and hyperlipidemia, particularly by regulating the activity of digestive enzymes, allows managing type 2 diabetes and obesity. The aim of this study was to assess the effects of TOTUM-63, a formulation of five plant extracts (*Olea europaea* L., *Cynara scolymus* L., *Chrysanthellum indicum* subsp. *afroamericanum* B.L.Turner, *Vaccinium myrtillus* L., and *Piper nigrum* L.), on enzymes involved in carbohydrate and lipid absorption. First, in vitro inhibition assays were performed by targeting three enzymes:  $\alpha$ -glucosidase,  $\alpha$ -amylase, and lipase. Then, kinetic studies and binding affinity determinations by fluorescence spectrum changes and microscale thermophoresis were performed. The in vitro assays showed that TOTUM-63 inhibited all three digestive enzymes, particularly  $\alpha$ -glucosidase (IC<sub>50</sub> of 13.1  $\mu$ g/mL). Mechanistic studies on  $\alpha$ -glucosidase inhibition by TOTUM-63 and molecular interaction experiments indicated a mixed (full) inhibition mechanism, and higher affinity for  $\alpha$ -glucosidase than acarbose, the reference  $\alpha$ -glucosidase inhibitor. Lastly, in vivo data using leptin receptor-deficient (db/db) mice, a model of obesity and type 2 diabetes, indicated that TOTUM-63 might prevent the increase in fasting glycemia and glycated hemoglobin (HbA1c) levels over time, compared with the untreated group. These results show that TOTUM-63 is a promising new approach for type 2 diabetes management via  $\alpha$ -glucosidase inhibition.

**Keywords:** plant extracts; type 2 diabetes; cardiometabolic diseases; enzyme assay;  $\alpha$ -glucosidase

**Citation:** Haguet, Q.;

Le Joubioux, F.; Chavanelle, V.;

Groult, H.; Schoonjans, N.;

Langhi, C.; Michaux, A.; Otero, Y.F.;

Boisseau, N.; Peltier, S.L.; et al.

Inhibitory Potential of  $\alpha$ -Amylase,

$\alpha$ -Glucosidase, and Pancreatic

Lipase by a Formulation of Five

Plant Extracts: TOTUM-63. *Int. J.*

*Mol. Sci.* **2023**, *24*, 3652. [https://](https://doi.org/10.3390/ijms24043652)

[doi.org/10.3390/ijms24043652](https://doi.org/10.3390/ijms24043652)

Academic Editors: Dake Qi and

Haibin Tong

Received: 18 January 2023

Revised: 6 February 2023

Accepted: 8 February 2023

Published: 11 February 2023



**Copyright:** © 2023 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article

distributed under the terms and

conditions of the Creative Commons

Attribution (CC BY) license

(<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction


Over the years, the worldwide prevalence of overweight and obesity has steadily increased, becoming a genuine pandemic. The growing incidence of patients with overweight/obesity (671 million people with obesity in 2016) represents an equal amount of potential patients with multiple metabolic pathologies in the future [1]. Currently, obesity is one of the leading causes of morbidity, including type 2 diabetes (T2D), prediabetes, cardiovascular diseases, and non-alcoholic fatty liver disease (NAFLD) [2–5]. Among them, T2D is a major concern due to its high prevalence (11.3% of the U.S. population in 2021) [6]. Hyperglycemia in T2D is caused by reduced insulin secretion and sensitivity, impaired glucose uptake, or increased glucose production associated with pathophysiological changes in multiple organs [7]. To manage these pathological states, various treatments are used with mixed results. Lifestyle changes lead to glycemia control

improvement without side effects, but long-term adherence is generally low. On the other hand, drug-based treatments are not fully satisfactory, mainly due to their side effects and cost [8–10]. As the mechanisms underlying disorders such as components of metabolic syndrome (NAFLD, hypertension or T2D) are complex and interconnected, one often being a risk factor for developing another [11], multiple strategies are available to target these pathophysiological states, and it has now become common for people with T2D to be treated with multiple drug classes [12,13]. Among such strategies, we focused on reducing the intestinal absorption of carbohydrates and lipids to manage diabetes and obesity [14–16]. Carbohydrate absorption is regulated mainly by the digestive enzymatic activity, stomach emptying rate, alimentary bolus composition, and nutrient transporters [17]. It is largely accepted that inhibiting digestive enzymes helps to reduce post-prandial glycemia and to achieve long-term glycemia control. Therefore, their inhibition represents an attractive approach for T2D management [16]. Two classes of enzymes are involved in digestion:  $\alpha$ -amylases (salivary and pancreatic) and  $\alpha$ -glucosidases (intestinal maltase-glucoamylase and sucrase-isomaltase). They successively hydrolyze food polysaccharides into absorbable sugar units [18,19]. Similarly, although numerous parameters regulate lipid absorption, the reduction of gastric and pancreatic lipase activity is a therapeutic strategy that has already shown its efficacy in body weight management [15,20]. Three steps can be distinguished in the digestion of dietary lipids: emulsification, hydrolysis, and micellization. They are followed by lipid uptake by enterocytes [21]. A significant reduction in body weight in patients with obesity and diabetes strongly improves the control of glycemia [22]. Therefore, products capable of reducing both carbohydrate and lipid absorption could be used to effectively manage T2D. Plant extracts seem to be good candidates. Among the many plant biomolecules, phenolic compounds display very potent inhibitory effects on digestive enzymes [23]. In this context, TOTUM-63, a new patented tool, was developed to be used as a dietary supplement for management of obesity and T2D [24]. TOTUM-63 is a naturally biomolecule-rich formulation of five plant extracts: *Olea europaea* L. (olive tree), *Cynara scolymus* L. (artichoke), *Chrysanthellum indicum subsp. afroamericanum* B.L. Turner (chrysanthellum), *Vaccinium myrtillus* L. (bilberry), and *Piper nigrum* L. (black pepper). Several of the extracts included in this formulation are commonly used in nutraceutical formulations or dietary supplements [25–28], but the interest of this project is to test the specific combination of these five plant extracts. A first pilot clinical study in volunteers with overweight showed that TOTUM-63 has good safety and tolerability profiles and improves the glucose and insulin responses to a carbohydrate tolerance test [29]. The present study describes complementary experiments to bring insights into its mechanism of action. Specifically, we assessed TOTUM-63 effects in vitro on three enzymes involved in carbohydrate and lipid absorption (an  $\alpha$ -glucosidase, an  $\alpha$ -amylase, and a lipase) and in vivo on glucose homeostasis (db/db mice). In vitro experiments revealed that TOTUM-63 can significantly reduce the activity of the three digestive enzymes, particularly  $\alpha$ -glucosidase, mainly through a mixed-type inhibition mechanism. The affinity of  $\alpha$ -glucosidase for this formulation was higher than for acarbose, a commercial inhibitor. Furthermore, we demonstrated that many molecules and families of molecules present in TOTUM-63 might participate in its activity. Lastly, in vivo data in db/db mice showed that TOTUM-63 might prevent the rise in fasting glycemia and glycated hemoglobin levels.

## 2. Results

### 2.1. In Vitro Enzymatic Inhibitions

The in vitro inhibitory effects of TOTUM-63 and of reference drugs (acarbose and orlistat, an  $\alpha$ -glucosidase/ $\alpha$ -amylase inhibitor and a lipase inhibitor, respectively) on  $\alpha$ -glucosidase,  $\alpha$ -amylase, and lipase activities are presented in Figure 1 and the  $IC_{50}$  values derived from these curves are in Table 1.



**Figure 1.** Inhibition of an  $\alpha$ -glucosidase by TOTUM-63 (a) and acarbose (b) of an  $\alpha$ -amylase by TOTUM-63 (c) and acarbose (d), and of a lipase by TOTUM-63 (e) and orlistat (f). Values are the mean  $\pm$  SEM (the error bars represent the SEM with  $n = 2$ –9 and the SEM values were  $<10\%$  of the mean values for acarbose or orlistat and  $<16\%$  of the mean values for TOTUM-63).

**Table 1.** IC<sub>50</sub> of TOTUM-63 and reference drugs for  $\alpha$ -glucosidase,  $\alpha$ -amylase, and lipase inhibition.

Tested Products	IC <sub>50</sub> Values in $\mu\text{g/mL}$		
	$\alpha$ -Glucosidase	$\alpha$ -Amylase	Lipase
TOTUM-63	13.1	380.0	276.5
Acarbose	28.8	0.110	-
Orlistat	-	-	0.00345

TOTUM-63 inhibited all three targets but displayed the highest effect on  $\alpha$ -glucosidase (TOTUM-63 IC<sub>50</sub> = 13.1  $\mu\text{g/mL}$  and acarbose IC<sub>50</sub> = 28.8  $\mu\text{g/mL}$ ). The inhibitory effect of TOTUM-63 on  $\alpha$ -amylase was also good (IC<sub>50</sub> = 380  $\mu\text{g/mL}$ ), although the acarbose effect was much stronger (IC<sub>50</sub> = 0.110  $\mu\text{g/mL}$ ). TOTUM-63 inhibition of the lipase was in the same order of magnitude (IC<sub>50</sub> = 276.5  $\mu\text{g/mL}$ ), but orlistat, the reference inhibitor, was much more effective (IC<sub>50</sub> = 0.00345  $\mu\text{g/mL}$ ).

Due to the significant  $\alpha$ -glucosidase inhibition by TOTUM-63, we then tested in vitro  $\alpha$ -glucosidase inhibition by various compounds representative of the families of molecules identified in TOTUM-63 (see TOTUM-63 characterization Table 2) using the same methodology.

**Table 2.** Phytochemical characterization of TOTUM-63.


Compound Types (Sorted by Families)	Extract Content (g/100 g)
<u>Total phenolic compounds</u>	9.1
Total anthocyanins	0.964
Monocaffeoylquinic acids	1.073
Chlorogenic acid	0.645
Other monocaffeoylquinic acids	0.428
Dicaffeoylquinic acids	0.917
Cynarine	0.112
Other dicaffeoylquinic acids	0.805
Caffeic acid	0.019
Oleuropein	3.645
Oleuropein isomers	0.519
Hydroxytyrosol	0.454
Luteolin	0.030
Luteolin-7-O-glucoside	0.656
Luteolin-7-O-glucuronide	0.440
Apigenin	0.016
Apigenin-7-O-glucoside	0.093
Apigenin-7-O-glucuronide	0.324
Apigenin-6-C-glucoside-8-C-arabinoside (Shaftoside)	0.029
Apigenin-6,8-C-diglucoside (Vicenin 2)	0.060
Eriodictyol	0.008
Eriodictyol-7-O-glucoside	0.590
Marein and Flavanomarein	0.318
Maritimein	0.129
Rutin	0.014
Verbascoside	0.046
<u>Terpenes and terpenoids</u>	
Oleanolic acid	2.004
Saponins	
Chrysanthellin A	0.553
Chrysanthellin B	0.507

Iridoids	
Oleoside	0.290
<u>Alkaloids</u>	
Piperine	0.007

Table 3 lists the IC<sub>50</sub> values obtained for five compounds: chlorogenic acid, luteolin, oleuropein, chrysanthellin B, and piperine (Figure 2). These five compounds are representative of the major families of compounds identified in TOTUM-63.

**Table 3.** IC<sub>50</sub> of five phytochemical compounds for  $\alpha$ -glucosidase inhibition.

Tested Compounds	IC <sub>50</sub> Values in $\mu\text{g/mL}$
Chlorogenic acid	0.3
Luteolin	0.4
Oleuropein	3.1
Chrysanthellin B	153.5
Piperine	0.9




**Figure 2.** Chemical structures of the compounds tested for  $\alpha$ -glucosidase inhibition.

The 5 tested compounds significantly inhibited  $\alpha$ -glucosidase activity, as indicated by their IC<sub>50</sub> values that ranged from 0.3  $\mu\text{g/mL}$  for chlorogenic acid to 153.5  $\mu\text{g/mL}$  for chrysanthellin B. Moreover, the inhibitory activity of 4 of these compounds (chlorogenic acid, luteolin, oleuropein, and piperine) was higher than that of acarbose (IC<sub>50</sub> = 0.3  $\mu\text{g/mL}$ , 0.4  $\mu\text{g/mL}$ , 3.1  $\mu\text{g/mL}$ , and 0.9  $\mu\text{g/mL}$ , respectively, versus IC<sub>50</sub> = 28.8  $\mu\text{g/mL}$  for acarbose).

## 2.2. Kinetics and Mechanisms of $\alpha$ -Glucosidase Inhibition

Figure 3 shows the Lineweaver–Burk plots of  $\alpha$ -glucosidase inhibition kinetics in the presence of various concentrations of substrate (*p*-nitrophenyl- $\alpha$ -D-glucopyranoside, PNPg) and of increasing concentrations of TOTUM-63 or acarbose.



**Figure 3.** Lineweaver–Burk plots of  $\alpha$ -glucosidase inhibition in the presence of TOTUM-63 (a) and acarbose (b). Values are the mean  $\pm$  SEM (the error bars represent the SEM with  $n = 9$ –25 for all points in both groups and SEM values were  $\leq 20\%$  of the mean values for acarbose and  $< 10\%$  of the mean values for TOTUM-63). PNP, *p*-nitrophenol; PNPg, *p*-nitrophenyl- $\alpha$ -D-glucopyranoside.

Analysis of these data with the SigmaPlot software 12 (Systat Software Inc., San Jose, CA, USA) indicated a mixed inhibitory mechanism for TOTUM-63 and a competitive inhibitory mechanism for acarbose (as the main mechanisms, they obtained the best  $R^2$  and Akaike Information Criterion corrected for small sample size, AICc values). As described for these mechanisms, the lines intersected to the left of the Y-axis and on the negative side of the X-axis for TOTUM-63, whereas they intersected on the Y-axis ( $X = 0$ ) for acarbose. Moreover, with increasing TOTUM-63 concentrations, both the Y-intercept ( $1/V_{max}$ ) and slope ( $K_m/V_{max}$ ) of the lines increased (decrease in velocity and substrate affinity), whereas with increasing acarbose concentrations, only the slope ( $K_m/V_{max}$ ) of the lines increased (decrease in substrate affinity). Table 4 shows the main kinetic parameters obtained.

**Table 4.** Kinetic parameters of  $\alpha$ -glucosidase inhibition in the presence of TOTUM-63 or acarbose.

Parameters	TOTUM-63	Acarbose
Inhibition type	Mixed (full)	Competitive (full)
$V_{max}$ (mM/min)	0.00259	0.00182
$K_m$ (mM)	0.14	0.06
$K_i$ ( $\mu\text{g/mL}$ )	2.7	4.5
Factor $\alpha$	2.9	none


$V_{max}$ , maximum reaction velocity;  $K_m$ , amount of substrate required to obtain half of the  $V_{max}$  (also called Michaelis constant);  $K_i$ , inhibitor dissociation constant; Factor  $\alpha$ , equilibrium between  $K_i'$  (the second dissociation constant in the mixed model) and  $K_i$ .

Lastly, the  $K_i$  value was lower for TOTUM-63 than for acarbose, and the factor  $\alpha$  value for TOTUM-63 was 2.9, indicating better affinity for the free enzyme than for the enzyme–substrate complex.

### 2.3. Interactions with $\alpha$ -Glucosidase Using Fluorescence Spectrum Changes

To assess the products' interactions with the enzyme, we recorded  $\alpha$ -glucosidase fluorescence emission in the ultraviolet range (320–440 nm), in the absence and presence of TOTUM-63 or acarbose, upon excitation at 285 nm.

The fluorescence intensity gradually decreased with increasing concentrations of TOTUM-63 (Figure 4a). We observed a similar trend with acarbose, but with a less obvious effect, and the fluorescence intensity remained high even at the highest concentrations (Figure 4b). We did not observe any significant shift of the maximum emission wavelength of  $\alpha$ -glucosidase in the presence of TOTUM-63 or acarbose.



**Figure 4.** Fluorescence spectra of  $\alpha$ -glucosidase in the presence of TOTUM-63 (a) and acarbose (b) at various concentrations, upon excitation at 285 nm and emission spectrum between 320 and 440 nm.


#### 2.4. Binding Affinity for $\alpha$ -Glucosidase Using Microscale Thermophoresis (MST)

As detailed by Jerabek-Willemsen et al., 2014 [30], MST is a technique to quantify biomolecular interactions based on thermophoresis. Thermophoresis is the directed movement of molecules through a temperature gradient. This phenomenon can be impacted by several molecular properties such as size, charge, hydration shell, or conformation of the molecule studied. Thus, this technique is very sensitive to any change in molecular properties and allows precise quantification of molecular events independent of the size or nature of the investigated sample. Therefore, MST can detect events such as the binding of molecules to enzymes. The experiment consisted in assessing changes in fluorescence signals caused by a laser that locally increased the temperature. The dissociation constant between a protein and a hypothetical ligand can be determined if an interaction occurs, as the temperature-induced behavior of the protein–ligand couple would differ from that of each free entity.

The binding affinity of TOTUM-63 and acarbose for  $\alpha$ -glucosidase was assessed using MST. We analyzed the results of the experiments with TOTUM-63 (i.e., the ligand) using the Initial Fluorescence Analysis mode in the MO Affinity Analysis software (NanoTemper Technologies, Munich, Germany) because ligand binding directly led to a change in the initial fluorescence (ligand-induced fluorescence changes). We verified that the fluorescence changes recorded for  $\alpha$ -glucosidase tagged with the fluorescent RED-NHS dye were ligand specific (TOTUM-63) and not due to any material loss. With acarbose we did not observe any change in the initial fluorescence and, therefore, we analyzed the data using the traditional MST mode based on the variation of the MST trace responses. Figure 5 shows the dose–response curves for TOTUM-63 and acarbose. The



software analysis gave  $K_d$  values of 0.0173  $\mu\text{g}/\text{mL}$  for TOTUM-63 and of 47.71  $\mu\text{g}/\text{mL}$  for acarbose (i.e., 2758 times lower affinity for acarbose).




**Figure 5.** Fraction of TOTUM-63 and acarbose bound to  $\alpha$ -glucosidase, measured by MST. Values are the mean  $\pm$  SEM.  $n = 2$  or 3 for all points in both groups. Raw fluorescence (in counts) for TOTUM-63 and variation of the normalized fluorescence ( $\Delta\text{FNorm}$ , in %) for acarbose were normalized by expressing them as the fraction bound to the target.

### 2.5. Glucose Homeostasis, Body Weight, and Body Composition in Diabetic db/db Mice

Body weight gain rate was significantly lower in db/db mice treated with TOTUM-63 ( $n = 10$ ; 2.7% of TOTUM-63 added in the food for 6 weeks) than in the untreated control group ( $n = 10$ ). Body weight started to be significantly lower in treated animals from week 4. The difference became more important at week 5 and week 6 (final measurement:  $44.66 \pm 1.99$  g in the control group vs.  $36.29 \pm 1.66$  g in the TOTUM-63 group,  $p < 0.0001$ ). The differences in body weight were only due to fat mass reduction by 26.8% at week 6 ( $26.08 \pm 1.52$  g in the control group vs.  $19.09 \pm 1.13$  g in the TOTUM-63 group,  $p < 0.01$ ). Indeed, food intake remained similar between groups, and the lean mass was not significantly different at the end of the study (week 6) ( $17.02 \pm 0.52$  g in the control group vs.  $16.47 \pm 0.65$  g in the TOTUM-63 group).

As observed for body weight, fasting glycemia was significantly lower in the TOTUM-63 group from week 4 compared with controls ( $132.1 \pm 14.4$  vs.  $266.0 \pm 40.0$  mg/dL,  $p < 0.0001$ ). At the study end (week 6), fasting blood glucose level was 2.74 times lower in the TOTUM-63 group ( $159.8 \pm 20.4$  mg/dL versus  $437.4 \pm 26.4$  mg/dL in the control group,  $p < 0.0001$ ) (Figure 6a). Glycohemoglobin (HbA1c) levels at week 6 were also significantly lower in the TOTUM-63 group than in controls ( $4.25 \pm 0.20\%$  vs.  $8.33 \pm 0.53\%$ ;  $-1.96\%$ ,  $p < 0.0001$ ) (Figure 6b).



**Figure 6.** Evolution of fasting blood glucose levels (a) and HbA1c levels at week 6 (b) in db/db mice. \*\*\*\*  $p < 0.0001$  (two-way ANOVA followed by the Sidak multiple comparisons test). \$\$\$\$  $p < 0.0001$  (Welch corrected unpaired  $t$ -test). Values are the mean  $\pm$  SEM.  $n = 9$  or 10 for all measurements. db/db, leptin receptor-deficient mice; HbA1c, glycated hemoglobin.

### 3. Discussion

In the present study, we showed that TOTUM-63 can inhibit some digestive enzymes related to the carbohydrate and lipid metabolisms. We observed the strongest effects of TOTUM-63 on  $\alpha$ -glucosidase activity. We thoroughly explored this inhibitory activity through in vitro interaction and affinity studies. Then, in vivo experiments confirmed the potential of TOTUM-63 to prevent body weight gain and glycemia increase in a mouse model of obesity and T2D.

Acarbose is one of the most widely known and used T2D management drugs, particularly through the inhibition of digestive enzymes [31]. In vitro, acarbose strongly inhibited  $\alpha$ -glucosidase ( $IC_{50} = 28.8 \mu\text{g/mL}$ ), but the inhibitory effect of TOTUM-63 was stronger ( $IC_{50} = 13.1 \mu\text{g/mL}$ ). The results obtained with acarbose are consistent with a previous study that reported comparable  $IC_{50}$  values [32]. As described in Table 2, the mixture of plant extracts in TOTUM-63 formulation is rich in phenolic compounds, terpenoids, and alkaloids. Any of these components may participate in the observed activities, as well as other non-listed constituents. Moreover, a synergy between components may exist [33,34]. Indeed, synergistic interactions are documented for biomolecules within a single plant extract, as well as between different plant extracts in a formulation. In these cases, the synergistic effects have been demonstrated by a better efficacy of the formulation than additional effects of the isolated constituents (biomolecules or plant extracts) [35]. Thus, we tested the inhibitory effect on  $\alpha$ -glucosidase of five compounds that are representative of the families of molecules identified in TOTUM-63. The three phenolic compounds (chlorogenic acid, luteolin, and oleuropein), a terpenoid (chrysanthellin B: a saponin based on oleanolic acid) and an alkaloid (piperine) showed significant inhibitory activity (Table 3) that was stronger than that of acarbose for four of them (chlorogenic acid, luteolin, oleuropein, and piperine). Several studies have already evaluated the effects of chlorogenic acid and luteolin on  $\alpha$ -glucosidase showing that these compounds have higher inhibitory activity than acarbose, the reference inhibitor [36–38]. The inhibitory activities of oleuropein and piperine on  $\alpha$ -glucosidase have been less studied and the results are conflicting. Hadrich et al. [39] reported that oleuropein  $IC_{50}$  was two times higher than that of acarbose, while Dekdouk et al. [40] found the opposite. In our in vitro inhibition experiments, the  $IC_{50}$  of oleuropein was 10 times lower than that of acarbose. Similarly, Kumar et al. [41] showed that the  $IC_{50}$  of piperine was 28 times higher than that of acarbose.

Conversely, Tolmie et al. [42] found similar inhibitory activities ( $K_i$ ) for piperine and acarbose, and our *in vitro* data gave an  $IC_{50}$  for piperine 30 times lower than that for acarbose. Despite these differences, all studies showed a significant  $\alpha$ -glucosidase inhibition by oleuropein and piperine. The inhibitory effect of chrysanthellin B on  $\alpha$ -glucosidase was five times lower than that of acarbose but was still significant. We did not find any previous studies on  $\alpha$ -glucosidase inhibition by chrysanthellin B. However, the effects of other molecules from the same sub-families (saponins based on oleanolic acid or oleanane) on this enzyme have been studied and the reported  $IC_{50}$  values were close to those of acarbose [43,44]. In addition, chlorogenic acid, luteolin, oleuropein, chrysanthellin B, and piperine are precursors or derivatives of several other compounds identified in TOTUM-63 (see Table 2) that could also potentially contribute to  $\alpha$ -glucosidase inhibition. For instance, cyanidin (one of TOTUM-63 anthocyanins), apigenin, luteolin-7-O-glucoside, and oleanolic acid exert strong inhibitory effects on  $\alpha$ -glucosidase [45–48]. Oleanolic acid showed the strongest effect with 162 times higher inhibitory activities on  $\alpha$ -glucosidase compared with acarbose ( $IC_{50} = 4.09 \mu\text{M}$  for oleanolic acid and  $IC_{50} = 665.56 \mu\text{M}$  for acarbose) [49]. Moreover, other studies have demonstrated major or minor inhibitory effects on  $\alpha$ -glucosidase for several other compounds identified in TOTUM-63: apigenin-7-O-glucuronide, dicaffeoylquinic acids (including cynarin), eriodictyol, marein, flavanomain, and apigenin-6,8-C-diglucoside (vicenin 2) [50–54]. All these results confirm that  $\alpha$ -glucosidase inhibition by TOTUM-63 is not due to a single specific molecule, but that many molecules and families of molecules present in the TOTUM-63 formulation contribute to this effect.

TOTUM-63 inhibitory effects on  $\alpha$ -amylase and lipase are comparable with those reported by other authors who tested other plant extracts, but they were much lower than those observed with the reference drugs acarbose and orlistat [55–57]. However, it should be noted that excessive inhibition of  $\alpha$ -amylase and lipase is associated with side effects. Therefore, the mild effect of TOTUM-63 on these enzymes could represent an advantage and might limit the risk of gastrointestinal disturbances and vitamin malabsorption, in particular [22,58,59]. Furthermore, as previously described for  $\alpha$ -glucosidase, several studies have demonstrated significant inhibitory effects on  $\alpha$ -amylase or lipase for many compounds identified in TOTUM-63. Significant  $\alpha$ -amylase inhibitory activities have been shown for apigenin, apigenin-7-O-glucuronide, oleanolic acid, anthocyanins, monocaffeoylquinic acids (including chlorogenic acid), dicaffeoylquinic acids (including cynarin), luteolin, eriodictyol, and marein [49–52,55,60–62]. The family of compounds showing the strongest effect were the di-caffeoylquinic acids with 4.8, 6.2, 7.2, 9.4, and 11.4 times higher inhibitory activities on  $\alpha$ -amylase than acarbose for cynarin, 1,5-dicaffeoylquinic acid, 3,4-dicaffeoylquinic acid, 3,5-dicaffeoylquinic acid, and 4,5-dicaffeoylquinic acid, respectively (based on  $IC_{50}$  values) [51]. Similarly, significant lipase inhibitory activities have been shown for apigenin, oleanolic acid, anthocyanins, monocaffeoylquinic acids (including chlorogenic acid), dicaffeoylquinic acids (including cynarin), luteolin-7-O-glucoside, luteolin, and Eriodictyol [63–68]. The highest lipase inhibitions were obtained using luteolin-7-O-glucoside and eriodictyol, two flavonoids, with more than two times higher inhibitory activities than orlistat for both (based on  $IC_{50}$  values) [68]. All these results confirm that  $\alpha$ -amylase and lipase inhibitions by TOTUM-63 are not due to a single specific molecule, but that many molecules and families of molecules present in TOTUM-63 formulation contribute to these effects.

Then, we studied TOTUM-63 mechanism of action by monitoring its interaction with  $\alpha$ -glucosidase because it was the most sensitive target enzyme. Analysis of  $\alpha$ -glucosidase inhibition kinetics showed the expected results for acarbose, confirming its widely described competitive inhibitory mechanism [45]. The results obtained for TOTUM-63 fitted with a mixed inhibitory mechanism. However, given the various biomolecules contained in this formulation, many other inhibitory mechanisms may also be involved to a lesser extent [46–48,55–57,69]. Indeed, the mixed model could be partially explained by the activities of cyanidin, luteolin-7-O-glucoside, oleanolic acid, or apigenin, known to inhibit

$\alpha$ -glucosidase according to this model [45–48]. For the other molecules with demonstrated inhibitory effect on  $\alpha$ -glucosidase, various inhibition mechanisms have been described in the literature. Chlorogenic acid, luteolin, and piperine are considered mixed, non-competitive, and mixed inhibitors, respectively [36,38,42]. Furthermore, molecular docking simulation studies showed that apigenin, luteolin, and cyanidin bind to sites close to the active site of  $\alpha$ -glucosidase, which may induce conformational changes and/or a channel closure to prevent the substrate access, eventually leading to  $\alpha$ -glucosidase inhibition [38,45,47]. In the same way, Deng et al. [49] performed docking simulation for  $\alpha$ -glucosidase using oleanolic acid oxime ester derivatives as potential inhibitors and they demonstrated the oleanolic acid part of the derivative located inside of the enzyme active site. Regarding caffeoylquinic acids (mono- or di-), molecular docking results showed different binding energies and modes with  $\alpha$ -glucosidase depending on their structures and especially the caffeoyl groups' distribution [70]. Thus, the various molecules identified in TOTUM-63 would act on several sites of the enzymes. This seems to confirm that the mechanism of  $\alpha$ -glucosidase inhibition by TOTUM-63 is influenced by all the compounds present in TOTUM-63, which predominantly exhibit mixed inhibition mechanisms.

The study of kinetic parameters revealed that the affinity of TOTUM-63 inhibitory biomolecules ( $K_i$ ) for  $\alpha$ -glucosidase was 1.67 times higher than that of acarbose, in agreement with the  $IC_{50}$  values ( $\sim 2.2$  times lower). The MST technique revealed that both TOTUM-63 and acarbose could bind to  $\alpha$ -glucosidase, modifying the enzyme response to a local temperature gradient. However, TOTUM-63's affinity for  $\alpha$ -glucosidase was much higher than that of acarbose (by 2758 times). Two hypotheses could be made. The main active TOTUM-63 component (for this target) has a very high capacity to bind to  $\alpha$ -glucosidase, but a relatively low effect on its activity compared with acarbose, thus leading to  $\sim$ two times higher inhibitory effect ( $IC_{50}$  and  $K_i$ ). Alternatively, many molecules in the plant extract mixture could bind to  $\alpha$ -glucosidase, possibly at multiple sites; however, most of them do not exert any effect on the enzyme activity. Indeed, changes in an enzyme's endogenous fluorescence intensity and spectra can be caused by quenching due to binding followed by conformational changes, or by a change in the enzyme microenvironment, such as a small distance ( $<10$  nm) between the ligand and the enzyme fluorescent amino acids [71,72]. Similarly, for the MST results, the more important fluorescence spectra changes upon excitation at 280 nm in the presence of TOTUM-63, compared with acarbose, could be explained by multiple molecules interacting with  $\alpha$ -glucosidase elsewhere than at the active site. This would be in line with the allosteric regulation expected for the mixed inhibition shown by TOTUM-63. Conversely, acarbose can only bind to the active site on  $\alpha$ -glucosidase, thus causing fewer fluorescence changes [46,73,74].

Based on these results, we hypothesized that potent activities could also be observed *in vivo*. Therefore, we assessed TOTUM-63's effects on glucose homeostasis and on body weight in db/db mice, a model of T2D [75]. At the end of the study, fasting blood glucose and HbA1c levels were significantly lower in mice supplemented with TOTUM-63 for 6 weeks compared with the control group, suggesting an improvement of glucose homeostasis. Moreover, we observed reduced fat mass gain, resulting in lower body weight gain in supplemented animals. Interestingly, these beneficial effects are often reported when studying the effects of drugs capable of inhibiting digestive enzymes such as glucosidases or lipases *in vivo*. Specifically,  $\alpha$ -glucosidase inhibitors have shown beneficial effects in blunting post-prandial hyperglycemia, resulting in modest improvements in HbA1c in db/db mice [76]. This class of drugs works by reducing the activity of sucrase and maltase in the small intestine [77], resulting in less monosaccharides being produced out of complex carbohydrates. Polysaccharides are unable to be up-taken by the intestinal transporters, hence  $\alpha$ -glucosidase inhibitors ingestion results in less sugar being absorbed overall. Interestingly in humans, long-term HbA1c reduction is also reported [78], sometimes associated to a modest decrease in body weight as well [79]. Similarly, pharmacologic inhibition of pancreatic lipase is a common and approved strategy to reverse obesity, resulting in body weight loss in humans [80]. Here again, the mechanism involved in the anti-

obesity effects is rather simple, as lipase inhibitors prevent part of the ingested triglycerides to be hydrolyzed into simple monoacylglycerol and fatty acids, the only forms able to be transported by the enterocyte [21]. This results in more fat being excreted into the feces, and hence, less energy being absorbed. Interestingly, just like in the present study, this reduction in body weight is associated with improved fasting glycemia and reduced HbA1c levels in humans, suggesting an improvement of glucose control [80].

Furthermore, in addition to all the hypotheses related to the digestive enzymes previously described, there are several other hypotheses that may explain the effect on the HbA1c level of TOTUM-63, a polyphenol-rich formulation. One of these hypotheses could be a direct effect of the phenolic compounds in TOTUM-63 on the glycation of hemoglobin. Indeed, several studies have already demonstrated the deglycation effects of several flavonoids such as luteolin on different proteins including hemoglobin [81,82].

The pathophysiology of metabolic diseases such as T2D being complex, interconnected, and rooted in several key organs, this work does not allow discrimination of the individual in vitro effect on each enzyme on glucose control or fat mass. Incidentally, a reduction in adiposity alone is a strong stimulus for improvement of glucose control [83], which has subsequently made body weight management an integral part of the recommendations of major health organizations worldwide for people with prediabetes or T2D, to improve glucose metabolism [84]. Hence, even though we could not demonstrate a direct and causal link between in vitro and in vivo effects, the fact that the inhibitory effect on digestive enzymes translates into improvements of body composition and glucose homeostasis in a well-known and extensively used mouse model of T2D certainly adds to the understanding of the complex mechanisms involved in the effects of Totum-63 in vivo.

## 4. Materials and Methods

### 4.1. Materials and Reagents

$\alpha$ -Glucosidase (maltase from *Saccharomyces cerevisiae*),  $\alpha$ -amylase (type VI-B, from porcine pancreas), lipase (type II, from porcine pancreas), *p*-nitrophenyl- $\alpha$ -D-glucopyranoside (PNPg), *p*-nitrophenol (PNP), 2-chloro-4-nitrophenyl-4-O- $\beta$ -D-galactopyranosyl- $\alpha$ -D-maltoside (GalG<sub>2</sub>CNP), 2-chloro-4-nitrophenol (CNP), 4-methylumbelliferyl oleate (4-MUO), 4-methylumbelliferone (4-MU), orlistat, acarbose, pluronic F-127, Folin–Ciocalteu reagent, Na<sub>2</sub>CO<sub>3</sub>, Ca(OAc)<sub>2</sub>, NaCl, gallic acid, potassium phosphate buffer, phosphate-citrate buffer, chlorogenic acid, luteolin, oleuropein, piperine, absolute ethanol, and dimethyl sulfoxide (DMSO) were obtained from Merck/Sigma-Aldrich (Darmstadt, Germany). Chrysanthellin B was purchased from Extrasynthese (Genay, France). 2-(N-morpholino)ethanesulfonic acid (MES) buffer was purchased from ThermoFisher Scientific (Waltham, Massachusetts, USA). NaN<sub>3</sub> was obtained from Carlo Erba Reagents (Val-de-Reuil, France). Pure water was obtained using an Evoqua ultra-pure water production system (Water Technologies, Günzburg, Germany). The RED-NHS protein labeling kit was from NanoTemper Technologies (Munich, Germany).

TOTUM-63 (Batch No. V190033) was supplied as a powder by VALBIOTIS (Perigny, France). This product is formulated with five standardized plant extracts (*Olea europaea* L., *Cynara scolymus* L., *Chrysanthellum indicum* subsp. *afroamericanum* B.L.Turner, *Vaccinium myrtillus* L., and *Piper nigrum* L.). Voucher specimens of batches from TOTUM-63 and all plant extracts were deposited and stored in the VALBIOTIS sample library. Table 2 shows the phytochemical characterization of TOTUM-63. Total phenolic compound levels (in gallic acid equivalent) were assessed using the Folin–Ciocalteu colorimetric method [85] and a more precise characterization of phytochemical compounds was performed by HPLC-UV/Visible-MS using a 1200 LC system with a 6110 Single Quad MS-ESI detector (Agilent Technologies, Santa Clara, CA, USA) with a C18 Prodigy reversed-phase column (250 mm × 4.6 mm, 5  $\mu$ m; Phenomenex, USA).

## 4.2. In Vitro Enzymatic Inhibition Assays

### 4.2.1. $\alpha$ -Glucosidase Inhibition Assay

$\alpha$ -Glucosidase hydrolytic activity was assessed by monitoring the conversion of *p*-nitrophenyl- $\alpha$ -D-glucopyranoside (PNP<sub>g</sub>) into *p*-nitrophenol (PNP). PNP levels were quantified at 405 nm. A commercial reference inhibitor, acarbose, was used as positive control, and potassium phosphate buffer (0.1 M, pH 6.8) was used as a reaction buffer. A series of PNP dilutions in buffer was used to generate the standard curve to convert PNP absorbance values into concentrations. The final  $\alpha$ -glucosidase concentration corresponded to an activity of 30  $\mu$ M of PNP/min for negative control without inhibitor. In each microplate well, 20  $\mu$ L of test products diluted in a 50:50 (v:v) water:ethanol mixture (or 20  $\mu$ L of this solvent mixture for the negative controls), 20  $\mu$ L of  $\alpha$ -glucosidase in buffer (or 20  $\mu$ L of buffer for the blanks), and 100  $\mu$ L of buffer were incubated in a microplate reader at 37 °C for 10 min. Then, 20  $\mu$ L of PNP<sub>g</sub> substrate (2.5 mM in buffer) was added to each well to initiate the reaction. A FLUO Star Omega (BMG LabTech, Champigny sur Marne, France) 96-well microplate reader, thermostatically controlled at 37 °C, was used to measure the absorbance at 405 nm, every minute for 30 min. The relative enzymatic activity was calculated using Equation (1), where the negative control activity was the initial reaction rate in the absence of any compound, and the residual activity was the initial reaction rate in the presence of test products. The IC<sub>50</sub> of each compound corresponded to the lowest concentration at which the glucosidase activity was halved.

$$\% \text{ Relative activity} = \frac{\text{Residual activity}}{\text{Negative control activity}} \times 100 \quad (1)$$

### 4.2.2. $\alpha$ -Amylase Inhibition Assay

$\alpha$ -amylase hydrolytic activity was assessed by monitoring the conversion of 2-chloro-4-nitrophenyl-4-O- $\beta$ -D-galactopyranosyl- $\alpha$ -D-maltoside (GalG<sub>2</sub>CNP) into 2-chloro-4-nitrophenol (CNP). CNP levels were quantified at 400 nm. A commercial reference inhibitor, acarbose, was used as the positive control, and MES buffer (50 mM, pH 6), with 5 mM Ca(OAc)<sub>2</sub>, 51.5 mM NaCl, and 152 mM NaN<sub>3</sub>, was used as the reaction buffer. A series of CNP dilutions in buffer was used to generate the standard curve to convert absorbance values into CNP concentrations. The final  $\alpha$ -amylase concentration corresponded to an activity of 200  $\mu$ M CNP/min for the negative control without inhibitor. In each microplate well, 20  $\mu$ L of test products diluted in a 50:50 (v:v) water:ethanol mixture (or 20  $\mu$ L of this solvent mixture for the negative controls), 20  $\mu$ L of  $\alpha$ -amylase in buffer (or 20  $\mu$ L of buffer for the blanks), and 100  $\mu$ L of buffer were incubated in a microplate reader at 37 °C for 10 min. Then, 20  $\mu$ L of GalG<sub>2</sub>CNP substrate (0.3125 mM in buffer) was added to each well to initiate the reaction. A FLUO Star Omega (BMG LabTech, Champigny sur Marne, France) 96-well microplate reader, thermostatically controlled at 37 °C, was used to measure the absorbance at 400 nm, every minute for 30 min. The relative enzymatic activity was calculated using Equation (1). The IC<sub>50</sub> of each compound corresponded to the lowest concentration at which the amylase activity was halved.

### 4.2.3. Lipase Inhibition Assay

Lipase hydrolytic activity was assessed by monitoring the conversion of 4-methylumbelliferyl oleate (4-MUO) into 4-methylumbelliferone (4-MU). The enzymatic activity was determined by measuring the 4-MU fluorescence signal at 460 nm (after excitation at 355 nm) over time. A commercial reference inhibitor, orlistat, was used as the positive control, and citrate-phosphate buffer (0.1 M, pH 7.4) was used as the reaction buffer. A series of 4-MU dilutions in buffer was used to generate the standard curve to convert absorbance values into 4-MU concentrations. A 4 g/L solution of lipase in buffer was centrifuged at 5000 $\times$  g and 10 °C for 10 min, then the supernatant was aliquoted. The final on-well concentration used in the assay was 250 mg/L. In each microplate well, 20

$\mu\text{L}$  of test products diluted in a 10:90 (v:v) DMSO:water mixture (or 20  $\mu\text{L}$  of this solvent mixture for the controls), 20  $\mu\text{L}$  of lipase in buffer (or 20  $\mu\text{L}$  of buffer for the blanks), and 110  $\mu\text{L}$  of buffer were incubated at 37 °C in a microplate reader for 10 min. Then, 10  $\mu\text{L}$  of 4-MUO substrate (48  $\mu\text{M}$  in DMSO) was added to each well to initiate the reaction. Fluorescence was measured at 355 nm/460 nm (excitation/emission) using a FLUO Star Omega (BMG LabTech, Champigny sur Marne, France) 96-well microplate reader, thermostatically controlled at 37 °C, every 30 s for 15 min. The relative enzymatic activity was calculated using Equation (1). The  $\text{IC}_{50}$  of each compound corresponded to the lowest concentration at which the lipase activity was halved.

#### 4.3. $\alpha$ -Glucosidase Inhibition Kinetics

$\alpha$ -glucosidase inhibition kinetics was investigated in vitro using the method described in Section 4.2.1. Inhibition models were determined by analysis of the Lineweaver–Burk plots using four concentrations of substrate and four concentrations of test products (including a “0”, without inhibitor). All initial rates were fitted to all enzymatic kinetic models with one substrate and one inhibitor provided by the Enzyme Kinetics Module of Sigma Plot 12 (Systat Software Inc., San Jose, California, USA), using non-linear least-square regression models to determine the inhibition model and calculate the inhibitory constant ( $K_i$ ). The mechanism of  $\alpha$ -glucosidase inhibition by TOTUM-63 was confirmed using Lineweaver–Burk plots ( $1/[S]$  against  $1/v$ ). The acarbose inhibition model was verified and compared with the literature to validate the obtained results [86].

Equations (2) and (3) (below) were used for data analysis and corresponded to the inhibition models that best fitted our data according to the SigmaPlot software 12 (Systat Software Inc., San Jose, California, USA)[73]. Equation (2), for acarbose, corresponded to the competitive (full) inhibition mechanism where  $v$  corresponds to the reaction velocity (also called reaction rate),  $V_{\text{max}}$  to the maximum reaction velocity,  $K_m$  to the amount of substrate required to obtain half of the  $V_{\text{max}}$  (also called Michaelis constant),  $K_i$  to the inhibitor dissociation constant,  $[S]$  to the substrate concentration, and  $[I]$  to the inhibitor concentration. Equation (3), for TOTUM-63, corresponded to the mixed (full) inhibition mechanism with a new factor,  $K_i'$ , introduced compared with Equation (2).  $K_i'$  corresponds to the second inhibitor dissociation constant because, in a mixed model, the constants for  $E + I \leftrightarrow EI$  and  $ES + I \leftrightarrow ESI$  are different. Factor  $\alpha$ , given by SigmaPlot (Systat Software Inc., San Jose, California, USA), represents the equilibrium between  $K_i'$  and  $K_i$ .

$$v = \frac{V_{\text{max}}}{1 + \left(\frac{K_m}{[S]}\right) \times \left(1 + \frac{[I]}{K_i}\right)} \quad (2)$$

$$v = \frac{V_{\text{max}}}{\left(1 + \frac{[I]}{K_i'}\right) + \frac{K_m}{[S]} \times \left(1 + \frac{[I]}{K_i}\right)} \quad (3)$$

#### 4.4. Determination of Interactions with $\alpha$ -Glucosidase Based on Fluorescence Spectrum Changes

The fluorescence spectra of  $\alpha$ -glucosidase in the absence and presence of TOTUM-63 or acarbose were recorded with a microplate fluorescence spectrophotometer (Spark 10M, Tecan, Männedorf, Switzerland). Potassium phosphate buffer (0.1 M, pH 6.8) was used as the reaction buffer. The final  $\alpha$ -glucosidase concentration in the wells was 1.275 mg/mL. For each microplate well, 20  $\mu\text{L}$  of test product in a 50:50 (v:v) water:ethanol mixture (or 20  $\mu\text{L}$  of this solvent mixture for the controls without products), 100  $\mu\text{L}$  of  $\alpha$ -glucosidase in buffer (or 100  $\mu\text{L}$  of buffer for the blanks), and 40  $\mu\text{L}$  of buffer were incubated at 25 °C in a microplate reader for 5 min. Then, fluorescence was measured at an excitation of 285 nm and an emission scan ranging from 320 nm to 440 nm.

#### 4.5. Determination of the Binding Affinity for $\alpha$ -Glucosidase Using Microscale Thermophoresis (MST)

The binding affinity of TOTUM-63 and acarbose for  $\alpha$ -glucosidase was assessed using MST.  $\alpha$ -glucosidase was labeled with the fluorescent RED-NHS dye using the RED-NHS protein labeling kit (NanoTemper Technologies, Munich, Germany) according to the manufacturer's instructions. Briefly, the dye and enzyme were incubated and the tagged  $\alpha$ -glucosidase was purified. The resultant tagged solution was then diluted in PBS supplemented with Pluronic F-127 to achieve a final  $\alpha$ -glucosidase concentration of 100 nM with 0.01125% Pluronic F-127 in the capillaries. The 2 test products, TOTUM-63 and acarbose, were prepared at their respective optimized concentrations of 0.3125 g/L and 10 g/L before 2-fold serial dilutions for 15 times to obtain ligand concentrations ranging from 0.3125 g/L to 9.5367  $\mu$ g/L and from 10 g/L to 305.1758  $\mu$ g/L, respectively. Standard capillaries for red fluorescence emission recording were filled with a 1:1 (v/v) mixture of tagged enzyme and diluted ligand. Fluorescence was measured using the NanoTemper Monolith NT.115 instrument (NanoTemper Technologies, Munich, Germany) at 37 °C, with a fluorescence excitation power of 80% and MST power set to Medium. Data were analyzed with the NanoTemper analysis software (NanoTemper Technologies, Munich, Germany).

#### 4.6. Animals and Diet

Five-week-old male db/db mice (BKS(D)-Leprdb/JOrlRj) were purchased from JANVIER LABS (Le Genest-Saint-Isle, France) and housed in individual cages with a normal 12 h/12 h light/dark cycle. After 1 week of acclimatization, 20 5-week-old male db/db mice (BKS(D)-Leprdb/JOrlRj) were randomly assigned to Control ( $n = 10$ ) or TOTUM-63 ( $n = 10$ ) groups to obtain 2 groups with comparable fasting glycemia and body weight. Animals belonging to the control group were fed a standard A03 diet enriched with 3% corn oil (Safe, France). Animals in the TOTUM-63 group were fed the same diet supplemented with 2.7% w/w TOTUM-63, incorporated directly into the chow by the manufacturer. The optimum dose (2.7%) was chosen during our preliminary experiments when plant extracts were tested individually and in combination at different ratios (unpublished data), to result in the final composition. All procedures were approved by the local ethics committee (C2EA-02, Auvergne, France) under number 00898.01.

#### 4.7. Glucose Homeostasis, Body Weight, and Body Composition in Diabetic db/db Mice

Body weight and food intake were recorded weekly. After an overnight fast, glycemia was measured weekly using a drop of whole blood taken from the tail and an Accu-Chek Performa blood glucose meter (Roche, Mannheim, Germany). After 6 weeks of treatment, fat and lean mass were recorded by MRI (EchoMRI 3-in-1 instrument; Echo Medical Systems, Houston, TX, USA). Finally, animals were fasted overnight (16 h) and euthanized by cervical dislocation. Whole blood was harvested right after the death of the mice and a small portion was collected in an EDTA-treated tube and put on ice until analysis. Glycated hemoglobin (HbA1c) levels were measured in EDTA-treated whole blood with a mouse Hemoglobin A1c (HbA1c) Assay Kit (Crystal Chem, Elk Grove Village, IL, USA) by comparing absorbance at 700 nm in a spectrometer with HbA1c controls according to the instructions of the manufacturer.

#### 4.8. Statistical Analysis of In Vivo Data

Prism V.7.0 and 8.0 (GraphPad Software, San Diego, CA, USA) and the SAS/STAT software V.9.4 (SAS Institute, Cary, NC, USA) were used for statistical analyses and figure drawing. For pair comparisons, the Shapiro–Wilk normality test was used to determine whether data followed a Gaussian distribution, and then the F-test (Fisher) was used to compare variances. If normal distribution was assumed and variances were homogenous, the unpaired t-test was used to compare groups. If variances were heterogenous, the unpaired t-test with Welch correction was used. If data were not consistent with a Gaussian



distribution, the Mann–Whitney test was used. For measurements repeated over time, differences were tested using a repeated-measure two-way ANOVA followed by the Sidak post hoc test for multiple comparisons if the interaction was statistically significant. Values are presented as the mean  $\pm$  SEM, unless specified otherwise. Differences were considered statistically significant at  $p < 0.05$ .

## 5. Conclusions

Controlling post-prandial hyperglycemia and hyperlipidemia allows managing T2D. Therefore, it is important to identify non-drug strategies to control these parameters. Our *in vivo* study in db/db mice (a model of obesity and T2D) showed that TOTUM-63, a naturally biomolecule-rich formulation of five plant extracts, improves glycemic control with fasting blood glucose and HbA1c levels were significantly lower compared with the control group. An inhibition of digestive enzyme activity may partly explain the improved glucose homeostasis. Indeed, *in vitro* study revealed that TOTUM-63 can reduce digestive enzyme activity, particularly  $\alpha$ -glucosidase activity with a mixed-type inhibition mechanism. TOTUM-63 acted on targets regulated by commonly used drugs (e.g., acarbose), with higher activity than these drugs which bodes well for its potential use in humans [31]. Thus, TOTUM-63 is an attractive candidate for the management of hyperglycemia and T2D.

**Author Contributions:** Conceptualization, S.L.P., P.S. and T.M.; methodology, Q.H., F.L.J., V.C., H.G. and N.S.; validation, Q.H., F.L.J. and P.S.; formal analysis, Q.H., F.L.J., V.C. and N.S.; investigation, Q.H., F.L.J., V.C., N.S., C.L. and A.M.; data curation, Q.H., F.L.J. and V.C.; writing—original draft preparation, Q.H. and F.L.J.; writing—review and editing, F.L.J., V.C., H.G., N.S., C.L., A.M., Y.F.O., N.B., P.S. and T.M.; visualization, Q.H.; supervision, P.S. and T.M.; funding acquisition, S.L.P. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was financially supported by Valbiotis (France). This study was also supported by a FEDER grant and the Auvergne Rhône Alpes region (France).

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the local Ethics Committee C2E2A, Auvergne, France (project n.r 01326.02 authorized on 13 January 2015).

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The data presented in this study are available on request to the corresponding author.

**Acknowledgments:** The authors wish to thank Mehdi Djelloul-Mazouz (UMR 1019 UNH, INRAE, Theix), Valérie Rousseau (UMR 0454, UCA, Clermont-Ferrand), and Christophe De L’Homme (UMR 1019 UNH, INRAE, Theix) for the care and the attention to the animals. We are also grateful to Geoffrey Delcros (Valbiotis, Riom), Monique Etienne (AME2P, UCA, Clermont-Ferrand), Clara Laurent (Valbiotis, Riom), and Mahëva Maura (Valbiotis, La Rochelle) for their technical skills in laboratory experiments. We thank Odd-Erik Johansen (Nestlé Health Science, Vevey) and Elisabetta Andermarcher for critical reading of the manuscript.

**Conflicts of Interest:** This work was financially supported by Valbiotis (France). Florian Le Joubioux, Vivien Chavanelle, Cédric Langhi, Arnaud Michaux, Yolanda Otero, and Pascal Sirvent are Valbiotis employees, whose CEO is Sébastien Peltier.

## References

1. Abarca-Gómez, L.; Abdeen, Z.A.; Hamid, Z.A.; Abu-Rmeileh, N.M.; Acosta-Cazares, B.; Acuin, C.; Adams, R.J.; Aekplakorn, W.; Afsana, K.; Aguilar-Salinas, C.A.; et al. Worldwide Trends in Body-Mass Index, Underweight, Overweight, and Obesity from 1975 to 2016: A Pooled Analysis of 2416 Population-Based Measurement Studies in 128·9 Million Children, Adolescents, and Adults. *Lancet* **2017**, *390*, 2627–2642. [https://doi.org/10.1016/S0140-6736\(17\)32129-3](https://doi.org/10.1016/S0140-6736(17)32129-3).
2. Wilson, P.W.F.; D’Agostino, R.B.; Parise, H.; Sullivan, L.; Meigs, J.B. Metabolic Syndrome as a Precursor of Cardiovascular Disease and Type 2 Diabetes Mellitus. *Circulation* **2005**, *112*, 3066–3072. <https://doi.org/10.1161/CIRCULATIONAHA.105.539528>.

3. Lonardo, A.; Nascimbeni, F.; Targher, G.; Bernardi, M.; Bonino, F.; Bugianesi, E.; Casini, A.; Gastaldelli, A.; Marchesini, G.; Marra, F.; et al. AISF Position Paper on Nonalcoholic Fatty Liver Disease (NAFLD): Updates and Future Directions. *Dig. Liver Dis.* **2017**, *49*, 471–483. <https://doi.org/10.1016/j.dld.2017.01.147>.
4. Lustig, R.H.; Collier, D.; Kassotis, C.; Roepke, T.A.; Kim, M.J.; Blanc, E.; Barouki, R.; Bansal, A.; Cave, M.C.; Chatterjee, S.; et al. Obesity I: Overview and Molecular and Biochemical Mechanisms. *Biochem. Pharmacol.* **2022**, *199*, 115012. <https://doi.org/10.1016/j.bcp.2022.115012>.
5. Abdelaal, M.; le Roux, C.W.; Docherty, N.G. Morbidity and Mortality Associated with Obesity. *Ann. Transl. Med.* **2017**, *5*, 161–161. <https://doi.org/10.21037/atm.2017.03.107>.
6. Centers for Disease Control and Prevention., National Diabetes Statistics Report, 2022: Atlanta. Available online: <https://www.cdc.gov/diabetes/library/features/diabetes-stat-report.html> (accessed on 15 May 2022).
7. Roden, M.; Shulman, G.I. The Integrative Biology of Type 2 Diabetes. *Nature* **2019**, *576*, 51–60. <https://doi.org/10.1038/s41586-019-1797-8>.
8. Faulkner, M.S.; Michaliszyn, S.F. Exercise Adherence in Hispanic Adolescents with Obesity or Type 2 Diabetes. *J. Pediatr. Nurs.* **2021**, *56*, 7–12. <https://doi.org/10.1016/j.pedn.2020.09.012>.
9. Brown, M.T.; Bussell, J.K. Medication Adherence: WHO Cares? *Mayo Clin. Proc.* **2011**, *86*, 304–314. <https://doi.org/10.4065/mcp.2010.0575>.
10. Kelly, M.P.; Barker, M. Why Is Changing Health-Related Behaviour so Difficult? *Public Health* **2016**, *136*, 109–116. <https://doi.org/10.1016/j.puhe.2016.03.030>.
11. Lonardo, A.; Nascimbeni, F.; Mantovani, A.; Targher, G. Hypertension, Diabetes, Atherosclerosis and NASH: Cause or Consequence? *J. Hepatol.* **2018**, *68*, 335–352. <https://doi.org/10.1016/j.jhep.2017.09.021>.
12. An, Y.; Li, Y.; Bian, N.; Ding, X.; Chang, X.; Liu, J.; Wang, G. Different Interactive Effects of Metformin and Acarbose With Dietary Macronutrient Intakes on Patients With Type 2 Diabetes Mellitus: Novel Findings From the MARCH Randomized Trial in China. *Front. Nutr.* **2022**, *9*, 861750. <https://doi.org/10.3389/fnut.2022.861750>.
13. Ku, E.J.; Lee, D.-H.; Jeon, H.J.; Oh, T.K. Effectiveness and Safety of Empagliflozin-Based Quadruple Therapy Compared with Insulin Glargine-Based Therapy in Patients with Inadequately Controlled Type 2 Diabetes: An Observational Study in Clinical Practice. *Diabetes Obes. Metab.* **2019**, *21*, 173–177. <https://doi.org/10.1111/dom.13476>.
14. Kaur, N.; Kumar, V.; Nayak, S.K.; Wadhwa, P.; Kaur, P.; Sahu, S.K. Alpha-amylase as Molecular Target for Treatment of Diabetes Mellitus: A Comprehensive Review. *Chem. Biol. Drug Des.* **2021**, *98*, 539–560. <https://doi.org/10.1111/cbdd.13909>.
15. Liu, T.-T.; Liu, X.-T.; Chen, Q.-X.; Shi, Y. Lipase Inhibitors for Obesity: A Review. *Biomed. Pharmacother.* **2020**, *128*, 110314. <https://doi.org/10.1016/j.biopha.2020.110314>.
16. Hossain, U.; Das, A.K.; Ghosh, S.; Sil, P.C. An Overview on the Role of Bioactive  $\alpha$ -Glucosidase Inhibitors in Ameliorating Diabetic Complications. *Food Chem. Toxicol.* **2020**, *145*, 111738. <https://doi.org/10.1016/j.fct.2020.111738>.
17. Holst, J.J.; Gribble, F.; Horowitz, M.; Rayner, C.K. Roles of the Gut in Glucose Homeostasis. *Diabetes Care* **2016**, *39*, 884–892. <https://doi.org/10.2337/dc16-0351>.
18. Nichols, B.L.; Avery, S.; Sen, P.; Swallow, D.M.; Hahn, D.; Sterchi, E. The Maltase-Glucoamylase Gene: Common Ancestry to Sucrase-Isomaltase with Complementary Starch Digestion Activities. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 1432–1437. <https://doi.org/10.1073/pnas.0237170100>.
19. Elferink, H.; Bruekers, J.P.J.; Veeneman, G.H.; Boltje, T.J. A Comprehensive Overview of Substrate Specificity of Glycoside Hydrolases and Transporters in the Small Intestine: “A Gut Feeling.” *Cell. Mol. Life Sci.* **2020**, *77*, 4799–4826. <https://doi.org/10.1007/s00018-020-03564-1>.
20. Thilavech, T.; Adisakwattana, S. Cyanidin-3-Rutinoside Acts as a Natural Inhibitor of Intestinal Lipid Digestion and Absorption. *BMC Complement. Altern. Med.* **2019**, *19*, 242. <https://doi.org/10.1186/s12906-019-2664-8>.
21. Iqbal, J.; Hussain, M.M. Intestinal Lipid Absorption. *Am. J. Physiol. Endocrinol. Metab.* **2009**, *296*, E1183–E1194. <https://doi.org/10.1152/ajpendo.90899.2008>.
22. Chaudhury, A.; Duvoor, C.; Reddy Dendi, V.S.; Kraleti, S.; Chada, A.; Ravilla, R.; Marco, A.; Shekhawat, N.S.; Montales, M.T.; Kuriakose, K.; et al. Clinical Review of Antidiabetic Drugs: Implications for Type 2 Diabetes Mellitus Management. *Front. Endocrinol.* **2017**, *8*. <https://doi.org/10.3389/fendo.2017.00006>.
23. You, Q.; Chen, F.; Wang, X.; Luo, P.G.; Jiang, Y. Inhibitory Effects of Muscadine Anthocyanins on  $\alpha$ -Glucosidase and Pancreatic Lipase Activities. *J. Agric. Food Chem.* **2011**, *59*, 9506–9511. <https://doi.org/10.1021/jf201452v>.
24. Peltier, S.; Sirvent, P.; Maugard, T. Composition Containing a Mixture of Plant Extracts or a Mixture of Molecules Contained in Said Plants, and Use for Controlling Glucose and/or Lipid Metabolism. WO 2016062958 A1, 2016, 2016. Available online: <https://patents.google.com/patent/WO2016062958A1/en> (accessed on 19 December 2022).
25. Frutos, M.J.; Ruiz-Cano, D.; Valero-Cases, E.; Zamora, S.; Pérez-Llamas, F. Artichoke (*Cynara scolymus* L.). In *Nonvitamin and Nonmineral Nutritional Supplements*; Elsevier: Amsterdam, The Netherlands, 2019; pp. 135–138 ISBN 978-0-12-812491-8.
26. Lattanzio, V.; Kroon, P.A.; Linsalata, V.; Cardinali, A. Globe Artichoke: A Functional Food and Source of Nutraceutical Ingredients. *J. Funct. Foods* **2009**, *1*, 131–144. <https://doi.org/10.1016/j.jff.2009.01.002>.
27. Rufino-Palomares, E.E.; Pérez-Jiménez, A.; García-Salguero, L.; Mokhtari, K.; Reyes-Zurita, F.J.; Peragón-Sánchez, J.; Lupiáñez, J.A. Nutraceutical Role of Polyphenols and Triterpenes Present in the Extracts of Fruits and Leaves of *Olea Europaea* as

- Antioxidants, Anti-Infectives and Anticancer Agents on Healthy Growth. *Molecules* **2022**, *27*, 2341. <https://doi.org/10.3390/molecules27072341>.
28. Brasanac-Vukanovic, S.; Mutic, J.; Stankovic, D.; Arsic, I.; Blagojevic, N.; Vukasinovic-Pesic, V.; Tadic, V. Wild Bilberry (*Vaccinium Myrtillus* L., Ericaceae) from Montenegro as a Source of Antioxidants for Use in the Production of Nutraceuticals. *Molecules* **2018**, *23*, 1864. <https://doi.org/10.3390/molecules23081864>.
29. Chavanelle, V.; Otero, Y.F.; Le Joubioux, F.; Ripoché, D.; Bargetto, M.; Vluggens, A.; Montaurier, C.; Pickering, G.; Ducheix, G.; Dubray, C.; et al. Effects of Totum-63 on Glucose Homeostasis and Postprandial Glycemia: A Translational Study. *Am. J. Physiol. Endocrinol. Metab.* **2021**, *320*, E1119–E1137. <https://doi.org/10.1152/ajpendo.00629.2020>.
30. Jerabek-Willemsen, M.; André, T.; Wanner, R.; Roth, H.M.; Dühr, S.; Baaske, P.; Breitsprecher, D. MicroScale Thermophoresis: Interaction Analysis and Beyond. *J. Mol. Struct.* **2014**, *1077*, 101–113. <https://doi.org/10.1016/j.molstruc.2014.03.009>.
31. Chiasson, J.-L.; Josse, R.G.; Gomis, R.; Hanefeld, M.; Karasik, A.; Laakso, M. Acarbose for Prevention of Type 2 Diabetes Mellitus: The STOP-NIDDM Randomised Trial. *Lancet* **2002**, *359*, 2072–2077. [https://doi.org/10.1016/S0140-6736\(02\)08905-5](https://doi.org/10.1016/S0140-6736(02)08905-5).
32. Chelladurai, G.R.M.; Chinnachamy, C. Alpha Amylase and Alpha Glucosidase Inhibitory Effects of Aqueous Stem Extract of *Salacia Oblonga* and Its GC-MS Analysis. *Braz. J. Pharm. Sci.* **2018**, *54*. <https://doi.org/10.1590/s2175-97902018000117151>.
33. Sung, Y.-Y.; Kim, S.-H.; Yoo, B.W.; Kim, H.K. The Nutritional Composition and Anti-Obesity Effects of an Herbal Mixed Extract Containing *Allium Fistulosum* and *Viola Mandshurica* in High-Fat-Diet-Induced Obese Mice. *BMC Complement. Altern. Med.* **2015**, *15*, 370. <https://doi.org/10.1186/s12906-015-0875-1>.
34. Adamska-Patruno, E.; Billing-Marczak, K.; Orłowski, M.; Gorska, M.; Krotkiewski, M.; Kretowski, A. A Synergistic Formulation of Plant Extracts Decreases Postprandial Glucose and Insulin Peaks: Results from Two Randomized, Controlled, Cross-Over Studies Using Real-World Meals. *Nutrients* **2018**, *10*, 956. <https://doi.org/10.3390/nu10080956>.
35. Williamson, E. Synergy and Other Interactions in Phytomedicines. *Phytomedicine* **2001**, *8*, 401–409. <https://doi.org/10.1078/0944-7113-00060>.
36. Wang, S.; Li, Y.; Huang, D.; Chen, S.; Xia, Y.; Zhu, S. The Inhibitory Mechanism of Chlorogenic Acid and Its Acylated Derivatives on  $\alpha$ -Amylase and  $\alpha$ -Glucosidase. *Food Chem.* **2022**, *372*, 131334. <https://doi.org/10.1016/j.foodchem.2021.131334>.
37. Proença, C.; Freitas, M.; Ribeiro, D.; Oliveira, E.F.T.; Sousa, J.L.C.; Tomé, S.M.; Ramos, M.J.; Silva, A.M.S.; Fernandes, P.A.; Fernandes, E.  $\alpha$ -Glucosidase Inhibition by Flavonoids: An in Vitro and in Silico Structure–Activity Relationship Study. *J. Enzym. Inhib. Med. Chem.* **2017**, *32*, 1216–1228. <https://doi.org/10.1080/14756366.2017.1368503>.
38. Yan, J.; Zhang, G.; Pan, J.; Wang, Y.  $\alpha$ -Glucosidase Inhibition by Luteolin: Kinetics, Interaction and Molecular Docking. *Int. J. Biol. Macromol.* **2014**, *64*, 213–223. <https://doi.org/10.1016/j.ijbiomac.2013.12.007>.
39. Hadrich, F.; Bouallagui, Z.; Junkyu, H.; Isoda, H.; Sayadi, S. The  $\alpha$ -Glucosidase and  $\alpha$ -Amylase Enzyme Inhibitory of Hydroxytyrosol and Oleuropein. *J. Oleo Sci.* **2015**, *64*, 835–843. <https://doi.org/10.5650/jos.ess15026>.
40. Dekdouk, N.; Malafronte, N.; Russo, D.; Faraone, I.; De Tommasi, N.; Ameddah, S.; Severino, L.; Milella, L. Phenolic Compounds from *Olea Europaea* L. Possess Antioxidant Activity and Inhibit Carbohydrate Metabolizing Enzymes In Vitro. *Evid. Based Complement. Altern. Med.* **2015**, *2015*, 1–9. <https://doi.org/10.1155/2015/684925>.
41. Kumar, S.; Sharma, S.; Vasudeva, N. Screening of Antidiabetic and Antihyperlipidemic Potential of Oil from *Piper Longum* and Piperine with Their Possible Mechanism. *Expert Opin. Pharmacother.* **2013**, *14*, 1723–1736. <https://doi.org/10.1517/14656566.2013.815725>.
42. Tolmie, M.; Bester, M.J.; Apostolides, Z. Inhibition of  $\alpha$ -glucosidase and  $\alpha$ -amylase by Herbal Compounds for the Treatment of Type 2 Diabetes: A Validation of in Silico Reverse Docking with in Vitro Enzyme Assays. *J. Diabetes* **2021**, *13*, 779–791. <https://doi.org/10.1111/1753-0407.13163>.
43. Guo, T.; Wu, S.; Guo, S.; Bai, L.; Liu, Q.; Bai, N. Synthesis and Evaluation of a Series of Oleanolic Acid Saponins as  $\alpha$ -Glucosidase and  $\alpha$ -Amylase Inhibitors: Oleanolic Acid Saponins as  $\alpha$ -Glucosidase Inhibitors. *Arch. Pharm. Chem. Life Sci.* **2015**, *348*, 615–628. <https://doi.org/10.1002/ardp.201500179>.
44. Wu, H.-B.; Liu, T.-T.; Wang, W.-S.; Feng, J.-C.; Tian, H.-M. Oleanane-Type Saponins from the Roots of *Ligulariopsis Shichuana* and Their  $\alpha$ -Glucosidase Inhibitory Activities. *Molecules* **2017**, *22*, 1981. <https://doi.org/10.3390/molecules22111981>.
45. Chen, J.; Wu, S.; Zhang, Q.; Yin, Z.; Zhang, L.  $\alpha$ -Glucosidase Inhibitory Effect of Anthocyanins from *Cinnamomum Camphora* Fruit: Inhibition Kinetics and Mechanistic Insights through in Vitro and in Silico Studies. *Int. J. Biol. Macromol.* **2020**, *143*, 696–703. <https://doi.org/10.1016/j.ijbiomac.2019.09.091>.
46. Zhang, H.; Wang, G.; Beta, T.; Dong, J. Inhibitory Properties of Aqueous Ethanol Extracts of Propolis on  $\alpha$ -Glucosidase. *Evid. Based Complement. Altern. Med.* **2015**, *2015*, 1–7. <https://doi.org/10.1155/2015/587383>.
47. Zeng, L.; Zhang, G.; Lin, S.; Gong, D. Inhibitory Mechanism of Apigenin on  $\alpha$ -Glucosidase and Synergy Analysis of Flavonoids. *J. Agric. Food Chem.* **2016**, *64*, 6939–6949. <https://doi.org/10.1021/acs.jafc.6b02314>.
48. Mwakalukwa, R.; Amen, Y.; Nagata, M.; Shimizu, K. Postprandial Hyperglycemia Lowering Effect of the Isolated Compounds from Olive Mill Wastes—An Inhibitory Activity and Kinetics Studies on  $\alpha$ -Glucosidase and  $\alpha$ -Amylase Enzymes. *ACS Omega* **2020**, *5*, 20070–20079. <https://doi.org/10.1021/acsomega.0c01622>.
49. Deng, X.-Y.; Ke, J.-J.; Zheng, Y.-Y.; Li, D.-L.; Zhang, K.; Zheng, X.; Wu, J.-Y.; Xiong, Z.; Wu, P.-P.; Xu, X.-T. Synthesis and Bioactivities Evaluation of Oleanolic Acid Oxime Ester Derivatives as  $\alpha$ -Glucosidase and  $\alpha$ -Amylase Inhibitors. *J. Enzym. Inhib. Med. Chem.* **2022**, *37*, 451–461. <https://doi.org/10.1080/14756366.2021.2018682>.

50. Feunaing, R.T.; Tamfu, A.N.; Gbaweng, A.J.Y.; Mekontso Magnibou, L.; Ntchapda, F.; Henoumont, C.; Laurent, S.; Talla, E.; Dinica, R.M. In Vitro Evaluation of  $\alpha$ -Amylase and  $\alpha$ -Glucosidase Inhibition of 2,3-Epoxyprocyanidin C1 and Other Constituents from *Pterocarpus Erinaceus* Poir. *Molecules* **2022**, *28*, 126. <https://doi.org/10.3390/molecules28010126>.
51. Olennikov, D.N.; Chirikova, N.K.; Kashchenko, N.I.; Nikolaev, V.M.; Kim, S.-W.; Vennos, C. Bioactive Phenolics of the Genus *Artemisia* (Asteraceae): HPLC-DAD-ESI-TQ-MS/MS Profile of the Siberian Species and Their Inhibitory Potential Against  $\alpha$ -Amylase and  $\alpha$ -Glucosidase. *Front. Pharmacol.* **2018**, *9*, 756. <https://doi.org/10.3389/fphar.2018.00756>.
52. Lim, J.; Ferruzzi, M.G.; Hamaker, B.R. Structural Requirements of Flavonoids for the Selective Inhibition of  $\alpha$ -Amylase versus  $\alpha$ -Glucosidase. *Food Chem.* **2022**, *370*, 130981. <https://doi.org/10.1016/j.foodchem.2021.130981>.
53. Miao, J.; Liu, J.; Gao, X.; Lu, F.; Yang, X. Effects of Different Drying Methods on Chemical Compositions, Antioxidant Activity and Anti- $\alpha$ -Glucosidase Activity of *Coreopsis Tinctoria* Flower Tea. *Heliyon* **2022**, *8*, e11784. <https://doi.org/10.1016/j.heliyon.2022.e11784>.
54. Islam, Md.N.; Ishita, I.J.; Jung, H.A.; Choi, J.S. Vicenin 2 Isolated from *Artemisia Capillaris* Exhibited Potent Anti-Glycation Properties. *Food Chem. Toxicol.* **2014**, *69*, 55–62. <https://doi.org/10.1016/j.fct.2014.03.042>.
55. Zheng, Y.; Yang, W.; Sun, W.; Chen, S.; Liu, D.; Kong, X.; Tian, J.; Ye, X. Inhibition of Porcine Pancreatic  $\alpha$ -Amylase Activity by Chlorogenic Acid. *J. Funct. Foods* **2020**, *64*, 103587. <https://doi.org/10.1016/j.jff.2019.103587>.
56. Tadera, K.; Minami, Y.; Takamatsu, K.; Matsuoka, T. Inhibition of .ALPHA.-Glucosidase and .ALPHA.-Amylase by Flavonoids. *J. Nutr. Sci. Vitam.* **2006**, *52*, 149–153. <https://doi.org/10.3177/jnsv.52.149>.
57. Seyedan, A.; Alshawsh, M.A.; Alshagga, M.A.; Koosha, S.; Mohamed, Z. Medicinal Plants and Their Inhibitory Activities against Pancreatic Lipase: A Review. *Evid. Based Complement. Altern. Med.* **2015**, *2015*, 1–13. <https://doi.org/10.1155/2015/973143>.
58. Ediger, T.R.; Erdman, S.H. Maldigestion and Malabsorption. In *Pediatric Gastrointestinal and Liver Disease*; Elsevier: Amsterdam, The Netherlands, 2021; pp. 321–338.e5; ISBN 978-0-323-67293-1.
59. Kim, K.S.; Kim, S.K.; Sung, K.M.; Cho, Y.W.; Park, S.W. Management of Type 2 Diabetes Mellitus in Older Adults. *Diabetes Metab. J.* **2012**, *36*, 336. <https://doi.org/10.4093/dmj.2012.36.5.336>.
60. Proença, C.; Freitas, M.; Ribeiro, D.; Tomé, S.M.; Oliveira, E.F.T.; Viegas, M.F.; Araújo, A.N.; Ramos, M.J.; Silva, A.M.S.; Fernandes, P.A.; et al. Evaluation of a Flavonoids Library for Inhibition of Pancreatic  $\alpha$ -Amylase towards a Structure–Activity Relationship. *J. Enzym. Inhib. Med. Chem.* **2019**, *34*, 577–588. <https://doi.org/10.1080/14756366.2018.1558221>.
61. Sui, X.; Zhang, Y.; Zhou, W. In Vitro and in Silico Studies of the Inhibition Activity of Anthocyanins against Porcine Pancreatic  $\alpha$ -Amylase. *J. Funct. Foods* **2016**, *21*, 50–57. <https://doi.org/10.1016/j.jff.2015.11.042>.
62. Yang, X.-W.; Huang, M.-Z.; Jin, Y.-S.; Sun, L.-N.; Song, Y.; Chen, H.-S. Phenolics from *Bidens Bipinnata* and Their Amylase Inhibitory Properties. *Fitoterapia* **2012**, *83*, 1169–1175. <https://doi.org/10.1016/j.fitote.2012.07.005>.
63. Li, M.-M.; Chen, Y.-T.; Ruan, J.-C.; Wang, W.-J.; Chen, J.-G.; Zhang, Q.-F. Structure-Activity Relationship of Dietary Flavonoids on Pancreatic Lipase. *Curr. Res. Food Sci.* **2023**, *6*, 100424. <https://doi.org/10.1016/j.crf.2022.100424>.
64. Yu, H.; Dong, S.; Wang, L.; Liu, Y. The Effect of Triterpenoid Saponins on Pancreatic Lipase in Vitro : Activity, Conformation, Kinetics, Thermodynamics and Morphology. *Biochem. Eng. J.* **2017**, *125*, 1–9. <https://doi.org/10.1016/j.bej.2017.05.010>.
65. Wang, Y.; Chen, L.; Liu, H.; Xie, J.; Yin, W.; Xu, Z.; Ma, H.; Wu, W.; Zheng, M.; Liu, M.; et al. Characterization of the Synergistic Inhibitory Effect of Cyanidin-3-O-Glucoside and Catechin on Pancreatic Lipase. *Food Chem.* **2023**, *404*, 134672. <https://doi.org/10.1016/j.foodchem.2022.134672>.
66. Ahmad, I.; Syakfanaya, A.M.; Azminah, A.; Saputri, F.C.; Mun'im, A. Optimization of Betaine-Sorbitol Natural Deep Eutectic Solvent-Based Ultrasound-Assisted Extraction and Pancreatic Lipase Inhibitory Activity of Chlorogenic Acid and Caffeine Content from Robusta Green Coffee Beans. *Heliyon* **2021**, *7*, e07702. <https://doi.org/10.1016/j.heliyon.2021.e07702>.
67. Hu, B.; Cui, F.; Yin, F.; Zeng, X.; Sun, Y.; Li, Y. Caffeoylquinic Acids Competitively Inhibit Pancreatic Lipase through Binding to the Catalytic Triad. *Int. J. Biol. Macromol.* **2015**, *80*, 529–535. <https://doi.org/10.1016/j.ijbiomac.2015.07.031>.
68. Ma, L.-J.; Hou, X.-D.; Qin, X.-Y.; He, R.-J.; Yu, H.-N.; Hu, Q.; Guan, X.-Q.; Jia, S.-N.; Hou, J.; Lei, T.; et al. Discovery of Human Pancreatic Lipase Inhibitors from Root of *Rhodiola Crenulata* via Integrating Bioactivity-Guided Fractionation, Chemical Profiling and Biochemical Assay. *J. Pharm. Anal.* **2022**, *12*, 683–691. <https://doi.org/10.1016/j.jpha.2022.04.002>.
69. Şöhretoğlu, D.; Sari, S.; Özel, A.; Barut, B.  $\alpha$ -Glucosidase Inhibitory Effect of *Potentilla Astracantha* and Some Isoflavones: Inhibition Kinetics and Mechanistic Insights through in Vitro and in Silico Studies. *Int. J. Biol. Macromol.* **2017**, *105*, 1062–1070. <https://doi.org/10.1016/j.ijbiomac.2017.07.132>.
70. Chen, Y.; Geng, S.; Liu, B. Three Common Caffeoylquinic Acids as Potential Hypoglycemic Nutraceuticals: Evaluation of  $\alpha$ -glucosidase Inhibitory Activity and Glucose Consumption in HepG2 Cells. *J. Food Biochem.* **2020**, *44*. <https://doi.org/10.1111/jfbc.13361>.
71. Dong, Y.; Huang, H.; Zhao, M.; Sun-Waterhouse, D.; Lin, L.; Xiao, C. Mechanisms Underlying the Xanthine Oxidase Inhibitory Effects of Dietary Flavonoids Galangin and Pinobanksin. *J. Funct. Foods* **2016**, *24*, 26–36. <https://doi.org/10.1016/j.jff.2016.03.021>.
72. Yan, J.; Zhang, G.; Hu, Y.; Ma, Y. Effect of Luteolin on Xanthine Oxidase: Inhibition Kinetics and Interaction Mechanism Merging with Docking Simulation. *Food Chem.* **2013**, *141*, 3766–3773. <https://doi.org/10.1016/j.foodchem.2013.06.092>.
73. Segel, I.H. *Enzyme Kinetics: Behavior and Analysis of Rapid Equilibrium and Steady-State Enzyme Systems*; Wiley: Hoboken, NJ, Canada, 1993.

74. Şöhretoğlu, D.; Sari, S.; Barut, B.; Özel, A. Discovery of Potent  $\alpha$ -Glucosidase Inhibitor Flavonols: Insights into Mechanism of Action through Inhibition Kinetics and Docking Simulations. *Bioorganic Chem.* **2018**, *79*, 257–264. <https://doi.org/10.1016/j.bioorg.2018.05.010>.
75. Wua, J.; Wang, H.-M.; Li, J.; Men, X.-L. [The research applications of db/db mouse]. *Sheng Li Ke Xue Jin Zhan* **2013**, *44*, 12–18.
76. Lee, S.M. The Effect of Chronic  $\alpha$ -Glycosidase Inhibition on Diabetic Nephropathy in the *Db/Db* Mouse. *Diabetes* **1982**, *31*, 249–254. <https://doi.org/10.2337/diab.31.3.249>.
77. Lee, S.M.; Bustamante, S.; Flores, C.; Bezerra, J.; Goda, T.; Koldovský, O. Chronic Effects of an Alpha-Glucosidase Inhibitor (Bay o 1248) on Intestinal Disaccharidase Activity in Normal and Diabetic Mice. *J. Pharm. Exp.* **1987**, *240*, 132.
78. Scheen, A.J. Clinical Efficacy of Acarbose in Diabetes Mellitus: A Critical Review of Controlled Trials. *Diabetes Metab.* **1998**, *24*, 311–320.
79. Zhang, F.; Xu, S.; Tang, L.; Pan, X.; Tong, N. Acarbose With Comparable Glucose-Lowering but Superior Weight-Loss Efficacy to Dipeptidyl Peptidase-4 Inhibitors: A Systematic Review and Network Meta-Analysis of Randomized Controlled Trials. *Front. Endocrinol.* **2020**, *11*, 288. <https://doi.org/10.3389/fendo.2020.00288>.
80. Padwal, R.S.; Rucker, D.; Li, S.K.; Curioni, C.; Lau, D.C. Long-Term Pharmacotherapy for Obesity and Overweight. *Cochrane Database Syst. Rev.* **2003**. <https://doi.org/10.1002/14651858.CD004094.pub2>.
81. Yeh, W.-J.; Hsia, S.-M.; Lee, W.-H.; Wu, C.-H. Polyphenols with Antiglycation Activity and Mechanisms of Action: A Review of Recent Findings. *J. Food Drug Anal.* **2017**, *25*, 84–92. <https://doi.org/10.1016/j.jfda.2016.10.017>.
82. Wu, C.-H.; Yen, G.-C. Inhibitory Effect of Naturally Occurring Flavonoids on the Formation of Advanced Glycation Endproducts. *J. Agric. Food Chem.* **2005**, *53*, 3167–3173. <https://doi.org/10.1021/jf048550u>.
83. Wilding, J.P.H. The Importance of Weight Management in Type 2 Diabetes Mellitus. *Int. J. Clin. Pract.* **2014**, *68*, 682–691. <https://doi.org/10.1111/ijcp.12384>.
84. Davies, M.J.; Aroda, V.R.; Collins, B.S.; Gabbay, R.A.; Green, J.; Maruthur, N.M.; Rosas, S.E.; Del Prato, S.; Mathieu, C.; Mingrone, G.; et al. Management of Hyperglycemia in Type 2 Diabetes, 2022. A Consensus Report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* **2022**, *45*, 2753–2786. <https://doi.org/10.2337/dci22-0034>.
85. Singleton, V.L.; Rossi, J.A. Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. *Am. J. Enol. Vitic.* **1965**, *16*, 144.
86. Bischoff, H. The mechanism of alpha-glucosidase inhibition in the management of diabetes. *Clin. Investig. Med. Med. Clin. Exp.* **1995**, *18*, 303–311.

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.