



**HAL**  
open science

**Methanomethylophilus alvi gen. nov., sp. nov., a Novel Hydrogenotrophic Methyl-Reducing Methanogenic Archaea of the Order Methanomassiliicoccales Isolated from the Human Gut and Proposal of the Novel Family Methanomethylophilaceae fam. nov.**

Guillaume Borrel, Khaled Fadhlou, Wajdi Ben Hania, Nadia Gaci, Gérard Péhau-arnaudet, Prem Prashant Chaudhary, Pascal Vandekerckove, Nathalie Ballet, Monique Alric, Paul William O'toole, et al.

► **To cite this version:**

Guillaume Borrel, Khaled Fadhlou, Wajdi Ben Hania, Nadia Gaci, Gérard Péhau-arnaudet, et al.. Methanomethylophilus alvi gen. nov., sp. nov., a Novel Hydrogenotrophic Methyl-Reducing Methanogenic Archaea of the Order Methanomassiliicoccales Isolated from the Human Gut and Proposal of the Novel Family Methanomethylophilaceae fam. nov.. Microorganisms, 2023, 11 (11), pp.2794. 10.3390/microorganisms11112794 . hal-04306127

**HAL Id: hal-04306127**

**<https://uca.hal.science/hal-04306127>**

Submitted on 3 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## Article

# *Methanomethylophilus alvi* gen. nov., sp. nov., a Novel Hydrogenotrophic Methyl-Reducing Methanogenic Archaea of the Order *Methanomassiliicoccales* Isolated from the Human Gut and Proposal of the Novel Family *Methanomethylophilaceae* fam. nov.

Guillaume Borrel <sup>1,\*</sup>, Khaled Fadhlouli <sup>2,3,4,†</sup>, Wajdi Ben Hania <sup>2,5</sup>, Nadia Gaci <sup>5</sup>, Gérard Pehau-Arnaudet <sup>6</sup>, Prem Prashant Chaudhary <sup>5,7</sup>, Pascal Vandekerckove <sup>8</sup>, Nathalie Ballet <sup>8</sup>, Monique Alric <sup>5</sup>, Paul William O'Toole <sup>9</sup>, Marie-Laure Fardeau <sup>2</sup>, Bernard Ollivier <sup>2</sup> and Jean-François Brugère <sup>5</sup>



**Citation:** Borrel, G.; Fadhlouli, K.; Ben Hania, W.; Gaci, N.; Pehau-Arnaudet, G.; Chaudhary, P.P.; Vandekerckove, P.; Ballet, N.; Alric, M.; O'Toole, P.W.; et al.

*Methanomethylophilus alvi* gen. nov., sp. nov., a Novel Hydrogenotrophic Methyl-Reducing Methanogenic Archaea of the Order *Methanomassiliicoccales* Isolated from the Human Gut and Proposal of the Novel Family *Methanomethylophilaceae* fam. nov. *Microorganisms* **2023**, *11*, 2794. <https://doi.org/10.3390/microorganisms11112794>

Academic Editor: Seong-Tshool Hong

Received: 28 October 2023  
Revised: 9 November 2023  
Accepted: 14 November 2023  
Published: 17 November 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

- <sup>1</sup> Institut Pasteur, Université Paris Cité, Evolutionary Biology of the Microbial Cell, 75015 Paris, France
  - <sup>2</sup> Aix Marseille Univ., Université de Toulon, CNRS, IRD, MIO, 13288 Marseille, France; khaled.fadhlouli@uca.fr (K.F.); bernard.ollivier@mio.osupytheas.fr (B.O.)
  - <sup>3</sup> Université Clermont Auvergne, INRA, MEDIS, 63000 Clermont-Ferrand, France
  - <sup>4</sup> Université Clermont Auvergne, CNRS, UMR 6023 CNRS-UCA, Laboratoire Microorganismes: Génome et Environnement LMGE, 63000 Clermont-Ferrand, France
  - <sup>5</sup> Université d'Auvergne, EA CIDAM, 63000 Clermont-Ferrand, France; j-francois.brugere@uca.fr (J.-F.B.)
  - <sup>6</sup> Institut Pasteur, Université Paris Cité, Ultrastructural Bioimaging, 75015 Paris, France
  - <sup>7</sup> Epithelial Therapeutics Unit, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892, USA
  - <sup>8</sup> Lesaffre International, Lesaffre Group, 59700 Marcq-en-Barœul, France
  - <sup>9</sup> APC Microbiome Ireland, University College Cork, T12 K8AF Cork, Ireland
- \* Correspondence: guillaume.borrel@pasteur.fr  
† These authors contributed equally to this work.

**Abstract:** The methanogenic strain Mx-05<sup>T</sup> was isolated from the human fecal microbiome. A phylogenetic analysis based on the 16S rRNA gene and protein marker genes indicated that the strain is affiliated with the order *Methanomassiliicoccales*. It shares 86.9% 16S rRNA gene sequence identity with *Methanomassiliicoccus luminyensis*, the only member of this order previously isolated. The cells of Mx-05<sup>T</sup> were non-motile cocci, with a diameter range of 0.4–0.7 µm. They grew anaerobically and reduced methanol, monomethylamine, dimethylamine, and trimethylamine into methane, using H<sub>2</sub> as an electron donor. H<sub>2</sub>/CO<sub>2</sub>, formate, ethanol, and acetate were not used as energy sources. The growth of Mx-05<sup>T</sup> required an unknown medium factor(s) provided by *Eggerthella lenta* and present in rumen fluid. Mx-05<sup>T</sup> grew between 30 °C and 40 °C (optimum 37 °C), over a pH range of 6.9–8.3 (optimum pH 7.5), and between 0.02 and 0.34 mol.L<sup>-1</sup> NaCl (optimum 0.12 mol.L<sup>-1</sup> NaCl). The genome is 1.67 Mbp with a G+C content of 55.5 mol%. Genome sequence annotation confirmed the absence of the methyl branch of the H<sub>4</sub>MPT Wood–Ljungdahl pathway, as described for other *Methanomassiliicoccales* members. Based on an average nucleotide identity analysis, we propose strain Mx-05<sup>T</sup> as being a novel representative of the order *Methanomassiliicoccales*, within the novel family *Methanomethylophilaceae*, for which the name *Methanomethylophilus alvi* gen. nov., sp. nov. is proposed. The type strain is Mx-05<sup>T</sup> (JCM 31474T).

**Keywords:** archaea; methanogens; human gut microbiota; *Methanomassiliicoccales*

## 1. Introduction

Only one species of the order *Methanomassiliicoccales* [1], *Methanomassiliicoccus luminyensis* B10<sup>T</sup>, has so far been isolated and described [2]. This archaeon was isolated

from human feces and belongs to the family *Methanomassiliicoccaceae* which is predominantly found in soil and sediment environments according to 16S rRNA gene surveys [3,4]. *Methanomassiliicoccus luminyensis* is rarely detected in the gut environment [5,6] but was found to be one of the most widely distributed representatives of the *Methanomassiliicoccales* in a diverse range of environments, indicating a generalist lifestyle [7]. Accordingly, a second strain of *M. luminyensis* (strain CZDD1) was isolated from arsenic-contaminated paddy field soil [8]. A closely related species, ‘*Candidatus Methanomassiliicoccus intestinalis*’, was enriched from a human fecal sample [9] and is more prevalent in human populations than *M. luminyensis* [5,6]. Two additional enrichment cultures of *Methanomassiliicoccaceae* were generated from freshwater sediment [7] and peat soil [10]. A second family-level lineage within the *Methanomassiliicoccales* family was retrieved primarily from samples of the gut environment. Initially, it was variously referred to as “rumen cluster C” [11], “host-associated clade” [12], or “intestinal clade” [13]. In 2012, one of the first enrichment cultures of a representative of this lineage (strain Mx1201) was obtained and its genome was sequenced [14]. The name ‘*Candidatus Methanomethylophilus alvus*’ was proposed for this methanogenic archaeon [14,15] and the name ‘*Candidatus Methanomethylophilaceae*’ [16] was subsequently proposed for the family to which it belongs. Other enrichment cultures of *Methanomassiliicoccales* representatives belonging to ‘*Ca. Methanomethylophilaceae*’ were obtained from fecal and digestive tract samples of different animals including ruminants and termites [4,13,17–19], as well as from anaerobic digester sludge from garbage slurry [1].

*Methanomassiliicoccus luminyensis* B10<sup>T</sup> and other *Methanomassiliicoccales* members characterized from enrichment cultures are hydrogenotrophic methyl-reducing methanogens (reducing methyl compounds into methane using H<sub>2</sub> as the electron donor). Genomic analyses of *Methanomassiliicoccales* suggested different adaptations to the gut environment in ‘*Ca. Methanomethylophilaceae*’ members, such as the presence of specific adhesins and bile acid resistance genes [5,20]. *Methanomassiliicoccales* members display several unusual characteristics, including a split 16S and 23S rRNA gene operon, the absence of known histone-like proteins [21], and a new mode of energy conservation relying on a homologue of the respiratory complex I, which does not transfer electrons to a membrane-soluble electron carrier [19,22,23]. In addition, *Methanomassiliicoccus luminyensis* displays tetraether lipids with a butanetriol or a pentanetriol replacing one of the two glycerols [24,25]. *Methanomassiliicoccales* genomes encode a particular *pyl* system responsible for the biosynthesis of pyrrolysine (Pyl, the 22nd proteogenic amino acid) and its incorporation in methyltransferase enzymes required for methylamine-based methanogenesis [26,27].

The *Methanomassiliicoccales* members have attracted recent investigations because of their potential biotechnological applications, their roles in bioremediation and environmental element cycling, and their health relationships as part of the gut microbiome. For example, the tRNA<sup>Pyl</sup>–PylRS pair of ‘*Ca. M. alvus*’ has attracted growing interest in synthetic biology as it can incorporate several non-canonical amino acids in proteins with high specificity and efficiency [28]. The tRNA<sup>Pyl</sup>–PylRS pair of ‘*Ca. M. alvus*’ is already used to develop applications for human in vivo studies and cancer therapeutics [29]. The environmental relevance is exemplified by the important role played by *M. luminyensis* in the multi-species demethylation process of the toxic dimethylarsinic acid observed in paddy field soil [8]. The addition of *M. luminyensis* CZDD1 to an arsenic-contaminated paddy soil was shown to alleviate rice straighthead disease caused by dimethylarsinic acid [8]. *Methanomassiliicoccales* can also play an important role in animal and human gut microbiomes. Twenty-seven potential archaeal species have been identified in the human gut on the basis of average nucleotide identity (ANI) comparisons between 1167 genomes reconstructed from fecal metagenomes originating from five continents [6]. Of these, 12 belong to the *Methanomassiliicoccales*, 10 of which have the genetic potential to use trimethylamine as an electron acceptor for methanogenesis. Trimethylamine (TMA) is a bacterial metabolite produced from different dietary compounds (e.g., carnitine, lecithin) and is indirectly involved in the development of kidney [30] and cardiovascular [31] diseases, as well as trimethylaminuria [32]. It was, thus, proposed that *Methanomassiliicoccales* could play a

beneficial role in human health by depleting this compound directly from its source [5,16]. Subjects with a higher abundance level of *Methanomassiliicoccales* in their gut have a lower TMA concentration in their feces than people without *Methanomassiliicoccales* [5]. Thus, we have proposed using these archaea [16] or genetically engineered bacteria expressing *Methanomassiliicoccales* genes [33] as probiotics. However, their therapeutic development and evaluation are impaired by the lack of isolated, well-characterized members of the host-associated clade of *Methanomassiliicoccales*. Beyond humans, a recent study on the distribution and abundance of gut methanogens within 250 animal species revealed that 'Ca. Methanomethylophilaceae' is one of the five main lineages of methanogens in this environment and can represent the large majority of the gut archaea in several animals [3].

Here, we report the isolation and formal taxonomic description of strain Mx-05<sup>T</sup> as the second isolated member pertaining to the *Methanomassiliicoccales* order and propose the name *Methanomethylophilus alvi* gen. nov., sp. nov., within the novel family *Methanomethylophilaceae* fam. nov.

## 2. Materials and Methods

### 2.1. Medium of Enrichment, Isolation, and Maintenance

Yeast extract, tryptone and peptone were purchased from Becton Dickinson (Miami, FL, USA). All other chemicals were purchased from Sigma-Aldrich (Steinheim, Germany).

The enrichment culture was performed in a medium containing the following components per liter: KH<sub>2</sub>PO<sub>4</sub> (0.5 g), MgSO<sub>4</sub>·7H<sub>2</sub>O (0.4 g), NaCl (0.4 g), NH<sub>4</sub>Cl (0.4 g), CaCl<sub>2</sub>·2H<sub>2</sub>O (0.05 g), sodium acetate (1 g), cysteine HCl (0.5 g), Na<sub>2</sub>S·9H<sub>2</sub>O (0.15 g), NaHCO<sub>3</sub> (4 g), yeast extract (1 g), resazurin (1 mg), Widdel vitamin solution (20 mL) [34], Widdel trace element solution (1 mL) [34], and tungstate–selenite solution (1 mL) [2], methanol (60 mmol.L<sup>-1</sup> final). The tungstate–selenite solution contained the following components per liter: NaOH (0.4 g), Na<sub>2</sub>SeO<sub>3</sub> (2 mg), and Na<sub>2</sub>WO<sub>4</sub>·5H<sub>2</sub>O (4 mg). Before autoclaving, all chemicals except NaHCO<sub>3</sub>, Na<sub>2</sub>S·9H<sub>2</sub>O, and vitamins were added and the pH of the medium was adjusted to 7.5 using 10 mol.L<sup>-1</sup> KOH. Subsequently, the medium was purged with oxygen-free N<sub>2</sub> gas and transferred into Hungate tubes (5 mL per tube) and 100 mL serum bottles (30 mL per bottle). After autoclaving, methanol, NaHCO<sub>3</sub>, Na<sub>2</sub>S·9H<sub>2</sub>O, and vitamins were added to the medium. The tubes were pressurized by adding H<sub>2</sub> after inoculation (200 kPa).

The roll tube medium corresponded to the enrichment medium supplemented with 2% (*v/v*) of an *Eggerthella* sp. culture filtrate and 2% (*w/v*) agar. The roll tubes were pressurized with H<sub>2</sub>/CO<sub>2</sub> (200 kPa) after inoculation. *Eggerthella lenta* was grown on commercial BHI medium (Sigma Aldrich), supplemented with sterile glucose (20 mM), Na<sub>2</sub>S·9H<sub>2</sub>O (0.4 g.L<sup>-1</sup>), NaHCO<sub>3</sub> (2 g.L<sup>-1</sup>), and Widdel vitamin (20 mL.L<sup>-1</sup>) solutions after autoclaving. This culture was filtered and used when performing the serial dilution of the enrichment culture in solid medium-containing roll tubes.

The axenic culture (strain Mx-05<sup>T</sup>) was checked periodically by monitoring the cell morphology under a bright-field microscope and by confirming the absence of fermentative contaminants when inoculating the modified BHI medium in the absence of H<sub>2</sub> and methanol.

Strain Mx-05<sup>T</sup> was then cultured in a medium containing the following per liter: KH<sub>2</sub>PO<sub>4</sub> (0.5 g), MgSO<sub>4</sub>·7H<sub>2</sub>O (0.4 g), NaCl (5 g), NH<sub>4</sub>Cl (0.4 g), CaCl<sub>2</sub>·2H<sub>2</sub>O (0.05 g), sodium acetate (1 g), yeast extract (2 g), tryptone (2 g), peptone (2 g), casamino acid (2 g), cysteine HCl (0.5 g), Na<sub>2</sub>S·9H<sub>2</sub>O (0.15 g), NaHCO<sub>3</sub> (4 g), yeast extract (1 g), resazurin (1 mg), Widdel vitamin solution (20 mL) [34], Widdel trace element solution (1 mL) [34], tungstate-selenite solution (1 mL), methanol (60 mmol.L<sup>-1</sup> final), and a filtrate of *Eggerthella lenta* DSM 2243<sup>T</sup>-grown culture (10% *v/v*).

### 2.2. Genome Sequencing, Genomic and Phylogenomic Analyses

The genome of strain Mx-05<sup>T</sup> was sequenced on an Illumina HiSeq instrument generating 2 × 150 bp reads. The reads were assembled with the GS De Novo Assembler v11/2015

(Roche) into a single circular chromosome. The genome was annotated with the Prokaryotic Genome Annotation Pipeline [35]. We built a phylogenomic tree of the *Methanomassiliicoccales* family using 41 phylogenetic marker proteins [36] from 54 *Methanomassiliicoccales* representatives, including Mx-05<sup>T</sup>, '*Ca. M. alvus*' Mx1201, *M. luminyensis* B10<sup>T</sup>, and eight genomes derived from previously described enrichment cultures, as well as from 43 genomes reconstructed from metagenomes (MAGs), covering the currently known phylogenetic diversity of the order (at least one genome per candidate genus in GTDB [37]). The tree was rooted with three members of '*Ca. Sysuiplasmatales*', the closest relatives to *Methanomassiliicoccales* family that do not carry *mcrABG* genes [38]. The protein sequences used to build the tree were retrieved with an HMM search (hmmer v3.3.2 [39]), aligned with MAFFT (v7.453) [40], trimmed with BMGE (v2) [41], and concatenated. A maximum likelihood tree was built with IQ-TREE (v2.0.6) [42] using the LG+F+R10 model. For the 16S rRNA gene tree, we extracted the 16S rRNA gene from the Mx-05<sup>T</sup> genome and other genomes selected for the phylogenomic analysis using Metaxa2 (v2.2) [43]. The maximum likelihood tree was built with IQ-TREE using the GTR+F+R4 model.

### 2.3. Microscopy

The cells were observed with an inverted contrasting microscope (Leica DMIRB, Leica, Germany) equipped with a Hamamatsu C13440 camera. For the cryo-electron microscopy (Cryo-EM), 4 µL of each sample were deposited on Quantifoil R2/1 glow discharge, then cryofixed using a Leica EMGP (Leica, Germany). The samples were visualized under a Tecnai F20 electron microscope (ThermoFisher, Waltham, MA, USA) with low-dose conditions and imaged using a Falcon II camera (ThermoFisher, Waltham, MA, USA).

### 2.4. Physiological Characterization

Test substrates for methanogenesis were added to the culture medium after autoclaving. In addition to H<sub>2</sub>/methanol (60 mmol.L<sup>-1</sup>), the tested substrates were trimethylamine (20 mmol.L<sup>-1</sup>), dimethylamine (20 mmol.L<sup>-1</sup>), monomethylamine (20 mmol.L<sup>-1</sup>), dimethyl sulfide (20 mM), CO<sub>2</sub> (20% of the gas phase), formate (20 mmol.L<sup>-1</sup>), ethanol (20 mmol.L<sup>-1</sup>), and acetate (20 mmol.L<sup>-1</sup>). Trimethylamine, dimethylamine, monomethylamine, dimethyl sulfide, and CO<sub>2</sub> were tested as potential electron acceptors in the presence of H<sub>2</sub>. Formate and ethanol were tested as electron donors for methanol reduction in the absence of H<sub>2</sub>. Acetate was tested solely as an energy source in the absence of H<sub>2</sub>.

The temperature range and optimum value for Mx-05<sup>T</sup> were determined at pH 7 and with 0.09 mol.L<sup>-1</sup> of NaCl, from 25 °C to 50 °C, with a 5 °C increment and measured at 37 °C. The pH range and optimum value were determined at 37 °C and with 0.09 mol.L<sup>-1</sup> of NaCl, from pH 5.3 to pH 9.5, with 0.2–0.3 pH steps. The salinity range and optimum value were determined at 37 °C and pH 7, from 0 to 0.86 mol.L<sup>-1</sup> NaCl.


The susceptibility to lysis with sodium dodecyl sulphate (SDS) was tested by adding SDS at a final concentration of 0.01% to a late-logarithmic-phase culture of Mx-05<sup>T</sup> and by following the decrease in OD. The susceptibility to lysis due to osmotic stress was also tested by resuspending cells in distilled water.

## 3. Results and Discussion

### 3.1. Selective Enrichment and Isolation

The inoculum for the enrichment culture was a fecal sample from an elderly subject [14]. The absolute abundances of total bacteria and *Methanomassiliicoccales* from "host-associated" and "free-living" clades were followed by qPCR three days after each transfer, using primers and conditions previously described [5]. Transferring the culture every three days led to the enrichment of '*Ca. M. alvus*' [14], while transfers every 10 days led to the enrichment of '*Ca. M. intestinalis*' [9] from the same initial sample. The latter was enriched up to 45% of the whole microbial community but it was lost after two years of subculturing. The 16S rRNA gene amplicon sequencing (MiSeq) was performed on enrichment cultures of the "host-associated" clade, amended with H<sub>2</sub> as the electron donor and either methanol

or trimethylamine as terminal electron acceptors, following the procedure described by Flemer et al. [44]. The enrichment culture on H<sub>2</sub>/trimethylamine had the highest relative abundance of *Methanomassiliicoccales* (Figure 1).




**Figure 1.** Proportional abundance of bacteria and archaea in enrichment cultures using either H<sub>2</sub>/methanol (a) or H<sub>2</sub>/trimethylamine (b) two years after the beginning of the enrichment culture. Percentage values next to the bacterial names correspond to the sequence identity they display to the closest species.

The main difference in the compositions of the two enrichment cultures was due to the massive out-growth of a species closely related to *Eubacterium limosum* in the presence of methanol, with all other bacteria being present in similar relative abundance levels between the two conditions. *Eubacterium limosum* can use methanol and possibly H<sub>2</sub> as sources of energy, making it a major competitor of the *Methanomassiliicoccales* members in the culture conditions that we used and potentially in the gut microbiome. After additional transfers (every three days for two years) and filtrations through 0.45 µm to deplete *Eubacterium* sp., 'Ca. M. alvus' Mx1201 represented 99% of the cells in the enrichment culture while retaining the presence of one rod-shaped microorganism. The first attempts to obtain the methanogen in axenic culture on a solid medium failed, which we hypothesized might be due to its dependence on bacteria for one or more growth factors. The remaining microbial contaminant was further isolated and was identified as an *Eggerthella* strain sharing more than 97% identity similarity with *Eggerthella lenta*. Thereafter, a filtrate (0.2 µm) from a culture supernatant of *Eggerthella* sp. was added to the roll tube solid medium during the process of the isolation of the methanogen. After one month of incubation at 37 °C, white and round colonies measuring 0.5–1 mm in diameter were obtained, transferred to a fresh liquid medium, and serially diluted. The addition of the culture filtrate of *Eggerthella* sp. was, therefore, essential to obtain the methanogenic strain Mx-05<sup>T</sup> in axenic culture.

### 3.2. Phylogeny and Genome Features

Mx-05<sup>T</sup> is a member of a clade that is composed almost exclusively of organisms from the gut microbiome [5,12] corresponding to taxa of the previously proposed 'Ca. Methanomethylophilaceae' (Figure 2). As expected, Mx-05<sup>T</sup> and 'Ca. M. alvus' Mx1201 are contiguous in the tree. Mx-05<sup>T</sup> is closely related to an MAG obtained from a human gut metagenome (GCA\_006954385) [5] and another taxon obtained from a rumen enrichment culture, 'Ca. Methanomethylophilus sp.' 1R26 (GCF\_001481295) [18]. Similar placements of Mx-05<sup>T</sup> and similar tree topologies for the *Methanomassiliicoccales* and proposed *Methanomethylophilaceae* families were obtained within a 16S rRNA gene tree (Figure S1). Mx-05<sup>T</sup> and *M. luminyensis* B10<sup>T</sup> share 86.9% nucleotide identity similarity on the 16S rRNA and 76.2% amino acid identity similarity on the McrA sequence. They share 66.3% ANI similarity with an alignment of 12% on the whole genome of Mx-05<sup>T</sup>.



**Figure 2.** *Methanomassiliicoccales* maximum likelihood tree built with an LG+F+R10 model from 41 concatenated protein sequences. The tree is rooted on ‘Candidatus Sysuiplasmatales’ [38], with 10,325 positions after trimming. Values on the tree indicate the bootstrap supports > 70%. The tree was formatted with Itol [45]. Isolated species are indicated in bold.

The Mx-05<sup>T</sup> genome is 1.67 Mbp in size with a G+C content of 55.5 mol%. Both the genome size and G+C content are smaller than those of *M. luminyensis* B10<sup>T</sup> (Table 1). More generally, the genome length within the proposed *Methanomethylophilaceae* family (average of 1.57 Mbp) is substantially smaller than that of the *Methanomassiliicoccaceae* (average of 2.12 Mbp, Mann–Whitney  $p = 0.0001$ , Figure 2), suggesting reductive genome evolution during the adaptation to the gut environment. The Mx-05<sup>T</sup> genome codes for 1597 predicted CDS, 45 tRNA genes, one 23S gene, one 16S gene, and two 5S rRNA genes. The rRNA genes do not form a single operon in Mx-05<sup>T</sup>, which is a characteristic shared with other members of the ‘*Ca. Thermoplasmatota*’ phylum [21,46]. Mx-05<sup>T</sup> genome has only 27 single-nucleotide polymorphisms (SNPs), with the genome of ‘*Ca. M. alvus*’ Mx-1201 obtained from the initial enrichment in 2012 (CP004049.1/GCA\_000300255.2) and a 32 bp insertion in the 5’ end of a hypothetical pyrrolysine-containing protein [21]. Twenty of the SNPs correspond to synonymous mutations, three are in non-coding regions, one is in the CRISPR repeat array, and one led to the deletion of the 5’ end of a gene coding for a GT2/GT4 glycosyltransferase. Accordingly, genomic features previously described for ‘*Ca. M. alvus*’ Mx1201 are shared by Mx-05<sup>T</sup> [12,27,47]. These include the lack of genes coding for the methyl branch of the H<sub>4</sub>MPT Wood–Ljungdahl pathway, which is involved in (i) the reduction of CO<sub>2</sub> into a methyl group during hydrogenotrophic CO<sub>2</sub>-reducing methanogenesis (H<sub>2</sub> + CO<sub>2</sub> → CH<sub>4</sub> + H<sub>2</sub>O) and (ii) the oxidation of a methyl group into CO<sub>2</sub> during methylotrophic methanogenesis (e.g., 4 CH<sub>3</sub>OH → 3CH<sub>3</sub> + CO<sub>2</sub>). Mx-05<sup>T</sup> codes for McrABG involved in methane production, for several methyltransferases used to transfer methyl groups from various substrates to coenzyme M (i.e., methanol

(MtaBC), monomethylamine (MtmBC), dimethylamine (MtbBC), trimethylamine (MttBC), and methylated thiol compounds (MtsAB)), as well as [NiFe]-hydrogenase (MvhAG) to use H<sub>2</sub> as an electron donor. The Mx-05<sup>T</sup> genome harbors genes for six predicted membrane-bound adhesin-like proteins with multiple Flg\_New repeats, including two (AYQ55217.1 and AYQ55598.1) that are longer than 1000 amino acids. Flg\_New repeats have been identified in the Pil3 protein of *Streptococcus gallolyticus*, playing an essential role in its adhesion to the human gut mucosa [48]. Proteins with Flg\_New repeats were also identified in distantly related gut *Methanimicrococcus* (*Methanosarcinales*) members [47], as well as in viruses of gut *Methanomassiliicoccales* members [49]. These proteins likely play an important role in the gut colonization of Mx-05<sup>T</sup>. Other characteristics are the absence of characterized histone or other nucleotide-associated proteins (NAPs) involved in the formation of chromatin. A large diversity of genes coding for these NAPs have been reported in other members of the ‘*Ca. Thermoplasmatota*’ phylum, including Alba, MC1, and HU [50]. This highlights a potential loss of the histone gene in the last common ancestor of the ‘*Ca. Thermoplasmatota*’ phylum, which was then replaced by various enzymes. *M. luminyensis* codes for an Alba protein but it is not expressed at a detectable level [50]. A potential new NAP was identified in ‘*Ca. M. alvus*’ Mx1021 [50] and corresponded to AYQ55638.1 in Mx-05<sup>T</sup>.


**Table 1.** Comparison of characteristics of strains Mx-05<sup>T</sup> (1) and *Methanomassiliicoccus luminyensis* B10<sup>T</sup> (2).

| Characteristics                    | 1  | 2  |
|------------------------------------|--|--|
| Source                             | Human feces  | Human feces  |
| Genome length (bp)                 | 1,666,767  | 2,637,810  |
| DNA G+C content (mol%)             | 55.5   | 59.93  |
| Morphology                         | Cocci, as single cells   | Cocci, as single cells   |
| Diameter (µm)                      | 0.4–0.7  | 0.7–1.0  |
| Motility                           | Non-motile   | Non-motile   |
| Growth temperature                 |  |  |
| Range                              | 30–40  | 30–40  |
| Optimum                            | 37   | 37   |
| Growth pH                          |  |  |
| Range                              | 6.9–8.3  | 7.2–8.4  |
| Optimum                            | 7.5  | 7.6  |
| Growth NaCl (mol.L <sup>-1</sup> ) |  |  |
| Range                              | 0.02–0.34  | 0.02–0.26  |
| Optimum                            | 0.12   | 0.17   |
| Required growth factor             | Filtrate from a grown <i>E. lenta</i> or rumen fluid   | acetate  |
| Substrates for methane production  | H <sub>2</sub> /methanol<br>H <sub>2</sub> /monomethylamine<br>H <sub>2</sub> /dimethylamine<br>H <sub>2</sub> /trimethylamine | H <sub>2</sub> /methanol<br>H <sub>2</sub> /monomethylamine<br>H <sub>2</sub> /dimethylamine<br>H <sub>2</sub> /trimethylamine |
| Lysed by osmotic shock/SDS         | yes  | yes  |

### 3.3. Morphology

Mx-05<sup>T</sup> cells are cocci measuring 0.4–0.7 µm that are not autofluorescent when excited at the 420 nm and 350 nm wavelengths. This absence of autofluorescence matched the absence of the genes coding for the F<sub>420</sub> biosynthesis pathway in the Mx-05<sup>T</sup> genome. Under Cryo-EM microscopy (Figure 3), we did not observe a flagellum (archaeallum), concordant with the absence of the corresponding genes in the genome. Long fibrillar appendices (up to 300 nm) are present on the cell surface (Figure 3b), which we speculate to represent the long adhesin-like proteins with Flg-new/List-Bac repeat domains (PF09479) coded by the genome of Mx-05<sup>T</sup>. No cell wall or S-layer structure was observed, which agrees with the results of cell lysing in SDS (0.01% (*w/v*)) and in distilled water. In contrast

to *M. luminyensis* B10<sup>T</sup> cells, which were reported to have two membranes [2], we only observed a single membrane in Mx-05<sup>T</sup> cells.



**Figure 3.** Cryo-electron microscopy pictures of strain Mx-05<sup>T</sup> showing the whole cell (a) and details (b). Black arrows (b) indicate potential adhesin-like proteins on the surface of the cell.

### 3.4. Physiology

Mx-05<sup>T</sup> is a strict anaerobe that uses methyl compounds as electron acceptors while oxidizing H<sub>2</sub> for methanogenesis. The methyl compounds used for methanogenesis comprise methanol, monomethylamine, dimethylamine, and trimethylamine but not dimethyl sulfide. The Mx-05<sup>T</sup> genome has homologues of *mtsAB* genes that are involved in the utilization of several methylated thiol compounds for methanogenesis in *Methanosarcina barkeri* [51]. Those genes are likely not involved in the transfer of methyl from dimethylsulfide to coenzyme M in Mx-05<sup>T</sup>, as neither growth nor methane production was observed when dimethylsulfide and H<sub>2</sub> were added to the culture medium. Instead, MtsAB homologues may catalyze methyl transfer from other types of methylated thiol to be tested (e.g., methylthiol, dimethylsulfoniopropionate). No growth was observed in the absence of H<sub>2</sub>. Formate and ethanol were not used as electron donors to reduce methyl compounds. Formate, H<sub>2</sub>/CO<sub>2</sub>, and acetate were not used for methanogenesis.

The growth of Mx-05<sup>T</sup> required an unknown factor that could be supplied by the culture supernatant filtrate of an *Eggerthella* sp., which had been isolated from the initial enrichment culture. A similar growth enablement was obtained using *E. lenta* DSM 2243<sup>T</sup>. The growth rate of Mx-05<sup>T</sup> was correlated with the proportional volume (2–10% (v/v)) of *E. lenta* supernatant filtrate added to the culture. The unknown growth factor could also be provided by rumen fluid, thereby indicating that *E. lenta* and possibly other bacteria growing in the rumen produce the factor(s) required by Mx-05<sup>T</sup>. In comparison to the *E. lenta* culture filtrate, the growth of Mx-05<sup>T</sup> was not influenced by the volume of rumen fluid added to the medium (in the range of 2–10% (v/v)), thereby indicating that the unknown growth factor is present at a higher concentration in the rumen fluid than in the *E. lenta* filtrate. The unknown factor does not appear to be sensitive to temperature, as the growth of the methanogenic isolate was similar when using an autoclaved culture filtrate of *E. lenta*.

The growth of Mx-05<sup>T</sup> was observed between 30 °C and 40 °C, with an optimum temperature at 37 °C. The pH range for growth was between 6.9 and pH 8.3, with an optimum value found at pH 7.5. At pH 6.9, Mx-05<sup>T</sup> grows almost as well as at pH 7.5, although a slightly lower pH is strongly inhibitory. Regarding salinity, Mx-05<sup>T</sup> can grow

in conditions of between 0.02 and 0.34 mol.L<sup>-1</sup> NaCl, with an optimum value of around 0.12 mol.L<sup>-1</sup> NaCl. Mx-05<sup>T</sup> grows in conditions almost similar to those of *M. luminyensis* (Table 1) [2]. Under optimal conditions (37 °C, pH 7.5, and 0.09 mmol.L<sup>-1</sup> NaCl), the maximum specific growth rate of Mx-05<sup>T</sup> was 0.026 h<sup>-1</sup> (i.e., doubling time of 26.7 h).

#### 4. Conclusions

Metagenomic and metataxonomic studies have revealed the global environmental distribution of *Methanomassiliicoccales* representatives [4,7,52,53] and a particular association with the gut microbiomes of animals [3,4,19], including ruminants [54,55] and humans [5,6,56]. Over the last 10 years, multiple enrichment cultures of *Methanomassiliicoccales* have been described [1,4,7,9,10,13,14,17,18], although only one generalist member (*M. luminyensis* B10<sup>T</sup>) of this order has been obtained in axenic culture and described so far [2]. Here, we report the description of a second hydrogenotrophic methyl-reducing methanogen representative of this order (strain Mx-05<sup>T</sup>), being the first isolate of a family-level clade associated with the gut microbiome of various animals. Based on physiological, phylogenetic, and genetic features, we propose strain Mx-05<sup>T</sup> as pertaining to a novel archaeal genus designated as *Methanomethylophilus* gen. nov., and named as *M. alvi* sp. nov. within the family *Methanomethylophilaceae* fam. nov. The originally proposed Candidatus name for this species (*M. alvus*) was changed to *M. alvi* following a recommendation by Oren et al. [15]. In contrast to *M. luminyensis* B10<sup>T</sup>, *Methanomethylophilus alvi* Mx-05<sup>T</sup> requires one or multiple unknown growth factors elaborated by *Eggerthella lenta*, of the family *Eggerthellaceae*, whose members are also recognized as members of the mammalian gut microbiome [57,58]. This requirement highlights the inter-species dependencies that can occur between members of the gut microbiota and warrants further biochemical and molecular investigations. The addition of the culture supernatant filtrate from the last co-cultured bacterium in the enrichment was critical to the isolation of *M. alvi* Mx-05<sup>T</sup>, as it eliminated the need to have this live bacterium in culture. Such a strategy could enable the isolation of other *Methanomassiliicoccales* members from animal or human gut microbiomes, which have so far only been obtained in mixed enrichment cultures, or more largely for other microorganisms that remain recalcitrant to isolation. Interestingly, in the case of *M. alvi* Mx-05<sup>T</sup>, these unknown growth factors were also present in rumen fluid, where they most probably result from microbial activities and may explain the presence of *Methanomethylophilus* spp. in both the ruminant and human gut environments. Further studies on *M. alvi* Mx-05<sup>T</sup> could be relevant for different human health aspects, such as its utilization as a probiotic (archaeobiotics, [16]) either to prevent the harmful effects of trimethylamine on human cardiovascular health or to develop methane mitigation strategies in ruminants.

##### 4.1. Description of *Methanomethylophilus* gen. nov.

*Methanomethylophilus* (Me.tha.no.me.thy.lo'phi.lus. N.L. pref. *methano-*, pertaining to methane (in compound words); N.L. neut. n. *methyl*, pertaining to the methyl group; N.L. masc. adj. suff. *-philus*, loving, from Gr. masc. adj. *philos*, on; N.L. masc. n. *Methanomethylophilus*, methane-producing organism-loving methyl groups). Strictly anaerobic and methanogenic cells. They only obtain energy by reducing methyl compounds into methane using H<sub>2</sub> as the electron donor (methyl-reducing hydrogenotrophic cells). Predominantly living in the intestinal microbiome. The type species is *Methanomethylophilus alvi*.

##### 4.2. Description of *Methanomethylophilus alvi* sp. nov.

*Methanomethylophilus alvi* (al'vi. L. gen. n. *alvi* of the bowels).

In addition to the characters described for the genus, the species is characterized by the following properties. The cells are single-membrane cocci measuring 0.4–0.7 µm, occurring alone. The cells lyse rapidly in SDS (0.01%) and in distilled water. Motility is not observed and genes coding for the archaellum are absent. Growth occurs between 30 °C and 40 °C (optimum, 37 °C), between pH 6.9 and 8.3 (optimum pH 7.5), and between 0.02 and 0.34 mol.L<sup>-1</sup> NaCl (optimum 0.12 mol.L<sup>-1</sup> NaCl). Obligate anaerobe. Produces methane

using hydrogen as an energy source and methanol, trimethylamine, dimethylamine, or monomethylamine only as electron acceptors. Requires the addition of the filtrate of a grown culture of *Eggerthella* sp., *Eggerthella lenta*, or rumen fluid for growth.

The type strain Mx-05<sup>T</sup> (=JCM 31474<sup>T</sup>) was isolated from a fecal sample of an elderly subject.

#### 4.3. Description of Methanomethylophilaceae fam. nov.

*Methanomethylophilaceae* (Me.tha.no.me.thy.lo.phi.la'ce.ae. N.L. fem. pl. n. *Methanomethylophilus* is the type genus of the family; -aceae ending to denote a family; N.L. fem. pl. n. *Methanomethylophilaceae* the family of the genus *Methanomethylophilus*).

Mesophilic methanogens non-motile. Found mostly in the animal digestive tract. Obligate anaerobes. All sequenced members lack the genes of the H<sub>4</sub>MPT-methyl branch of the Wood–Ljungdahl pathway and likely cannot disproportionate methyl compounds into methane and CO<sub>2</sub> or reduce CO<sub>2</sub> while oxidizing H<sub>2</sub>. Most have genes coding for methyltransferases involved in the transfer of methyl groups from methyl compound substrates to coenzyme M and genes coding for hydrogenases to reduce the methyl groups of these compounds using H<sub>2</sub> as electron donor.

The type genus is *Methanomethylophilus*.

This article is dedicated to the American microbiologist Professor R.A. Mah, for his important contribution to the physiology and taxonomy of methanogens.

## 5. Patents

One patent resulted from the work reported here, under the reference WO2019106181A1.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/microorganisms11112794/s1>. Figure S1: Position of Mx-05<sup>T</sup> in the 16S rRNA gene phylogeny of the *Methanomassiliicoccales*. Maximum likelihood tree built with GTR+F+R4 model. The tree is rooted on 'Candidatus Sysuiplasmatales' [38], with 1550 positions after trimming. Values on the tree indicate bootstrap support > 70%.

**Author Contributions:** Conceptualization, G.B., B.O. and J.-F.B.; formal analysis, G.B. and K.F.; investigation, G.B. and K.F.; methodology, G.B., K.F., W.B.H., N.G., G.P.-A., P.P.C., M.-L.F., B.O. and J.-F.B.; supervision, G.B., B.O. and J.-F.B.; validation, G.B., K.F., B.O. and J.-F.B.; visualization, G.B., K.F. and G.P.-A.; writing—original draft, G.B.; writing—review and editing, G.B., P.V., N.B., M.A., P.W.O., B.O. and J.-F.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by French National Agency for Research Grants, grant number ANR-19-CE02-0005-01 (Methevol Project). Lesaffre International made a financial contribution to this work through a research collaboration. PWOT is supported by Science Foundation Ireland through a Centre award to the APC.

**Data Availability Statement:** The GenBank accession number for the complete genome sequence of *Methanomethylophilus alvi* Mx-05<sup>T</sup> is CP017686.1. The strain was deposited at the Japan Collection of Microorganisms (JCM) under the number JCM 31474T.

**Acknowledgments:** We would like to thank M. Nilges and the Equipment d'excellence CACSICE (grant 924 number ANR-11-EQPX-0008) for providing the Falcon II direct electron detector. The authors are grateful to Lise Cao Van Tuat and Caroline Soulard for their valuable technical help. J.-F.B. and B.O. thank Lesaffre International, and more specifically J-Luc Blond, for the collaboration and help to patent some resulting findings.

**Conflicts of Interest:** P.V. and N.B. are full-time employees of Lesaffre. The other authors report no conflict of interest.

## References

1. Iino, T.; Tamaki, H.; Tamazawa, S.; Ueno, Y.; Ohkuma, M.; Suzuki, K.; Igarashi, Y.; Haruta, S. *Candidatus Methanogranum caenicola*: A Novel Methanogen from the Anaerobic Digested Sludge, and Proposal of *Methanomassiliicoccales* fam. nov. and *Methanomassiliicoccales* ord. nov., for a Methanogenic Lineage of the Class *Thermoplasmata*. *Microbes Environ.* **2013**, *28*, 244–250. [[CrossRef](#)] [[PubMed](#)]

2. Dridi, B.; Fardeau, M.L.; Ollivier, B.; Raoult, D.; Drancourt, M. *Methanomassiliicoccus luminyensis* gen. nov., sp. nov., a Methanogenic Archaeon Isolated from Human Faeces. *Int. J. Syst. Evol. Microbiol.* **2012**, *62*, 1902–1907. [[CrossRef](#)]
3. Thomas, C.M.; Desmond-Le Quéméner, E.; Gribaldo, S.; Borrel, G. Factors Shaping the Abundance and Diversity of the Gut Archaeome across the Animal Kingdom. *Nat. Commun.* **2022**, *13*, 3358. [[CrossRef](#)]
4. Söllinger, A.; Schwab, C.; Weinmaier, T.; Loy, A.; Tveit, A.T.; Schleper, C.; Urich, T. Phylogenetic and Genomic Analysis of *Methanomassiliicoccales* in Wetlands and Animal Intestinal Tracts Reveals Clade-Specific Habitat Preferences. *FEMS Microbiol. Ecol.* **2016**, *92*, fiv149. [[CrossRef](#)] [[PubMed](#)]
5. Borrel, G.; McCann, A.; Deane, J.; Neto, M.C.; Lynch, D.B.; Brugère, J.F.; O’Toole, P.W. Genomics and Metagenomics of Trimethylamine-Utilizing Archaea in the Human Gut Microbiome. *ISME J.* **2017**, *11*, 2059–2074. [[CrossRef](#)] [[PubMed](#)]
6. Chibani, C.M.; Mahnert, A.; Borrel, G.; Almeida, A.; Werner, A.; Brugère, J.-F.; Gribaldo, S.; Finn, R.D.; Schmitz, R.A.; Moissl-Eichinger, C. A Catalogue of 1,167 Genomes from the Human Gut Archaeome. *Nat. Microbiol.* **2022**, *7*, 48–61. [[CrossRef](#)]
7. Cozannet, M.; Borrel, G.; Roussel, E.; Moalic, Y.; Allieux, M.; Sanvoisin, A.; Toffin, L.; Alain, K. New Insights into the Ecology and Physiology of *Methanomassiliicoccales* from Terrestrial and Aquatic Environments. *Microorganisms* **2021**, *9*, 30. [[CrossRef](#)]
8. Chen, C.; Li, L.; Wang, Y.; Dong, X.; Zhao, F.-J. Methylophilic Methanogens and Bacteria Synergistically Demethylate Dimethylarsenate in Paddy Soil and Alleviate Rice Straighthead Disease. *ISME J.* **2023**, *17*, 1851–1861. [[CrossRef](#)]
9. Borrel, G.; Harris, H.M.B.; Parisot, N.; Gaci, N.; Tottey, W.; Mihajlovski, A.; Deane, J.; Bardot, O.; Peyretailade, E.; et al. Genome Sequence of “*Candidatus* Methanomassiliicoccus intestinalis” Isoire-Mx1, a Third *Thermoplasmatales*-Related Methanogenic Archaeon from Human Feces. *Genome Announc.* **2013**, *1*, e00453-13. [[CrossRef](#)]
10. Weil, M.; Hoff, K.J.; Meißner, W.; Schäfer, F.; Söllinger, A.; Wang, H.; Hagenau, L.; Kuss, A.W.; Urich, T. Full Genome Sequence of a *Methanomassiliicoccales* Representative Enriched from Peat Soil. *Microbiol. Resour. Announc.* **2021**, *10*, e00443-21. [[CrossRef](#)]
11. Janssen, P.H.; Kirs, M. Structure of the Archaeal Community of the Rumen. *Appl. Environ. Microbiol.* **2008**, *74*, 3619–3625. [[CrossRef](#)] [[PubMed](#)]
12. Borrel, G.; O’Toole, P.W.; Harris, H.M.B.; Peyret, P.; Brugère, J.F.; Gribaldo, S. Phylogenomic Data Support a Seventh Order of Methylophilic Methanogens and Provide Insights into the Evolution of Methanogenesis. *Genome Biol. Evol.* **2013**, *5*, 1769–1780. [[CrossRef](#)] [[PubMed](#)]
13. Paul, K.; Nonoh, J.O.; Mikulski, L.; Brune, A. “*Methanoplasmatales*,” *Thermoplasmatales*-Related Archaea in Termite Guts and Other Environments, Are the Seventh Order of Methanogens. *Appl. Environ. Microbiol.* **2012**, *78*, 8245–8253. [[CrossRef](#)]
14. Borrel, G.; Harris, H.M.B.; Tottey, W.; Mihajlovski, A.; Parisot, N.; Peyretailade, E.; Peyret, P.; Gribaldo, S.; O’Toole, P.W.; Brugère, J.F. Genome Sequence of “*Candidatus* Methanomethylophilus alvus” Mx1201, a Methanogenic Archaeon from the Human Gut Belonging to a Seventh Order of Methanogens. *J. Bacteriol.* **2012**, *194*, 6944–6945. [[CrossRef](#)]
15. Oren, A.; Garrity, G.M.; Parker, C.T.; Chuvochina, M.; Trujillo, M.E. Lists of Names of Prokaryotic Candidatus Taxa. *Int. J. Syst. Evol. Microbiol.* **2020**, *70*, 3956–4042. [[CrossRef](#)] [[PubMed](#)]
16. Brugère, J.F.; Borrel, G.; Gaci, N.; Tottey, W.; O’Toole, P.W.; Malpuech-Brugère, C. Archaeobiotics: Proposed Therapeutic Use of Archaea to Prevent Trimethylaminuria and Cardiovascular Disease. *Gut Microbes* **2014**, *5*, 5–10. [[CrossRef](#)]
17. Li, Y.; Leahy, S.C.; Jeyanathan, J.; Henderson, G.; Cox, F.; Altermann, E.; Kelly, W.J.; Lambie, S.C.; Janssen, P.H.; Rakonjac, J.; et al. The Complete Genome Sequence of the Methanogenic Archaeon ISO4-H5 Provides Insights into the Methylophilic Lifestyle of a Ruminant Representative of the *Methanomassiliicoccales*. *Stand. Genom. Sci.* **2016**, *11*, 59. [[CrossRef](#)]
18. Noel, S.J.; Højberg, O.; Urich, T.; Poulsen, M. Draft Genome Sequence of “*Candidatus* Methanomethylophilus” Sp. 1R26, Enriched from Bovine Rumen, a Methanogenic Archaeon Belonging to the *Methanomassiliicoccales* Order. *Genome Announc.* **2016**, *4*, e01734-15. [[CrossRef](#)]
19. Lang, K.; Schuldes, J.; Klingl, A.; Poehlein, A.; Daniel, R.; Brune, A. New Mode of Energy Metabolism in the Seventh Order of Methanogens as Revealed by Comparative Genome Analysis of “*Candidatus* Methanoplasma Termitum”. *Appl. Environ. Microbiol.* **2015**, *81*, 1338–1352. [[CrossRef](#)]
20. de la Cuesta-Zuluaga, J.; Spector, T.D.; Youngblut, N.D.; Ley, R.E. Genomic Insights into Adaptations of Trimethylamine-Utilizing Methanogens to Diverse Habitats, Including the Human Gut. *mSystems* **2021**, *6*, e00939-20. [[CrossRef](#)]
21. Borrel, G.; Parisot, N.; Harris, H.M.; Peyretailade, E.; Gaci, N.; Tottey, W.; Bardot, O.; Raymann, K.; Gribaldo, S.; Peyret, P.; et al. Comparative Genomics Highlights the Unique Biology of *Methanomassiliicoccales*, a *Thermoplasmatales*-Related Seventh Order of Methanogenic Archaea That Encodes Pyrrolysine. *BMC Genom.* **2014**, *15*, 679. [[CrossRef](#)] [[PubMed](#)]
22. Kröninger, L.; Berger, S.; Welte, C.; Deppenmeier, U. Evidence for the Involvement of Two Heterodisulfide Reductases in the Energy-Conserving System of *Methanomassiliicoccus luminyensis*. *FEBS J.* **2016**, *283*, 472–483. [[CrossRef](#)] [[PubMed](#)]
23. Kröninger, L.; Steiniger, F.; Berger, S.; Kraus, S.; Welte, C.U.; Deppenmeier, U. Energy Conservation in the Gut Microbe *Methanomassiliicoccus luminyensis* Is Based on Membrane-bound Ferredoxin Oxidation Coupled to Heterodisulfide Reduction. *FEBS J.* **2019**, *286*, 3831–3843. [[CrossRef](#)] [[PubMed](#)]
24. Becker, K.W.; Elling, F.J.; Yoshinaga, M.Y.; Söllinger, A.; Urich, T.; Hinrichs, K.U. Unusual Butane- and Pentanetriol-Based Tetraether Lipids in *Methanomassiliicoccus luminyensis*, a Representative of the Seventh Order of Methanogens. *Appl. Environ. Microbiol.* **2016**, *82*, 4505–4516. [[CrossRef](#)]
25. Coffinet, S.; Mühlhena, L.; Lipp, J.S.; Weil, M.; Neubauer, C.; Urich, T.; Hinrichs, K.-U. Evidence for Enzymatic Backbone Methylation of the Main Membrane Lipids in the Archaeon *Methanomassiliicoccus luminyensis*. *Appl. Environ. Microbiol.* **2022**, *88*, e02154-21. [[CrossRef](#)]

26. Brugère, J.-F.; Atkins, J.F.; O'Toole, P.W.; Borrel, G. Pyrrolysine in Archaea: A 22nd Amino Acid Encoded through a Genetic Code Expansion. *Emerg. Top. Life Sci.* **2018**, *2*, 607–618.
27. Borrel, G.; Gaci, N.; Peyret, P.; O'Toole, P.W.; Gribaldo, S.; Brugère, J.F. Unique Characteristics of the Pyrrolysine System in the 7th Order of Methanogens: Implications for the Evolution of a Genetic Code Expansion Cassette. *Archaea* **2014**, *2014*, 374146. [[CrossRef](#)]
28. Willis, J.C.W.; Chin, J.W. Mutually Orthogonal Pyrrolysyl-TRNA Synthetase/TRNA Pairs. *Nat. Chem.* **2018**, *10*, 831–837. [[CrossRef](#)]
29. Li, S.; Wang, N.; Yu, B.; Sun, W.; Wang, L. Genetically Encoded Chemical Crosslinking of Carbohydrate. *Nat. Chem.* **2023**, *15*, 33–42. [[CrossRef](#)]
30. Tang, W.H.W.; Wang, Z.; Kennedy, D.J.; Wu, Y.; Buffa, J.A.; Agatista-Boyle, B.; Li, X.S.; Levison, B.S.; Hazen, S.L. Gut Microbiota-Dependent Trimethylamine N-Oxide (TMAO) Pathway Contributes to Both Development of Renal Insufficiency and Mortality Risk in Chronic Kidney Disease. *Circ. Res.* **2015**, *116*, 448–455. [[CrossRef](#)]
31. Zhu, W.; Gregory, J.C.; Org, E.; Buffa, J.A.; Gupta, N.; Wang, Z.; Li, L.; Fu, X.; Wu, Y.; Mehrabian, M.; et al. Gut Microbial Metabolite TMAO Enhances Platelet Hyperreactivity and Thrombosis Risk. *Cell* **2016**, *165*, 111–124. [[CrossRef](#)]
32. Mackay, R.J.; McEntyre, C.J.; Henderson, C.; Lever, M.; George, P.M. Trimethylaminuria: Causes and Diagnosis of a Socially Distressing Condition. *Clin. Biochem. Rev.* **2011**, *32*, 33–43.
33. Fadhlou, K.; Arnal, M.-E.; Martineau, M.; Camponova, P.; Ollivier, B.; O'Toole, P.W.; Brugère, J.-F. Archaea, Specific Genetic Traits, and Development of Improved Bacterial Live Biotherapeutic Products: Another Face of next-Generation Probiotics. *Appl. Microbiol. Biotechnol.* **2020**, *104*, 4705–4716. [[CrossRef](#)] [[PubMed](#)]
34. Widdel, F.; Pfennig, N. Studies on Dissimilatory Sulfate-Reducing Bacteria That Decompose Fatty Acids: I. Isolation of New Sulfate-Reducing Bacteria Enriched with Acetate from Saline Environments. Description of *Desulfobacter postgatei* Gen. Nov., Sp. Nov. *Arch. Microbiol.* **1981**, *129*, 395–400. [[CrossRef](#)]
35. Tatusova, T.; DiCuccio, M.; Badretdin, A.; Chetvernin, V.; Nawrocki, E.P.; Zaslavsky, L.; Lomsadze, A.; Pruitt, K.D.; Borodovsky, M.; Ostell, J. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res.* **2016**, *44*, 6614–6624. [[CrossRef](#)] [[PubMed](#)]
36. Borrel, G.; Adam, P.S.; McKay, L.J.; Chen, L.-X.; Sierra-García, I.N.; Sieber, C.M.K.; Letourneur, Q.; Ghoulane, A.; Andersen, G.L.; Li, W.-J.; et al. Wide Diversity of Methane and Short-Chain Alkane Metabolisms in Uncultured Archaea. *Nat. Microbiol.* **2019**, *4*, 603–613. [[CrossRef](#)]
37. Parks, D.H.; Chuvochina, M.; Rinke, C.; Mussig, A.J.; Chaumeil, P.-A.; Hugenholtz, P. GTDB: An Ongoing Census of Bacterial and Archaeal Diversity through a Phylogenetically Consistent, Rank Normalized and Complete Genome-Based Taxonomy. *Nucleic Acids Res.* **2022**, *50*, D785–D794. [[CrossRef](#)]
38. Yuan, Y.; Liu, J.; Yang, T.-T.; Gao, S.-M.; Liao, B.; Huang, L.-N. Genomic Insights into the Ecological Role and Evolution of a Novel *Thermoplasmata* Order, “*Candidatus Sysuiplasmatales*”. *Appl. Environ. Microbiol.* **2021**, *87*, e01065-21. [[CrossRef](#)] [[PubMed](#)]
39. Johnson, L.S.; Eddy, S.R.; Portugaly, E. Hidden Markov Model Speed Heuristic and Iterative HMM Search Procedure. *BMC Bioinform.* **2010**, *11*, 431. [[CrossRef](#)] [[PubMed](#)]
40. Katoh, K.; Standley, D.M. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Mol. Biol. Evol.* **2013**, *30*, 772–780. [[CrossRef](#)]
41. Criscuolo, A.; Gribaldo, S. BMGE (Block Mapping and Gathering with Entropy): A New Software for Selection of Phylogenetic Informative Regions from Multiple Sequence Alignments. *BMC Evol. Biol.* **2010**, *10*, 210. [[CrossRef](#)]
42. Nguyen, L.T.; Schmidt, H.A.; Von Haeseler, A.; Minh, B.Q. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol. Biol. Evol.* **2015**, *32*, 268–274. [[CrossRef](#)]
43. Bengtsson-Palme, J.; Hartmann, M.; Eriksson, K.M.; Pal, C.; Thorell, K.; Larsson, D.G.J.; Nilsson, R.H. METAXA2: Improved Identification and Taxonomic Classification of Small and Large Subunit rRNA in Metagenomic Data. *Mol. Ecol. Resour.* **2015**, *15*, 1403–1414. [[CrossRef](#)] [[PubMed](#)]
44. Flemer, B.; Gaci, N.; Borrel, G.; Sanderson, I.R.; Chaudhary, P.P.; Tottey, W.; O'Toole, P.W.; Brugère, J.F. Fecal Microbiota Variation across the Lifespan of the Healthy Laboratory Rat. *Gut Microbes* **2017**, *8*, 428–439. [[CrossRef](#)] [[PubMed](#)]
45. Letunic, I.; Bork, P. Interactive Tree of Life (ITOL) v4: Recent Updates and New Developments. *Nucleic Acids Res.* **2019**, *47*, 256–259. [[CrossRef](#)]
46. Jüttner, M.; Ferreira-Cerca, S. Looking through the Lens of the Ribosome Biogenesis Evolutionary History: Possible Implications for Archaeal Phylogeny and Eukaryogenesis. *Mol. Biol. Evol.* **2022**, *39*, msac054. [[CrossRef](#)] [[PubMed](#)]
47. Thomas, C.M.; Taib, N.; Gribaldo, S.; Borrel, G. Comparative Genomic Analysis of *Methanimicrococcus blatticola* Provides Insights into Host-Adaptation in Archaea and the Evolution of Methanogenesis. *ISME Commun.* **2021**, *1*, 47. [[CrossRef](#)]
48. Martins, M.; Aymeric, L.; Du Merle, L.; Danne, C.; Robbe-Masselot, C.; Trieu-Cuot, P.; Sansonetti, P.; Dramsi, S. *Streptococcus gallolyticus* Pil3 Pilus Is Required for Adhesion to Colonic Mucus and for Colonization of Mouse Distal Colon. *J. Infect. Dis.* **2015**, *212*, 1646–1655. [[CrossRef](#)]
49. Medvedeva, S.; Borrel, G.; Krupovic, M.; Gribaldo, S. A Compendium of Viruses from Methanogenic Archaea Reveals Their Diversity and Adaptations to the Gut Environment. *Nat. Microbiol.* **2023**, *8*, 2170–2182. [[CrossRef](#)]
50. Hocher, A.; Borrel, G.; Fadhlou, K.; Brugère, J.-F.; Gribaldo, S.; Warnecke, T. Growth Temperature and Chromatinization in Archaea. *Nat. Microbiol.* **2022**, *7*, 1932–1942. [[CrossRef](#)]

51. Tallant, T.C.; Krzycki, J.A. Methylthiol: Coenzyme M Methyltransferase from *Methanosarcina barkeri*, an Enzyme of Methanogenesis from Dimethylsulfide and Methylmercaptopropionate. *J. Bacteriol.* **1997**, *179*, 6902–6911. [[CrossRef](#)] [[PubMed](#)]
52. Weil, M.; Wang, H.; Zak, D.; Urich, T. Spatial and Temporal Niche Separation of *Methanomassiliicoccales* Phylotypes in Temperate Fens. *FEMS Microbiol. Ecol.* **2023**, *99*, fiad049. [[CrossRef](#)] [[PubMed](#)]
53. Zhang, C.-J.; Pan, J.; Liu, Y.; Duan, C.-H.; Li, M. Genomic and Transcriptomic Insights into Methanogenesis Potential of Novel Methanogens from Mangrove Sediments. *Microbiome* **2020**, *8*, 94. [[CrossRef](#)] [[PubMed](#)]
54. Seedorf, H.; Kittelmann, S.; Janssen, P.H. Few Highly Abundant Operational Taxonomic Units Dominate within Rumen Methanogenic Archaeal Species in New Zealand Sheep and Cattle. *Appl. Environ. Microbiol.* **2015**, *81*, 986–995. [[CrossRef](#)]
55. Poulsen, M.; Schwab, C.; Jensen, B.B.; Engberg, R.M.; Spang, A.; Canibe, N.; Højberg, O.; Milinovich, G.; Fragner, L.; Schleper, C.; et al. Methylotrophic Methanogenic *Thermoplasmata* Implicated in Reduced Methane Emissions from Bovine Rumen. *Nat. Commun.* **2013**, *4*, 1428. [[CrossRef](#)]
56. Borrel, G.; Gribaldo, S.; Brugère, J.-F.; Schmitz, R.; Moissl-Eichinger, C. The Host-Associated Archaeome. *Nat. Rev. Microbiol.* **2020**, *18*, 622–636. [[CrossRef](#)]
57. Jackson, M.A.; Jeffery, I.B.; Beaumont, M.; Bell, J.T.; Clark, A.G.; Ley, R.E.; O’Toole, P.W.; Spector, T.D.; Steves, C.J. Signatures of Early Frailty in the Gut Microbiota. *Genome Med.* **2016**, *8*, 8. [[CrossRef](#)]
58. Wang, K.; Zhang, H.; Hu, L.; Zhang, G.; Lu, H.; Luo, H.; Zhao, S.; Zhu, H.; Wang, Y. Characterization of the Microbial Communities along the Gastrointestinal Tract in Crossbred Cattle. *Animals* **2022**, *12*, 825. [[CrossRef](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.