



Implant-supported prostheses in patients with special needs: A systematic literature review of protocols and outcomes

Marie-Sophie Bogner, Vivien Chambas, Emmanuel Nicolas, Denise Faulks,
Martine Hennequin

► To cite this version:

Marie-Sophie Bogner, Vivien Chambas, Emmanuel Nicolas, Denise Faulks, Martine Hennequin. Implant-supported prostheses in patients with special needs: A systematic literature review of protocols and outcomes. Journal of Prosthetic Dentistry, 2023, 10.1016/j.jprosdent.2023.09.013 . hal-04323669

HAL Id: hal-04323669

<https://uca.hal.science/hal-04323669v1>

Submitted on 12 Dec 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

SYSTEMATIC REVIEW

Implant-supported prostheses in patients with special needs: A systematic literature review of protocols and outcomes

Marie-Sophie Bogner, BDS, MSc,^a Vivien Chambas, BDS,^b Emmanuel Nicolas, BDS, MSc, PhD, HDR,^c
Denise Faulks, BDS, MSc, PhD, HDR,^d and Martine Hennequin, BDS, MSc, PhD, HDR^e

Patients with special needs are at higher risk of missing teeth when compared with the general population because genetic disorders can be associated with agenesis and aggressive periodontal disease.^{1,2} Early tooth loss may also be associated with poor access to care, leading to more advanced oral disease when services are first contacted. In addition, these patients are more likely to undergo extractions than periodontal, endodontic, or restorative treatment compared with the general population.³ Missing teeth that are not replaced may lead to social and functional impairments and disability, and functional oral rehabilitation is challenging to provide for patients with special needs.^{4–6} In particular, removable dentures are often declined or abandoned after a short period of use.^{7–9} Handling and

ABSTRACT

Statement of problem. Implant-supported prostheses for individuals with special needs have been considered high-risk because of the prevalence of poor oral hygiene and parafunction in this population; however, systematic evidence on protocols and outcomes is scarce.

Purpose. The purpose of this systematic review was to summarize implant procedures and outcomes for the special needs population.

Material and methods. A systematic literature review was conducted using PubMed, Embase, and Scopus databases in accordance with the preferred reporting items for systematic reviews and meta-analyses (PRISMA) methodological framework. Data collected included patient information, location and number of implants, surgical and prosthetic procedures, behavior support techniques, postoperative follow-up, complications, and criteria for outcome assessment. Survival rates were pooled for quantitative analysis, and a separate analysis pooling data from groups of special needs patients that present similar risks was undertaken.

Results. The qualitative analysis included 79 studies involving 759 patients and 2269 implants. The quantitative analysis included 65 studies concerning 666 patients with special needs and 1886 implants. The overall weighted implant survival rate was 95.4%, with a mean follow-up duration of 40.7 ± 29.5 months.

Conclusions. In individuals with special needs considered suitable for implant-supported prostheses by their clinician, implant survival rates were similar to those reported in the general population. (*J Prosthet Dent* xxxx;xxx:xxx-xxx)

cleaning removable dentures can be challenging for patients with physical or cognitive impairments. In addition, drug-associated xerostomia can lead to unfavorable outcomes.¹⁰ The benefits of implant-supported or

Funding: This research was self-supported by University of Clermont Auvergne. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors have no conflict of interest to declare.

^aSpecialist, Centre for Research in Clinical Dentistry (CROC), University of Clermont Auvergne BP 10448, Clermont-Ferrand, France.

^bSpecialist, Centre for Research in Clinical Dentistry (CROC), University of Clermont Auvergne BP 10448, Clermont-Ferrand, France; and Private practice, Clermont-Ferrand, France.

^cProfessor, Centre for Research in Clinical Dentistry (CROC), University of Clermont Auvergne BP 10448, Clermont-Ferrand, France; and CHU Clermont-Ferrand, Service d'Odontologie, CHU Estaing, Clermont-Ferrand, France.

^dSpecialist Consultant, Centre for Research in Clinical Dentistry (CROC), University of Clermont Auvergne BP 10448, F-63000 Clermont-Ferrand, France; and CHU Clermont-Ferrand, Service d'Odontologie, CHU Estaing, Clermont-Ferrand, France.

^eProfessor, Centre for Research in Clinical Dentistry (CROC), University of Clermont Auvergne BP 10448, Clermont-Ferrand, France; and CHU Clermont-Ferrand, Service d'Odontologie, CHU Estaing, Clermont-Ferrand, France.

Clinical Implications

Complex medical conditions, cognitive and physical disabilities, and genetic disorders should not necessarily be contraindications to providing implant-supported prostheses.

implant-retained prostheses with improved oral health-related quality of life compared with removable dentures have been established in the general population, elderly patients with edentulism, and patients with tumors.^{11–19}

The high prevalence of poor oral hygiene, lack of cooperation, and the presence of parafunction have been commonly voiced against implant placement in patients with special needs.^{20–25} However, international evidence-based recommendations or guidelines for this population are lacking. This systematic review aimed to summarize implant procedures and outcomes for patients with special needs. Considering the diversity and multiplicity of risk factors for poor oral hygiene and parafunction in patients with special needs, the research hypothesis was that implant survival rates would be lower in patients with special needs than in the general population.

MATERIAL AND METHODS

A systematic literature review was conducted using the PubMed, Embase, and Scopus databases in accordance with the preferred reporting items for systematic reviews and meta-analyses (PRISMA) methodological framework in October 2021.²⁶ A follow-up search was conducted in November 2022 to identify any additional studies or clinical reports published after the initial search.

Two reviewers conducted the study selection independently (M-S.B., V.C.). A consensus on any uncertainties was obtained after discussion with a third researcher (M.H.). The electronic search in the databases had the following key words: (dental implant) AND ((disabled) OR (disability) OR (disabilities) OR (handicapped) OR (special need/s) OR (specific need/s) OR (medical compromised) OR (cerebral palsy) OR (phobia) OR (Down syndrome) OR (autism) OR (osteogenesis imperfecta) OR (cleidocranial dysplasia) OR (polyarthritis). A hand search was also undertaken in the *Journal of Disability and Oral Health*, which was not indexed in PubMed. Title and abstract were screened to identify studies with the following criteria: publications in English or French; clinical studies, case reports, and case series; patients with special needs undergoing dental implant placement; detailed information about the number of implants placed and loaded; rehabilitation finished, including prosthetic loading. Exclusion criteria were articles

not in English or French; in vitro or animal studies; use of implants other than dental endosseous titanium implants; no prosthetic loading; and systematic reviews of implant survival rates within specific subgroups of special care patients. The full text of the articles was assessed for eligibility. The references cited within each article were also assessed to avoid missing studies. Critical appraisal was assessed according to the JBI Critical Appraisal Checklists for case control studies, case reports, and case series.^{27,28}

Two investigators (M-S.B., V.C.) screened each included study independently. The following information was recorded: patient data: sociodemographic data and description of special needs; implant data: number per participant and location in the mouth; clinical procedures: surgical and prosthetic procedures, number and details of surgical and prosthetic complications that arose during the postoperative period; behavioral support procedures: general anesthesia (GA), conscious sedation, premedication, local anesthesia; study indicators: criteria used for outcome assessment.

Descriptive statistics were performed by measuring the weighted survival rate of the implants. Articles were eligible if they reported a follow-up duration (at least 1 follow-up appointment after prosthetic loading). For studies including patients with and without special needs, only those with special needs were included. For cohort studies, only the most recent data were included. The survival rate was calculated for each cohort or patient with a simple analysis of the number of implants lost weighted by the number of implants placed. The data were pooled with a separate analysis by pooling data from groups of patients with special needs that presented similar risks: patients able to perform oral hygiene and with no immune or metabolic/bone problems mentioned or deducted; patients needing support for oral hygiene because of dexterity or cognitive impairment and with no immune or metabolic/bone problems mentioned or deducted; and patients with impaired immunity and/or with a syndrome and/or medication affecting bone quality.

RESULTS

Fifty-four of the 124 articles assessed for eligibility were excluded, 15 of which reported systematic reviews of implant survival rates within specific subgroups of patients with special needs (geriatric patients²⁹; people taking antiresorptive drugs³⁰; those with Down syndrome,^{31,32} systemic conditions,^{33–36} oral manifestations of autoimmune diseases,^{37–42} and osteogenesis imperfecta).⁴³ Seventy-nine studies were included in the qualitative synthesis: 16 observational studies (2 case control studies and 14 case series),^{20,25,31,44–56} 59 case reports,^{57–115} and 4 articles presenting more than 1 clinical report.^{116–119} The quality of the included studies varied. Overall, studies described practices from 24

Table 1. Description of patients

Author	Country	No. of Patients	Age (Years)	Sex (M/W)	Special Needs as Described by Authors
Observational studies					
Cune et al, 2009 ²⁰	The Netherlands	61	43.2	NR	Severe epilepsy and multiple disabilities.
Corcueria-Flores et al, 2017 ⁴⁴	Spain	19	up to 40	36 implants/66 implants	Thirteen patients with cerebral palsy (71 implants), 6 with Down syndrome (31 implants).
De Bruyn et al, 2019 ⁴⁵	Belgium	8	45	6/2	Down syndrome.
Ekfeldt, 2005 ⁴⁶	Sweden	14	44.1	5/9	Neurologic disabilities.
Ekfeldt, 2013 ⁴⁷	Sweden	27	46	14/13	Neurologic disabilities.
Enkling et al, 2013 ⁴⁸	Switzerland	30	50.5	17/13	Fifteen patients with dental phobia versus 15 non-fearful patients.
Feijoo et al, 2012 ⁴⁹	Spain	8	36.7	5/3	Non-syndromic intellectual disability.
Jagadeesh et al, 2020 ⁵⁰	India	342	NR	NR	Patients with various medical conditions, diabetes mellitus, hypertension, mental disability, oral cancer, osteomyelitis.
Kim et al, 2017 ⁵¹	South Korea	19	32.9	8/11	Eleven patients with mental retardation, 3 with autism, 2 with cerebral palsy, 2 with schizophrenia, and 1 with brain disorder.
Krennmaier et al, 2010 ⁵²	Austria	34	58.1	0/34	Rheumatoid arthritis.
Limeres Posse et al, 2016 ⁵³	Spain	25	34	13/12	Down syndrome.
López-Jiménez et al, 2003 ²¹	Spain	18	34.7	7/11	Six patients with cerebral palsy, 3 with head injuries, 4 with Down syndrome, 3 with dementia, 1 with pyknodysostosis, 1 with Rieger's syndrome.
Öczakir et al, 2005 ⁵⁴	Switzerland	25	55.6	8/17	Cleft lip/palate (n=8), Down syndrome (n=3), Sjogren syndrome and scleroderma (n=2), ectodermal dysplasia (n=4), developmental retardation (n=2), chronic leukemia (n=2), lichen planus (n=1), cerebral palsy (n=1), deaf-muteness (n=1), amyotrophic lateral sclerosis (n=1).
Packer et al, 2009 ²⁵	UK	9	63	NR	Parkinson's disease.
Van de Velde et al, 2007 ⁵⁵	Belgium	18	39-77	8/10	Among patients, 1 with Down syndrome.
Weinländer et al, 2010 ⁵⁶	Austria	22	55.6	0/22	Rheumatoid arthritis and connective tissue diseases.
Case reports					
Apaydin et al, 2014 ⁵⁷	Turkey	1	17	W	Amelogenesis imperfecta.
Barker et al, 2008 ⁵⁸	UK	1	NR	NR	Ankylosing spondylitis.
Rogers, 1995 ⁵⁹	UK	1	64	NR	Athetoid cerebral palsy.
Hong et al, 2017 ⁶⁰	Republic of Korea	1	33	W	Cerebral palsy and mental retardation.
Eder and Watzek, 1999 ⁶¹	Austria	1	81	W	Chronic polyarthritis.
Ambard et al, 2019 ⁶²	USA	1	21	M	Cleidocranial dysostosis.
Angle and Rebello, 2005 ⁶³	USA	1	17	W	Cleidocranial dysostosis.
Ahmad et al, 2019 ⁶⁴	Sudan	1	24	W	Cleidocranial dysplasia.
Atil et al, 2018 ⁶⁵	Turkey	1	48	W	Cleidocranial dysplasia.
Calvo-Guirado et al, 2013 ⁶⁶	Spain	1	31	W	Cleidocranial dysplasia.
Daskalogiannakis et al, 2006 ⁶⁷	Canada	1	39	W	Cleidocranial dysplasia.
Lee et al, 2015 ⁶⁸	Republic of Korea	1	29	M	Cleidocranial dysplasia.
Lombardas and Toothaker, 1997 ⁶⁹	USA	1	15	M	Cleidocranial dysplasia.
Noh et al, 2014 ⁷⁰	South Korea	1	27	W	Cleidocranial dysplasia.
Petropoulos et al, 2004 ⁷¹	USA	1	42	W	Cleidocranial dysplasia.
Petropoulos et al, 2011 ⁷²	USA	1	45	W	Cleidocranial dysplasia.
Schnuttenhaus et al, 2015 ⁷³	Germany	1	43	M	Cleidocranial dysplasia.
Schnuttenhaus et al, 2021 ⁷⁴	Germany	1	26	M	Cleidocranial dysplasia.
Tan et al, 2020 ⁷⁵	Singapore	1	24	M	Cleidocranial dysplasia.
Balshi and Wolfinger, 2002 ⁷⁶	USA	1	30	W	Dental phobia.
Alqahtani et al, 2018 ⁷⁷	Saudi Arabia	1	44	M	Down syndrome.
Altintas et al, 2019 ⁷⁸	Turkey	1	37	W	Down syndrome.
Comparin et al, 2022 ⁷⁹	Brazil	1	50	M	Down syndrome.
Lustig et al, 2002 ⁸⁰	Israel	1	16	M	Down syndrome.
Reiche et al, 2015 ⁸¹	Hungary	1	18	M	Down syndrome.
Ribeiro et al, 2011 ⁸²	Brazil	1	36	W	Down syndrome.
Saponaro et al, 2016 ⁸³	USA	1	27	W	Down syndrome.
Schmidt et al, 2020 ⁸⁴	Germany	1	52	M	Down syndrome.
Soares et al, 2010 ⁸⁵	Brazil	1	22	M	Down syndrome.
Thomas and Simons, 2011 ⁸⁶	UK	1	49	M	Down syndrome.
Brahim et al, 1992 ⁸⁷	USA	1	38	M	Erdheim-Chester disease.
Engel et al, 1992 ⁸⁸	USA	1	30	W	Fanconi's anemia.
Gallotini et al, 2018 ⁸⁹	Brazil	1	16	W	Fraser syndrome.
Güven et al, 2010 ⁹⁰	Turkey	1	26	M	Freeman-Sheldon syndrome.
Merlini et al, 2016 ⁹¹	Italy	1	32	M	Gnathodiaphyseal dysplasia.
Deniz E et al, 2011 ⁹²	Turkey	1	67	W	Huntington's disease.
Jackowski et al, 2001 ⁹³	Germany	1	56	M	Huntington's disease.
Peñarrocha et al, 2001 ⁹⁴	Spain	1	52	W	Idiopathic torsion dystonia.
Károlyházy et al, 2014 ⁹⁵	Hungary	1	NR	NR	Intellectual disability with epilepsy.
Shek et al, 2014 ⁹⁶	USA	1	79	W	Involuntary mandibular movements.
Oelgiesser et al, 2006 ⁹⁷	Israel	1	19	W	Maple syrup urine disease.
Poluha et al, 2019 ⁹⁸	Brazil	1	40	W	Microstomia.
Peñarrocha et al, 2001 ⁹⁹	Spain	1	67	W	Oromandibular dystonia with blepharospasm.
Ullbro et al, 2000 ¹⁰⁰	Sweden	1	25	W	Papillon-Lefèvre syndrome.
Woo et al, 2003 ¹⁰¹	USA	1	13	M	Papillon-Lefèvre syndrome.
Applebaum et al, 1997 ¹⁰²	USA	1	72	W	Parkinson's disease.
Chu et al, 2004 ¹⁰³	China	1	84	W	Parkinson's disease.
Fiberg, 1994 ¹⁰⁴	Sweden	1	57	W	Polyarthritis

Table 1 (Continued)

Author	Country	No. of Patients	Age† (Years)	Sex (M/W)	Special Needs as Described by Authors
Bencharit et al, 2010 ¹⁰⁵	USA	1	74	W	Polymyalgia rheumatica.
Abdel-Kader et al 2019 ¹⁰⁶	Egypt	1	22	W	Regional odontodysplasia.
Gaur et al, 2021 ¹⁰⁷	India	1	60	W	Rheumatoid arthritis
Ella et al, 2011 ¹⁰⁸	France	1	56	W	Rheumatoid polyarthritis
Choi et al, 2019 ¹⁰⁹	South Korea	1	62	M	Schizophrenia.
Tizzoni et al, 2014 ¹¹⁰	Italy	1	54	M	Severe physical disability caused by neoplastic disease.
Thor et al, 2005 ¹¹¹	Sweden	1	60	W	Systemic mastocytosis.
Kelleher et al, 1998 ¹¹²	UK	1	69	W	Tardive dyskinesia.
Misch et al, 1998 ¹¹³	USA	1	64	W	Thalassemia.
Flanagan, 2011 ¹¹⁴	USA	1	24	W	Traumatic brain injury
Takeuchi et al, 2019 ¹¹⁵	USA	1	53	M	Traumatic brain injury
Durham et al, 2006 ¹¹⁶	USA	6	68	NR	Various cognitive disabilities.
Friberg, 2013 ¹¹⁷	Sweden	6	42.7	NR	Three patients with X-linked hypophosphatemia, 2 patients with Langerhans cell histiocytosis, 1 patient with osteogenesis/dentinogenesis imperfecta. Parkinson's disease.
Heckmann et al, 2000 ¹¹⁸	USA	3	75.7	2/1	
Owens et al, 2009 ¹¹⁹	Ireland	6	38.8	3/3	One patient with cerebral palsy and epilepsy, 1 patient with Morquio's syndrome, 2 with moderate learning disability, 1 with a factor VIII deficiency hemophilia associated to HIV and Hepatitis C, 1 with diabetes, kidney and pancreas transplant.

M, Man; No, Number; NR, Not reported; UK, the United Kingdom; USA, the United States of America; W, Woman.

† Mean age given if more than one patient included.

countries with different health systems (Table 1). Sixty-five articles were included in the quantitative synthesis.

The articles included 759 patients with a mean \pm standard deviation age of 42.4 ± 18.8 years. Sex was given for 313 patients, with a majority of women (n=196). The description of the special needs of the patients included in the studies varied (Table 1). Descriptions included physical disabilities and individuals with movement disorders, neurodegenerative diseases, learning disabilities, genetic disorders, and dental anxiety. Oral hygiene status was specified in 30 articles.^{20,25,44,45,48,49,51–54,56,58,60–62,67,77,79–83,86,90,109,110,114,117–119}

It was considered good or satisfactory for 121, and poor or inadequate for 80 patients. Support for oral hygiene was explicitly described for 32 patients. Good plaque control was a criterion for preselection of patients receiving implants in 2 studies.^{44,48} Inability to perform oral self-care was an inclusion criterion for 1 observational study.⁵¹ Parafunctional status (such as bruxism or thumb-sucking) was detailed for 18 patients, 14 of whom had parafunction.^{46,83,85} Atypical swallowing was not mentioned. Smoking status was specified for 422 patients, 11 of whom were smokers.^{52,73,76,111,115} Cooperation during dental treatment was described for 131 patients, 57 of whom were uncooperative.^{20,51,59,60,77,84,110} For patients with cognitive impairment, information about their ability to communicate was available for 37 patients and was described as difficult for 4 of them.^{59,60,79,84}

Overall, 2269 implants were placed, all endosseous titanium implants of different brands and models. The implants were equally distributed between the maxillary (n=712) and mandibular arches (n=718), although this information was not available in 4 articles^{20,21,45,50} and

the location was not specified for 2 of the 105 implants placed in 1 study.⁵⁴

Protocols for surgery, type of prosthodontic rehabilitation, and sedation are shown in *Supplemental Table 1* (available online). The surgical procedures consisted of 2-stage surgery, 1-stage surgery, or implant placement with immediate loading. If immediate loading was not used, healing times until abutment connection and prosthetic rehabilitation were reported in 45 articles^{21,25,45–47,51,53,60,61,63,65,66,68–70,74,77–80,82–84,87–90,92–94,97–103,106,108–110,112–114,117} and ranged from 1 month^{53,92} to 13 months,¹¹⁷ with a mean \pm standard deviation time of 5.2 ± 2.4 months.

Implant-supported prostheses included single crowns (SCs), fixed partial dentures (FPDs), fixed complete arch complete dentures (FCD), and overdentures (IOD).

Surgical complications were reported in 6 articles.^{46,47,57,93,105,116} Flap dehiscence with infection and implant exposure was specified,^{46,47} as were lack of primary stability,¹⁰⁵ sublingual hematoma,⁹³ temporary paraesthesia,⁵⁷ fracture of the mandible during the healing phase associated with trauma during an alteration, and upper airway obstruction.¹¹⁶

Prosthetic complications were reported in 29 articles, specifying one or more biological implant-related complications or failures (*Supplemental Table 2*). When the exact number of implants was specified, complications occurred in 308 implants. Implant fixture complications were marginal bone loss (194 implants), implant loss (90 implants), perimucositis (14 implants), peri-implantitis (5 implants), pain (4 implants), and soft tissue alterations (1 implant). Complications were identified at different stages: in the healing phase, by stage-2 surgery,

when changing the prosthesis a second time, or during an evaluation appointment. Report of complications was not always directly associated with failure, and the implants were not systematically removed.^{47,82,105} Twenty-two articles specified one or more lost implants in patients with special needs. In the 19 articles for which the information was available, 56 implants were lost before they had been loaded, and 31 implants were lost after osseointegration.

Behavioral management and anesthesia were described in 44 articles (*Supplemental Table 1*). Protocols differed depending on the treatment phase. For the implant surgery, GA was most frequently reported, followed by management under local anesthesia (LA), and then under conscious sedation (CS). Lack of cooperation during dental treatment was not the only reason to proceed with GA.^{57,64,71,109} When detailed, sedation was described as 'light'⁴⁹ or 'deep'^{53,60} without precision; as inhalation sedation with nitrous oxide;¹¹⁰ and as intravenous sedation^{20,21,25,51,80,92,103,110,113,119} with midazolam only^{20,25,92,103,110,119} or in association with pethidine⁹⁰ or nitrous oxide.¹⁰⁸ Propofol sedation was provided for a cone beam computer tomography scan and for periodontal surgery undertaken separately from the implant surgery phases.⁶⁰ When 2-stage surgery was provided, 8 articles^{45,46,65,77,79,80,84,119} specifically described that pharmacological behavioral support with GA^{45,84} or intravenous sedation^{79,80,119} was necessary to uncover the implants. Four articles described that wax occlusal registrations and impressions were made during the implant surgery with sedation.^{20,45,46,76}

Nine articles reported that pharmacological behavioral support was also required for the prosthetic phases of treatment (impression making or prosthesis placement) when GA or CS had been previously necessary for implant placement.^{21,45,46,49,51,80,84,109,119} Depending on the patient, the techniques used were GA,^{45,46,49,51,80,84,109} intravenous sedation,^{21,119} light sedation,⁴⁹ or oral premedication.²¹

Nine clinical reports did not specify follow-up, but all the implants were successfully loaded. Dental examinations were timed on an individual basis (*Supplementary Table 2*). The time between 2 appointments was generally decided by the clinician according to the patient's oral condition and was reduced if the patient had difficulty maintaining oral hygiene at home. The patients were closely monitored during the first year. Thereafter, the appointments became less frequent. They took place, or were expected to take place, every 3 months, 6 months, or annually.

The evaluation criteria of the success or failure of an implant were not always explicitly stated. The definitions and the clinical and radiological criteria were not standardized, although they were often similar (*Supplemental Table 3*). Reported survival rates^{44,45,48,49,52,54} ranged from 84.4%⁴⁵ to 100%.^{48,52,56} An estimation of the cumulative

survival rate was specified in 5 articles,^{20,47,51,52,54} ranging from 85.5%⁴⁷ to 100%⁵² at respectively 5 to 15 years⁴⁷ and 1 to 7 years⁵² after implant placement and loading. Only 2 articles^{52,56} made a clear distinction between survival rate and success rate. Whereas the authors of both articles reported a survival rate of 100%, the 3-year success rates were 93.8%⁵² and 96.1%.⁵⁶

One article specified the mastication ability and improved quality of life as reported by the patient's relatives.⁷⁹ Two articles evaluated the subjective experience of the patients using satisfaction questionnaires.^{52,118} One article investigated the oral health-related quality of life of a patient with the Oral Health Impact Profile-14.¹¹⁵ Another evaluated improvement in mastication through a quantifiable criterion of gastrointestinal deficiency.^{118,120}

The single treatment of a patient with special needs in a cohort of patients from the general population was included with those of the clinical reports.⁵⁵ Fourteen of the 79 articles included in the qualitative synthesis were excluded from the quantitative analysis, 12 of which did not report follow-up duration. Two articles were cohort studies.

Overall, 65 studies involving 666 patients with special needs and 1886 implants were included. Eighty-six implants were lost or removed. The weighted survival rate was 95.4% with a mean \pm standard deviation follow-up duration of 40.7 \pm 29.5 months.

The survival rate among patients who were able to perform oral hygiene and with no immune or metabolic bone disease (18 patients, 75 implants) was 100% for a mean \pm standard deviation follow-up time of 21.7 \pm 18.6 months.^{48,76,111,113} A 96.1% survival rate was found for the 196 patients (715 implants) needing oral hygiene support because of dexterity or cognitive impairment, but with no immune or metabolic bone disease.^{20,21,25,44,47,49,51,54,59,60,92-99,103,109,112,114,116,118,119} Their mean \pm standard deviation follow-up time was 39.5 \pm 29.8 months. A 93.9% survival rate was reported for the 425 patients (1096 implants) whose immune system was impaired and/or with a syndrome and/or medication affecting bone quality for a mean \pm standard deviation follow-up time of 43.4 \pm 29.6 months.^{44,45,50,52,53,55,58,61,64,67-70,72,74,75,78-89,91,100,101,104,105,107,108,110,117,119}

DISCUSSION

The study hypothesis that implant survival rates would be lower in patients with special needs than in the general population was rejected. The results show an encouraging implant survival rate in a population that includes individuals with dental anxiety, genetic disorders, and cognitive and/or physical impairment. This review is possibly the most exhaustive yet of the use of dental implants in




Figure 1. Preferred reporting items for systematic reviews and meta-analyses 2020 flow diagram.

patients with special needs. The population of patients with special needs is varied and difficult to classify. Patients tend to share common characteristics, including poor co-operation and barriers to oral self-care. The fact that 79 articles could be included in the qualitative synthesis is a testimony to a real and worldwide clinical interest in implant rehabilitation in special care dentistry that has steadily increased over the past decade. Nevertheless, the low number of randomized controlled clinical trials compared with the high number of case reports illustrates the challenges of undertaking clinical research into oral rehabilitation for populations with special needs. Case reports advance knowledge when little evidence exists, but they tend to lean strongly toward positive outcomes and therefore this systematic review was subject to risk of publication bias.¹²¹ Further studies with higher quality evidence are needed.

This literature review reported that, as with the general population, patients with special needs are prone to both biological and technical complications.^{122,123} The wide heterogeneity of the population included implied differences in risk factor exposure. Patients with bone pathology have additional risks for the biological complications of dental implants.¹²⁴

Previous systematic reviews and meta-analyses have focused on subgroups of patients with special needs. The reported survival rate ranged from 65% to 100% in these groups.^{29,31,35–37,39,40,43,125–133} Systematic reviews of the survival rates of dental implants in the general population show the following results: 95.6%¹³⁴ or 91.27% to 96.36% depending on the implant-supported prosthesis after 5 years.¹³⁵ The value ranges from 93.1%¹³⁴ to 94.6%¹³⁶ after 10 years. The present results showed survival rates consistent with those for patients with systemic conditions and within the range of survival rates for healthy patients. The findings suggested that complex medical conditions, cognitive and physical disabilities, and genetic disorders should not be considered immediate contraindications to oral implant rehabilitation and that implants are stable once osseointegrated.^{45,47}

Poor oral hygiene, poor cooperation, and parafunction are arguments commonly heard against implant placement in patients with special needs. However, few authors mentioned these factors, and the evident lack of available data hinders an evidence-based conclusion. All patients with dental implants should be followed up at regular intervals to monitor peri-implant status, the

condition of the implant-supported prosthesis, and plaque control.¹³⁷ Patients with special needs may have reduced capacity to express pain.¹³⁸ The awareness of their need for care may rely on the assessment of care givers.¹³⁹ Follow-up appointments therefore need to be more frequent and information needs to be provided to help carers understand, be involved, and perform oral hygiene when necessary.

The evaluation criteria for treatment outcomes differ widely in the included articles. All focused on biological instead of physiological criteria despite the fact that improving function is a primary aim when treating patients. The success and survival rates reported were related to the implant only and not to the implant-supported prosthesis. Several alternatives exist for the evaluation of oral implant rehabilitation. For individuals without cognitive impairments, psychometric evaluations can be used, and validated questionnaires of oral health-related quality of life have been used.¹⁴⁰ A tool based upon the International Classification of Functioning, Disability and Health describing the functional impact of oral health is currently being developed and could be used in further studies.^{141–144} For individuals with cognitive and/or learning disabilities, objective evaluation is needed. Mastication can be judged objectively with several validated methods,^{145,146} some of which have already been used in implant dentistry.^{147–151}

A multidisciplinary approach is necessary to assess the benefit-risk ratio of any oral rehabilitation for patients with special needs, to choose the most suitable option between implant-supported fixed prostheses, implant-supported overdentures, fixed partial dentures, removable dentures, or no rehabilitation. It has been well documented that patients with missing posterior teeth and an unstable occlusion are at higher risk of dysphagia, choking, food aspiration, and severe bronchopulmonary disease.^{152–155} These factors can only be magnified in patients with preexisting neuromotor conditions. The consequences of not providing oral rehabilitation must therefore be weighed against mental health, quality of life, and the risk of implant failure when making clinical decisions regarding patients with special needs.^{92,93,112} This assessment may be one path to ensuring more equitable oral health outcomes for patients with special healthcare needs (Fig. 1).

CONCLUSIONS

Based on the findings of this systematic literature review of protocols and outcomes, the following conclusion was drawn:

This review should enrich the discussion between professionals and patients for whom daily oral hygiene and parafunction are difficult to control

when making informed decisions between the different treatment options for oral rehabilitation.

DATA AVAILABILITY STATEMENT

The review was not registered. Protocol, critical appraisal review and data presented in this study are available on request from the corresponding author.

APPENDIX A. SUPPORTING INFORMATION

Supplemental data associated with this article can be found in the online version at [doi:10.1016/j.jprosdent.2023.09.013](https://doi.org/10.1016/j.jprosdent.2023.09.013).

REFERENCES

- Kapferer-Seebacher I, Foradori L, Zschocke J, Schilke R. Rare genetic disorders affecting the periodontal supporting tissues in adolescence. *Front Dent Med.* 2021;2:687510.
- Shimizu T, Maeda T. Prevalence and genetic basis of tooth agenesis. *Jpn Dental Sci Rev.* 2009;45:52–58.
- Peretz B, Spierer A, Spierer S, Rakoczi M. Dental treatment of patients with systemic diseases compared to patients with developmental disabilities under general anesthesia. *Spec Care Dentist.* 2012;32:21–25.
- Niesten D, van Mourik K, van der Sanden W. The impact of frailty on oral care behavior of older people: A qualitative study. *BMC Oral Health.* 2013;13:61.
- Gerritsen AE, Allen PF, Witter DJ, Bronkhorst EM, Creugers NH. Tooth loss and oral health-related quality of life: A systematic review and meta-analysis. *Health Qual Life Outcomes.* 2010;8:126.
- Naka O, Anastassiou V, Pissiotis A. Association between functional tooth units and chewing ability in older adults: A systematic review. *Gerodontology.* 2014;31(3):166–177.
- Meguro A, Ohara Y, Edahiro A, et al. Factors associated with denture non-use in older adults requiring long-term care. *Arch Gerontol Geriatr.* 2021;95:104412.
- Mac Giolla Phadraig C, Nunn J, Carroll R, McCarron M, McCallion P. Why do edentulous adults with intellectual disabilities not wear dentures? Wave 2 of the IDS TILDA cohort study. *J Prosthodont Res.* 2017;61:61–66.
- Minakuchi S, Takaoka S, Shimoyama K, Uematsu H. Factors affecting denture use in some institutionalized elderly people. *Spec Care Dentist.* 2006;26:101–105.
- Tanaka A, Kellesarian SV, Arany Szilvia. Xerostomia and patients' satisfaction with removable denture performance: Systematic review. *Quintessence Int.* 2020;52:46–55.
- Oh SH, Kim Y, Park JY, Jung YJ, Kim SK, Park SY. Comparison of fixed implant-supported prostheses, removable implant-supported prostheses, and complete dentures: Patient satisfaction and oral health-related quality of life. *Clin Oral Implants Res.* 2016;27:e31–e37.
- Mericske-Stern R. Prosthetic considerations. *Aust Dental J.* 2008;53:S49–S59.
- Wöstmann B, Budtz-Jørgensen E, Jepson N, et al. Indications for removable partial dentures: A literature review. *Int J Prosthodont.* 2005;18:139–145.
- Shaghaghian S, Taghva M, Abduo J, Bagheri R. Oral health-related quality of life of removable partial denture wearers and related factors. *J Oral Rehabil.* 2015;42:40–48.
- Batisse C, Bonnet G, Veyrune JL, Nicolas E, Bessadet M. Predictive parameters of oral health quality of life in complete mandibular denture wearers stabilized by mini-implants: A two-year follow-up study. *Materials (Basel).* 2017;10:1197.
- Heydecke G, Locker D, Awad MA, Lund JP, Feine JS. Oral and general health-related quality of life with conventional and implant dentures: Oral and general health of seniors with implant dentures. *Community Dent Oral Epidemiol.* 2003;31:161–168.
- Fillion M, Aubazac D, Bessadet M, Allègre M, Nicolas E. The impact of implant treatment on oral health related quality of life in a private dental practice: A prospective cohort study. *Health Qual Life Outcomes.* 2013;11:197.

18. Gonçalves GSY, De Magalhães KMF, Rocha EP, Dos Santos PH, Assunção WG. Oral health-related quality of life and satisfaction in edentulous patients rehabilitated with implant-supported full dentures all-on-four concept: A systematic review. *Clin Oral Invest.* 2022;26:83–94.
19. Schoen PJ, Raghoebar GM, Bouma J, et al. Prosthodontic rehabilitation of oral function in head-neck cancer patients with dental implants placed simultaneously during ablative tumour surgery: An assessment of treatment outcomes and quality of life. *Int J Oral Maxillofac Surg.* 2008;37:8–16.
20. Cune MS, Strooker H, de Putter C, Laine ML, Verhoeven JW. Dental implants in persons with severe epilepsy and multiple disabilities: A long-term retrospective study. *Int J Oral Maxillofac Implants.* 2009;24:534–540.
21. Jiménez JL, Domínguez AR, Prats MG. Implantes en pacientes discapacitados. *Med Oral.* 2003;8:288–293.
22. Ekdeldt A, Johansson LÅ, Christiansson U, et al. A retrospective analysis of factors associated with multiple implant failures in maxillae: Factors associated with multiple implant failures in maxillae. *Clin Oral Implants Res.* 2001;1:462–467.
23. Brägger U, Aeschlimann S, Bürgin W, Hämmeler CHF, Lang NP. Biological and technical complications and failures with fixed partial dentures (FPD) on implants and teeth after four to five years of function: Biological and technical complications. *Clin Oral Implants Res.* 2001;12:26–34.
24. Gläuser R, Ree A, Lundgren A, Gottlow J, Hammerle CHR, Schärer P. Immediate occlusal loading of bränemark implants applied in various jawbone regions: A prospective, 1-year clinical study. *Clin Implant Dent Relat Res.* 2001;3:204–213.
25. Packer M, Nikitin V, Coward T, Davis DM, Fiske J. The potential benefits of dental implants on the oral health quality of life of people with Parkinson's disease. *Gerodontology.* 2009;26:11–18.
26. Page MJ, Moher D, Bossuyt PM, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ.* 2021;1:160.
27. Munn Z, Barker TH, Moola S, et al. Methodological quality of case series studies: An introduction to the JBI critical appraisal tool. *JBI Evidence Synthesis.* 2020;18:2127–2133.
28. Moola S, Munn Z, Tuñanaru C, et al. Chapter 7: Systematic reviews of etiology and risk. In: Aromataris E, , Munn Z. (Editors). JBI Manual for Evidence Synthesis. JBI, 2020. Available from (<https://synthesismanual.jbi.global/>). Accessed Sept 10, 2021.
29. Srinivasan M, Meyer S, Mombelli A, Müller F. Dental implants in the elderly population: A systematic review and meta-analysis. *Clin Oral Impl Res.* 2017;28:920–930.
30. Stavropoulos A, Berti K, Pietschmann P, Pandis N, Schiødt M, Klinge B. The effect of antiresorptive drugs on implant therapy: Systematic review and meta-analysis. *Clin Oral Impl Res.* 2018;29:54–92.
31. Najeeb S, Khurshid Z, Siddiqui F, Zohaib S, Zafar MS. Outcomes of dental implant therapy in patients with down syndrome: A systematic review. *J Evid Based Dent Pract.* 2017;17:317–323.
32. Sales PH da H, Barros AWP, de Lima FJC, Carvalho A, de AT, Leão JC. Is Down syndrome a risk factor or contraindication for dental implants? A systematic review. *Int J Prosthet Dent.* 2023;128:531–537.
33. Turri A, Rossetti P, Canullo L, Grusovin M, Dahlén C. Prevalence of peri-implantitis in medically compromised patients and smokers: A systematic review. *Int J Oral Maxillofac Implants.* 2016;31:111–118.
34. Guobis Z, Pacauskiene I, Astramskaite I. General diseases influence on peri-implantitis development: A systematic review. *JOMR.* 2016;7.
35. Aghaloo T, Pi-Anfruns J, Moshaverinia A, Sim D, Grogan T, Hadaya D. The effects of systemic diseases and medications on implant osseointegration: A systematic review. *Int J Oral Maxillofac Implants.* 2019;34:s35–s49.
36. Kotsakis GA, Ioannou AL, Hinrichs JE, Romanos GE. A systematic review of observational studies evaluating implant placement in the maxillary jaws of medically compromised patients: Maxillary implants in compromised patients. *Clin Implant Dent Relat Res.* 2015;17:598–609.
37. Chrcanovic BR, Kisch J, Wennerberg A. Dental implants in patients with Sjögren's syndrome: A case series and a systematic review. *Int J Oral Maxillofac Surg.* 2019;48:1250–1259.
38. Almeida D, Vianna K, Arriaga P, Moraschini V. Dental implants in Sjögren's syndrome patients: A systematic review. *PLoS ONE.* 2017;12:e0189507.
39. Reicht PA, Schmidt-Westhausen AM, Khongkhunthian P, Strietzel FP. Dental implants in patients with oral mucosal diseases - A systematic review. *J Oral Rehabil.* 2016;43:388–399.
40. Strietzel F, Schmidt-Westhausen A, Neumann K, Reicht P, Jackowski J. Implants in patients with oral manifestations of autoimmune or mucocutaneous diseases – A systematic review. *Med Oral.* 2019;1:e217–e230.
41. Dubourg S, Huck O, Jung S. Implant-based oral rehabilitation in systemic sclerosis patients: A systematic review. *J Oral Implantol.* 2022;48:251–260.
42. Anitua E, Alkhraisat MH, Piñas L, Torre A, Eguia A. Implant-prosthetic treatment in patients with oral lichen planus: A systematic review. *Spec Care Dentist.* 2022;42:60–72.
43. Oelerich O, Kleinheinz J, Bohner L, Wiesmüller V, Hanisch M. Dental implants in people with osteogenesis imperfecta: A systematic review. *In J Environ Res Public Health.* 2022;19:1563.
44. Corcera-Flores JR, López-Giménez J, López-Jiménez J, López-Giménez A, Silvestre-Rangil J, Machuca-Portillo G. Four years survival and marginal bone loss of implants in patients with Down syndrome and cerebral palsy. *Clin Oral Invest.* 2017;21:1667–1674.
45. De Bruyn H, Glibert M, Matthijs L, Filipi M, Christiaens V, Marks L. Clinical guidelines for implant treatment in patients with Down syndrome. *Int J Periodontics Restorative Dent.* 2019;39:361–368.
46. Ekdeldt A. Early experience of implant-supported prostheses in patients with neurologic disabilities. *Br Dent J.* 2005;199: 151–151.
47. Ekdeldt A, Zellmer M, Carlsson G. Treatment with implant-supported fixed dental prostheses in patients with congenital and acquired neurologic disabilities: A prospective study. *Int J Prosthodont.* 2013;26:517–524.
48. Enkling N, Hardt K, Katsoulis J, et al. Dental phobia is no contraindication for oral implant therapy. *Quintessence Int.* 2013;44:363–371.
49. Feijoo JF, Limeres J, Diniz M, Del Llano A, Seoane J, Díz P. Osseointegrated dental implants in patients with intellectual disability: A pilot study. *Disabil Rehabil.* 2012;34:2025–2030.
50. Jagadeesh KN, Parihar AS, Verma AK, Das AC, Razi MA. Assessment of the survival rate of short dental implants in medically compromised patients. *J Contemp Dent Pract.* 2020;21:880–883.
51. Kim I hyung, Kuk TS, Park SY, Choi Y suk, Kim HJ, Seo KS. Prognosis following dental implant treatment under general anesthesia in patients with special needs. *J Dent Anesth Pain Med.* 2017;17:205.
52. Krennmaier G, Seemann R, Piehslinger E. Dental implants in patients with rheumatoid arthritis: Clinical outcome and peri-implant findings. *J Clin Periodontol.* 2010;37:928–936.
53. Limeres Posse J, López Jiménez J, Ruiz Villandiego JC, et al. Survival of dental implants in patients with Down syndrome: A case series. *J Prosthetic Dent.* 2016;116:880–884.
54. Özcakir C, Balmer S, Mericske-Stem R. Implant-prosthetic treatment for special care patients: A case-series study. *Br Dent J.* 2006;200: 145–145.
55. Van de Velde T, Collaert B, De Bruyn H. Immediate loading in the completely edentulous mandible: technical procedure and clinical results up to 3 years of functional loading. *Clin Oral Implants Res.* 2007;18:295–303.
56. Weinländer M, Krennmaier G, Piehslinger E. Implant prosthodontic rehabilitation of patients with rheumatic disorders: A case series report. *Int J Prosthodont.* 2009;23:7.
57. Apaydin A, Sermet B, Ureturk S, Kundakcioglu A. Correction of malocclusion and oral rehabilitation in a case of amelogenesis imperfecta by insertion of dental implants followed by Le Fort I distraction osteogenesis of the edentulous atrophic maxilla. *BMC Oral Health.* 2014;14:116.
58. Barker D, Nohl FSA, Postlethwaite KR, Smith DG. Case report of multiple implant failure in a patient with ankylosing spondylitis. *Eur J Prosthodont Restor Dent.* 2008;16:20–23.
59. Rogers JO. Implant-stabilized complete mandibular denture for a patient with cerebral palsy. *Dent Update.* 1995;23–26.
60. Hong YJ, Dan JB, Kim MJ, Kim HJ, Seo KS. Prognosis after treatment with multiple dental implants under general anesthesia and sedation in a cerebral palsy patient with mental retardation: A case report. *J Dent Anesth Pain Med.* 2017;17:149.
61. Eder A, Watzek G. Treatment of a patient with severe osteoporosis and chronic polyarthritides with fixed implant-supported prosthesis: A Case report. *Int J Oral Maxillofac Implants.* 1999;14:587–590.
62. Ambard AJ, Clemens S, Phillips DS. Multidisciplinary implant rehabilitation of a patient with cleidocranial dysostosis: A journey from age 13 to 21. *J Prosthodont.* 2019;28:361–364.
63. Angle AD, Rebello J. Dental team management for a patient with cleidocranial dysostosis. *Am J Orthod Dentofacial Orthop.* 2005;128:110–117.
64. Ahmad AG, Osman M, Awadalkreem F. Full-mouth rehabilitation of a patient with cleidocranial dysplasia using immediately loaded basal implant-supported fixed prostheses: A case report. *Int J Surg Case.* 2019;65:344–348.
65. Atıl F, Culhaoglu A, Kocigit ID, Adisen Z, Misirlioglu M, Yilmaz B. Oral rehabilitation with implant-supported fixed dental prostheses of a patient with cleidocranial dysplasia. *J Prosthet Dent.* 2018;119:12–16.
66. Calvo-Guirado JL, Ramos-Oltra ML, Negri B, et al. Platform switching in the treatment of cleidocranial dysplasia: A case report. *Journal of Osseointegration.* 2013;5:27–30.
67. Daskalogiannakis J, Piedade L, Lindholm TC, Sándor GKB, Carmichael RP. Cleidocranial dysplasia: 2 generations of management. *J Can Dent Assoc.* 2006;72:337–342.
68. Lee C, Jung H sup, Baek JA, Leem DH, Ko SO. Manifestation and treatment in a cleidocranial dysplasia patient with a RUNX2 (T420I) mutation. *Maxillofac Plast Reconstr Surg.* 2015;37:41.
69. Lombardas P, Toothaker RW. Bone grafting and osseointegrated implants in the treatment of cleidocranial dysplasia. *Compend Contin Educ Dent.* 1997;18:509–514.

70. Noh K, Kwon KR, Ahn H, Paek J, Pae A. Prosthetic rehabilitation of a cleidocranial dysplasia patient with vertical maxillofacial deficiency: A clinical report: Prosthetic Rehabilitation of a CCD Patient. *J Prosthodont.* 2014;23:64–70.
71. Petropoulos VC, Balshi TJ, Balshi SF, Wolfinger GJ. Treatment of a patient with cleidocranial dysplasia using osseointegrated implants: A patient report. *Int J Oral Maxillofac Implants.* 2004;19:282–287.
72. Petropoulos VC, Balshi TJ, Wolfinger GJ, Balshi SF. Treatment of a patient with cleidocranial dysplasia using a single-stage implant protocol. *Treatment of a Patient with Cleidocranial Dysplasia.* *J Prosthodont.* 2011;20:S26–S31.
73. Schnuttenhaus S, Luthardt RG, Rudolph H, Götz W. Histological examination and clinical evaluation of the jawbone of an adult patient with cleidocranial dysplasia: A case report. *Int J Clin Exp Pathol.* 2015;8:8521–8531.
74. Schnuttenhaus S, Götz W, Luthardt RG. Prosthetic rehabilitation of a patient with cleidocranial dysplasia using dental implants—A clinical report. *Int J Implant Dent.* 2021;7:5.
75. Tan MH, Seah TE, Goh BT. Staged orthodontic, orthognathic and open septorhinoplasty for management of cleidocranial dysplasia: A case report. *Int J Oral and Maxillofac Surg.* 2019;48:26.
76. Balshi TJ, Wolfinger GJ. Immediate placement and implant loading for expedited patient care: A patient report. *Int J Oral Maxillofac Implants.* 2002;17:587–592.
77. Alqahtani NM, Alsayed HD, Levon JA, Brown DT. Prosthodontic rehabilitation for a patient with Down syndrome. *J Prosthodont.* 2018;27:681–687.
78. Altintas NY, Kilic S, Altintas SH. Oral rehabilitation with implant-retained overdenture in a patient with Down syndrome : Implant therapy in a patient with Down syndrome. *J Prosthodont.* 2019;28:e617–e621.
79. Comparin LL, Filho RP, Deliberador T, et al. Sinus lift technique and dental implants for rehabilitation for a Down syndrome patient with tomographic 3-years follow-up. *Spec Care Dentist.* 2022;42:524–530.
80. Lustig JP, Yanko R, Zilberman U. Use of dental implants in patients with Down syndrome: A case report. *Spec Care Dentist.* 2002;22:201–204.
81. Reiche W, Parkanyi L, Braunitzer G, Nagy K, Karpati K. Oligodontia in a teenager with Down syndrome – A case study. *Klin Padiatr.* 2014;227:35–37.
82. Ribeiro CG, Siqueira AF, Bez L, Cardoso AC, Ferreira CF. Dental implant rehabilitation of a patient with Down syndrome: A case report. *J Oral Implantol.* 2011;37:481–487.
83. Saponaro PC, Deguchi T, Lee DJ. Implant therapy for a patient with Down syndrome and oral habits: A clinical report. *J Prosthet Dent.* 2016;116:320–324.
84. Schmidt P, Goedicke-Padigur G, Schulte AG, Benz K, Jackowski J. Implant-supported prosthetic rehabilitation of a senescent patient with Down syndrome. *Quintessence Int.* 2020;51:170–177.
85. Soares MRPS, Paula FO de, Chaves M das GAM, Assis NM de SP, Chaves Filho HD de M. Patient with Down syndrome and implant therapy: A case report. *Braz Dent J.* 2010;21:550–554.
86. Thomas S, Simons D. Implant, surgical and prosthodontic treatment for a patient with Down syndrome - a case report. *J Disability Oral Health.* 2011;12:91–94.
87. Brahim J, Guckles A, Ruby S. Implant Rehabilitation in Erdheim-Chester Disease: A Clinical Report. *J Prosthet Dent.* 1992;68:399–401.
88. Engel JD, Ruskin JD, Tu HK. Hematologic management of a patient with Fanconi's anemia undergoing bone grafting and implant surgery. *J Oral Maxillofac Surg.* 1992;50:288–292.
89. Gallottini M, Llanos AH, Romito GA, Romano MM, de Oliveira FB, de Rezende NPM. Oral manifestations and rehabilitation in Fraser syndrome: A case report: Oral manifestations and rehabilitation. *Spec Care Dentist.* 2018;38:249–254.
90. Güven O, Tekin U, Hatipoğlu M. Surgical and prosthodontic rehabilitation in a patient with Freeman-Sheldon syndrome. *J Craniofac Surg.* 2010;21:1571–1574.
91. Merlini A, Garibaldi J, Giorgi L, Balbi P. Gnathodiaphyseal dysplasia: Surgical treatment and prosthetic rehabilitation of 2 members of the same family. *J Oral Maxillofac Surg.* 2016;74:2441–2446.
92. Deniz E, Kokat AM, Noyan A. Implant-supported overdenture in an elderly patient with Huntington's disease: Oral rehabilitation in Huntington's disease. *Gerodontology.* 2011;28:157–160.
93. Jackowski J, Dent DM, Andrich J, et al. Implant-supported denture in a patient with Huntington's disease: Interdisciplinary aspects. *Spec Care Dentist.* 2001;21:15–20.
94. Peñarrocha M, Sanchis JM, Rambla J, Guarinos J. Oral rehabilitation using osseointegrated implants in a patient with idiopathic torsion dystonia. *Int J Oral Maxillofac Implants.* 2001;16:433–435.
95. Károlyházy K, Schmidt P, Bogdán S, Hermann P, Arányi Z. Prosthodontic treatment of an edentulous epileptic patient with an implant-retained overdenture. A case report. *Ideggyógy Sz.* 2014;67:342–346.
96. Shek JW, Plesh O, Curtis DA. Immediately loaded implants in a patient with involuntary mandibular movements: A clinical report. *J Prosthet Dent.* 2014;112:14–17.
97. Oelgiesser D, Levin L, Elpeleg O, Schwartz-Arad D. Posttraumatic dental implant placement in a patient with maple syrup urine disease. *Implant Dent.* 2006;15:143–147.
98. Poluha RL, Eidl JMS, Danieleto-Zanna CF, Ferreira GZ, Takizawa OK, Farah GJ. Use of osseointegrated implants in the intermaxillary suture in a patient with microstomia: Case report with a 2-year follow-up. *Spec Care Dentist.* 2019;39:533–537.
99. Peñarrocha M, Sanchis JM, Rambla J, Sánchez MA. Oral rehabilitation with osseointegrated implants in a patient with oromandibular dystonia with blepharospasm (Brueghel's syndrome): A patient history. *Int J Oral Maxillofac Implants.* 2001;16:115–117.
100. Ullbro C, Crossner CG, Lundgren T, Stålblad PÅ, Renvert S. Osseointegrated implants in a patient with Papillon-Lefèvre syndrome: A 4½-year follow-up. *J Clin Periodontol.* 2000;27:951–954.
101. Woo I, Brunner DP, Yamashita DDR, Le BT. Dental implants in a young patient with Papillon-Lefèvre syndrome: A case report. *Implant Dent.* 2003;12:140–144.
102. Applebaum G, Langsam B, Huba G. The implant retained UCLA-type clip bar overdenture. *Oral Health.* 1997;65–72.
103. Chu FCS, Deng FL, Siu ASC, Chow TW. Implant-tissue supported, magnet-retained mandibular overdenture for an edentulous patient with Parkinson's disease: A clinical report. *J Prosthet Dent.* 2004;91:219–222.
104. Friberg B. Treatment with dental implants in patients with severe osteoporosis: A case report. *Int J Periodontics Restorative Dent.* 1994;14:348–353.
105. Bencharit S, Reside GJ, Howard-Williams EL. Complex prosthodontic treatment with dental implants for a patient with polymyalgia rheumatica: A clinical report. *Int J Oral Maxillofac Implants.* 2010;25:1241–1245.
106. Abdel-Kader MA, Abdelazeem AF, Ahmed NEB, Khalil YM, Mostafa MI. Oral rehabilitation of a case with regional odontodysplasia using a regenerative approach—A case report and a review of literature. *Spec Care Dentist.* 2019;39:330–339.
107. Gaur V, Singh N, Doshi AG, Chandradas B. Immediate rehabilitation of a rheumatoid arthritis patient with single-piece implants. *Int J Surg Case Rep.* 2021;82:105874.
108. Ella B, Lasserre JF, Blanchard JP, Fricain JC. A 4-year follow-up of two complete mandibular implant-supported removable prostheses in a patient with severe rheumatoid polyarthritis: Case report. *Int J Oral Maxillofac Implants.* 2011;26:e19–e22.
109. Choi Y suk, Kim H, Rhee SH, et al. Multiple implant therapy with multiple inductions of general anesthesia in non-compliant patients with schizophrenia: A case report. *J Dent Anesth Pain Med.* 2019;19:239.
110. Tizzoni R, Veneroni L, Clerici CA. A patient with meningeal melanomatosis treated for periodontal disease with a bone regeneration procedure and dental implants: Clinical and behavioral management to support medical compliance. *Oral Implantol ((Rome)).* 2014;6:75–81.
111. Thor A, Stenport VF, Larsson A, Bostrom A. Bone graft and implants in a patient with systemic mastocytosis. *Clin Implant Dent Rel Res.* 2005;7:79–86.
112. Kelleher M, Scott B, Djemal S. Case report: Complications of rehabilitation using osseointegrated implants - Tardive dyskinesia. *Eur J Prosthodont Rest Dent.* 1998;6:133–136.
113. Misch CM, Jolly RL, Williams DR, Chorazy CJ. Maxillary implant surgery on a patient with thalassemia. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1998;86:401–405.
114. Flanagan D. Oral Implantology for the traumatic brain injured patient: A case report. *J Oral Implantol.* 2011;37:713–716.
115. Takeuchi S, Planerova A, Malmstrom H, Saunders R. Improving oral health-related quality of life for a traumatic brain injury patient: A case report. *Spec Care Dentist.* 2019;39:617–623.
116. Durham TM, King T, Salinas T, Franco T, Ross J. Dental implants in edentulous adults with cognitive disabilities: Report of a pilot project. *Spec Care Dentist.* 2006;26:40–46.
117. Friberg B. Bränemark system implants and rare disorders: A report of six cases. *Int J Periodontics Restorative Dent.* 2013;33:139–148.
118. Heckmann SM, Heckmann JG, Weber HP. Clinical outcomes of three Parkinson's disease patients treated with mandibular implant overdentures: Outcome of 3 PD patients treated with mandibular overdentures. *Clin Oral Implants Res.* 2000;11:566–571.
119. Owens L, Claffey N, O'Sullivan M, Houston F, Nunn J. The placement of implants in patients who are medically or intellectually compromised. A review of the literature and case reports. *J Disability Oral Health.* 2009;10:107–114.
120. Hoehn MM, Yahr MD. Parkinsonism: onset, progression, and mortality. *Neurology.* 1967;17: 427–427.
121. Murad MH, Sultan S, Haffar S, Bazerbachi F. Methodological quality and synthesis of case series and case reports. *BMJ Evid Based Med.* 2018;23:60–63.

122. Sakka S, Baroudi K, Nassani MZ. Factors associated with early and late failure of dental implants. *J Investig Clin Dent.* 2012;3:258–261.
123. Pjetursson BE, Heimisdottir K. Dental implants - Are they better than natural teeth? *Eur J Oral Sci.* 2018;126(Suppl 1):81–87.
124. Ngeow WC, Lim D, Tan CC, Shetty N, Marla V. Dental implant modifications for medically compromised patients. *Dental Implants.* Elsevier; 2020:255–286.
125. Schimmel M, Srinivasan M, McKenna G, Müller F. Effect of advanced age and/or systemic medical conditions on dental implant survival: A systematic review and meta-analysis. *Clin Oral Impl Res.* 2018;29:311–330.
126. Singh K, Rao J, Afsheen T, Tiwari B. Survival rate of dental implant placement by conventional or flapless surgery in controlled type 2 diabetes mellitus patients: A systematic review. *Indian J Dent Res.* 2019;30:600.
127. Bornstein MM, Cionca N, Mombelli A. Systemic conditions and treatments as risks for implant therapy. *Int J Oral Maxillofac Implants.* 2009;24(Suppl):12–27.
128. Kim SH, Oh NS, Kim HJ. Survival rates and clinical outcomes of implant overdentures in old and medically compromised patients. *Int J Environ Res Public Health.* 2022;19:11571.
129. Sivakumar I, Arunachalam S, Choudhary S, Buzayan MM. Does HIV infection affect the survival of dental implants? A systematic review and meta-analysis. *J Prosthet Dent.* 2021;125:862–869.
130. de Medeiros FCFL, Kudo GAH, Leme BG, et al. Dental implants in patients with osteoporosis: A systematic review with meta-analysis. *Int J Oral and Maxillofac Surg.* 2018;47:480–491.
131. Esimekara JFO, Perez A, Courvoisier DS, Scolozzi P. Dental implants in patients suffering from autoimmune diseases: A systematic critical review. *J Stomatol Oral Maxillofac Surg* 2022;S2468785522000052.
132. Toneatti DJ, Graf RR, Burkhard JP, Schaller B. Survival of dental implants and occurrence of osteoradionecrosis in irradiated head and neck cancer patients: A systematic review and meta-analysis. *Clin Oral Invest.* 2021;25:5579–5593.
133. Wang Y, He J, Decker AM, Hu JC, Zou D. Clinical outcomes of implant therapy in ectodermal dysplasia patients: A systematic review. *Int J Oral Maxillofac Surg.* 2016;45:1035–1043.
134. Pjetursson BE, Thoma D, Jung R, Zwahlen M, Zembic A. A systematic review of the survival and complication rates of implant-supported fixed dental prostheses (FDPs) after a mean observation period of at least 5 years. *Clin Oral Implants Res.* 2012;23:22–38.
135. Muddugangadhar BC, Amarnath GS, Sonika R, Chheda PS, Garg A. Meta-analysis of failure and survival rate of implant-supported single crowns, fixed partial denture, and implant tooth-supported prostheses. *J Int Oral Health.* 2015;7:11–17.
136. Moraschini V, Poubel LA, da C, Ferreira VF, Barboza E, dos SP. Evaluation of survival and success rates of dental implants reported in longitudinal studies with a follow-up period of at least 10 years: A systematic review. *Int J Oral and Maxillofac Surg.* 2015;44:377–388.
137. Cohen RE. Research, science and therapy committee. Position paper: Periodontal maintenance. *J Periodontol.* 2003;74:1395–1401.
138. Hennequin M, Faulks D, Roux D. Accuracy of estimation of dental treatment need in special care patients. *J Dent.* 2000;28:131–136.
139. Brahm CO, Klingberg G, Efkeldt A. Prosthodontic treatment of patients with disabilities at dental specialist clinics in the County of Västra Götaland. *Swed Dent J.* 2009;33:11–18.
140. Sidenö L, Hmaidouch R, Brandt J, von Krockow N, Weigl P. Satisfaction level in dental-phobic patients with implant-supported rehabilitation performed under general anaesthesia: A prospective study. *BMC Oral Health.* 2018;18:182.
141. World Health Organization, ed. International Classification of Functioning, Disability and Health: ICF. World Health Organization; 2001.
142. Dougal A, Martinez Pereira F, Molina G, Eschevins C, Daly B, Faulks D. Identifying common factors of functioning, participation and environment amongst adults requiring specialist oral health care using the International Classification of Functioning, disability and health. Jan YK, ed. *PLoS ONE* 2018;13:e0199781.
143. Faulks D, Scambler S, Daly B, Jamieson L, Hennequin M, Tsakos G. Measuring oral health - How can the International Classification of Functioning help? *Community Dent Oral Epidemiol.* 2023;51:153–164.
144. Faulks D., Norderyd J., Molina G., et al. Using the International Classification of Functioning, Disability and Health (ICF) to Describe Children Referred to Special Care or Paediatric Dental Services. *Glogauer M, ed. PLoS ONE* 2013;8:e61993.
145. Van Der BILT A. Assessment of mastication with implications for oral rehabilitation: A review: Mastication and oral rehabilitation. *J Oral Rehabil.* 2011;38:754–780.
146. Schimmel M, Christou P, Miyazaki H, Halazonetis D, Hermann FR, Müller F. A novel colourimetric technique to assess chewing function using two-coloured specimens: Validation and application. *J Dent.* 2015;43:955–964.
147. Boven GC, Raghoebar GM, Vissink A, Meijer HJA. Improving masticatory performance, bite force, nutritional state and patient's satisfaction with implant overdentures: A systematic review of the literature. *J Oral Rehabil.* 2015;42:220–233.
148. Gonçalves TMSV, Campos CH, Rodrigues Garcia RCM. Mastication and jaw motion of partially edentulous patients are affected by different implant-based prostheses. *J Oral Rehabil.* 2014;41:507–514.
149. Lundqvist S. Speech and other oral functions. Clinical and experimental studies with special reference to maxillary rehabilitation on osseointegrated implants. *Swed Dent J Suppl.* 1993;91:1–39.
150. Veyrune JL, Opé S, Nicolas E, Woda A, Hennequin M. Changes in mastication after an immediate loading implantation with complete fixed rehabilitation. *Clin Oral Invest.* 2013;17:1127–1134.
151. Tan D, Foster S, Korgaonkar MS, Oxenham V, Whittle T, Klineberg I. The role of progressive oral implant rehabilitation in mastication, cognition and oral health-related quality of life outcomes—A pilot to define the protocol. *J Oral Rehabil.* 2020;47:1368–1381.
152. Faulks D, Veyrune JL, Hennequin M. Consequences of oral rehabilitation on dyskinesia in adults with Down's syndrome: A clinical report. *J Oral Rehabil.* 2002;29:209–218.
153. Binkley CJ, Haugh GS, Kitchens DH, Wallace DL, Sessler DI. Oral microbial and respiratory status of persons with mental retardation/intellectual and developmental disability: An observational cohort study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2009;108:722–731.
154. Tamura F, Mizukami M, Ayano R, Mukai Y. Analysis of feeding function and jaw stability in bedridden elderly. *Dysphagia.* 2002;17:235–241.
155. Terpenning MS, Taylor GW, Lopatin DE, Kerr CK, Dominguez BL, Loesche WJ. Aspiration pneumonia: Dental and oral risk factors in an older veteran population. *J Am Geriatr Soc.* 2001;49:557–563.

Corresponding author:

Dr Marie-Sophie Bogner
Centre for Research in Clinical Dentistry (CROC)
University of Clermont Auvergne
2 Rue Braga
Clermont-Ferrand 63000
FRANCE
Email: msj.bogner@gmail.com.

CRediT authorship contribution statement

All listed authors made considerable and valuable contributions to the development of the manuscript. **Martine Hennequin** conceived the design and idea. **Marie-Sophie Bogner** and **Vivien Chambas** performed the literature search and carried out data collection. **Marie-Sophie Bogner**, **Vivien Chambas**, **Martine Hennequin** and **Denise Faulks** were involved in drafting the manuscript. **Marie-Sophie Bogner** and **Emmanuel Nicolas** participated in meta-analysis. **Marie-Sophie Bogner**, **Martine Hennequin** and **Denise Faulks** performed a critical revision of the article for important intellectual content. All authors read the final manuscript and approved it.

Copyright © 2023 by the Editorial Council of *The Journal of Prosthetic Dentistry*.
All rights reserved.
<https://doi.org/10.1016/j.jprosdent.2023.09.013>