



**HAL**  
open science

## FAAH inhibitors in the limelight, but regrettably

Christophe Mallet, Claude Dubray, Christian Dualé

► **To cite this version:**

Christophe Mallet, Claude Dubray, Christian Dualé. FAAH inhibitors in the limelight, but regrettably. *International Journal of Clinical Pharmacology and Therapeutics*, 2016, 54 (07), pp.498 - 501. 10.5414/cp202687 . hal-04337687

**HAL Id: hal-04337687**

**<https://uca.hal.science/hal-04337687v1>**

Submitted on 12 Dec 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



©2016 Dustri-Verlag Dr. K. Feistle  
ISSN 0946-1965

DOI 10.5414/CP202687  
e-pub: May 18, 2016

# FAAH inhibitors in the limelight, but regrettably

Christophe Mallet<sup>1,2</sup>, Claude Dubray<sup>1,2,3</sup>, and Christian Dualé<sup>1,3</sup>

<sup>1</sup>Inserm U1107 «Neuro-Dol», <sup>2</sup>Univ-Clermont1, and <sup>3</sup>CHU Clermont-Ferrand, Centre de Pharmacologie Clinique (Inserm CIC1405), Clermont-Ferrand, France

## Key words

fatty acid amide hydroxylase – endocannabinoids – phase I – serious adverse event – brain injury

**Abstract.** This short review focuses on the recent drug development of FAAH inhibitors, as recent serious adverse events have been reported in a phase I study with a compound of this class. The authors overview the potential interest in targeting FAAH inhibition, the current programs, and the available information on the recent dramatic events.

In January 2016, we were informed of the exceptional occurrence of serious adverse events (SAEs) in a phase I clinical trial conducted by the Biotrial Pharmacology Center (Rennes, France) on behalf of Bial-Portela & Ca. SA (São Mamede do Coronado, Portugal). The trial involved the compound BIA 10-2474, a drug designed to inhibit fatty acid amide hydrolase (FAAH). After two initial phases (single escalating doses up to 100 mg, and kinetics-food interaction studies) without any untoward SAE, the phase in question, which intended to examine the effect of multiple doses (5 or 6 daily doses), resulted in SAEs in 6 participants, who had all been administered the highest tested dose (50 mg). This was a threshold effect, since no SAE had been reported with the lower dose of 20 mg given to the volunteers previously. The most serious symptoms had central neurological features, the worst being those associated with a single case of coma which rapidly lead to brain death. Of the other 5 hospitalized participants, 2 had serious neurological damage (with clinical improvement apparently occurring within a few days). Because of these events, the trial was immediately suspended. Further information (including the protocol of the trial) is available on the website of the French National Agency for Medicine and Health Product Safety (ANSM)<sup>1</sup>. The agency has also recently published the summarized

conclusions of a temporary specialized scientific committee<sup>2</sup>. Although no definitive conclusion can be drawn at the present time regarding the causes, the seriousness of the events has undoubtedly and, in this case regrettably, brought the development of FAAH inhibitors/inactivators into the limelight.

FAAH inhibitors/inactivators have been developed because of their ability to increase the concentration of endocannabinoids. Endocannabinoids are lipid mediators released on demand from membrane phospholipid precursors. Their targets are the cannabinoid receptors CB<sub>1</sub> and CB<sub>2</sub>, but other receptors can be involved in their action, such as GPR<sub>55</sub>, peroxisome proliferator-activated receptors (PPARs) and vanilloid receptors (TRPV<sub>1</sub>). This system has been implicated in a wide range of physiological processes such as those associated with chronic pain, metabolic disorders, psychoses, nausea and vomiting, depression, and anxiety disorders (see [1, 2, 3, 4, 5] for reviews). Some exogenous cannabinoids acting on CB<sub>1-2</sub> are currently used in therapeutics (e.g., Bedrocane<sup>®</sup>, Bedrobinol<sup>®</sup>, Bediol<sup>®</sup>, Bedica<sup>®</sup>, Cesamet<sup>®</sup>, Marinol<sup>®</sup>, Sativex<sup>®</sup>) involving a variety of indications such as anorexia, neuropathic pain and multiple sclerosis, depending on the country in which the drugs are marketed. However, such treatments may have neurological side effects (including impairment of cognition and motor functions and a predisposition to psychoses), notably when these agents are used for long-term treatment [6, 7].

Received  
April 7, 2016;  
accepted  
April 13, 2016

Correspondence to  
Dr. Christian Dualé  
CHU Clermont-Ferrand,  
Centre de Pharmacologie  
Clinique, 28, Place  
Henri-Dunant BP 38,  
63001 Clermont-Ferrand,  
France  
cduale@chu-clermont-  
ferrand.fr

<sup>1</sup><http://ansm.sante.fr/S-informer/Actualite/Essai-clinique-BIA-102474-101-Bial-Biotrial-Chronologie>

<sup>2</sup>[http://ansm.sante.fr/Dossiers/Essai-Clinique-Bial-Biotrial/Essai-clinique-BIA-102474-101-du-laboratoire-BIAL/\(offset\)/0](http://ansm.sante.fr/Dossiers/Essai-Clinique-Bial-Biotrial/Essai-clinique-BIA-102474-101-du-laboratoire-BIAL/(offset)/0)

Increasing the concentration of endocannabinoids, rather than administering exogenous agonistic agents, would reduce cannabinoid-like adverse events. This strategy can be achieved through the inhibition of catabolic enzymes, notably FAAH, an integral membrane enzyme that hydrolyzes the fatty amide family of lipid transmitters including the most thoroughly studied endocannabinoid, N-arachidonylethanolamide (anandamide) [8]. FAAH also degrades several related fatty acid amides which have diverse biological functions and mechanisms of action [9]. FAAH-deficient mice have enhanced levels of anandamide and display a CB<sub>1</sub> receptor-mediated hypoalgesic phenotype [10, 11]. Pharmacological inhibition of FAAH increases fatty acid amide concentrations in both rats and humans [12, 13]. This strategy seems to be successful in animal models of anxiety and depression [3], sleep disorders [14], and nociceptive or neuropathic pain [12, 15, 16]. Interest in this pharmacological pathway is illustrated by the numerous molecules under development. Among the most advanced programs are those concerning the compounds PF-04457845, JNJ-42165279, SSR-411298, V-158866, and URB597<sup>3</sup>. The disorders for which these agents are being tested are mostly neuropsychiatric, such as pain conditions, depression, anxiety disorders, and phobias, Tourette syndrome, and symptoms associated with cannabis withdrawal. The available data from completed clinical trials indicate that FAAH inhibitors are well tolerated. A phase I study of PF-04457845 (developed by Pfizer, New York, NY, USA) showed that, compared to placebo, the increase in somnolence was only mild, and that there were no effects on cognitive function [13]. The corresponding phase II study demonstrated that this agent had a safety profile indistinguishable from placebo, where the main treatment-related side effect was dizziness [17]. A phase I study of JNJ-42165279 (developed by Johnson & Johnson Pharmaceutical, New York, NY, USA) found few side effects and all were of mild intensity. A slight and transient

increase in liver transaminases was observed at the highest doses in a few cases [18]. In a phase II study with SSR-411298 (Sanofi, Gentilly, France) in patients with depression, the rate of adverse events was similar in the treatment and the placebo group. Headache, suicidal ideation, diarrhea, dizziness, and nausea were the most frequent adverse effects observed in the treatment group [19]. A major observation was that FAAH inhibitors do not apparently induce those adverse effects commonly associated with exogenous cannabinoids, such as impairment in cognition, motor coordination, and psychoses. However, it must be noted that the effects of chronic treatment are still largely unknown. It is also of interest that in two randomized, double-blind, placebo- and active drug-controlled clinical trials, no clinical efficacy of the tested FAAH inhibitor could be demonstrated. This was the case with PF-04457845 on pain due to osteoarthritis [17], and with SSR-411298 in elderly patients with major depressive disorders [19]. Following announcement of the accident at Biotrial, the two JNJ-42165279 trials were also suspended as a precautionary measure.

Since these events came to light, many researchers and clinicians have questioned whether the type of adverse events observed are directly related to an inhibition of FAAH, and speculate that they may result from an interaction with other pharmacological targets, specific to BIA 10-2474 and which only become important above a certain concentration threshold. An answer to this question is highly important since on the one hand, this type of adverse reaction could potentially occur with all substances belonging to this class of drugs but on the other, this would not apply if a different target, specific to this particular drug was involved. The putative role of a metabolite has also been mentioned by the experts commissioned by the ANSM. They have since rejected the hypothesis of a manufacturing defect, and state that the preclinical studies – including that in which dogs died – had met the prerequisites for a first-in-man study.

The protocol document of the above mentioned phase I study of BIA 10-2474 shows that the study design is conventional and was apparently strictly adhered to. However, more precise information concerning

<sup>3</sup>See refs. on Clinicaltrials.gov: NCT02134080, NCT01618656, NCT00981357, NCT02498392, NCT02432703, NCT00822744, NCT01439919, NCT01748695 & NCT00916201.

the extent of the brain damage in the healthy study volunteers who suffered neurological symptoms in the days following the first 5 or 6 BIA 10-2474 administrations is still awaited. The short report which has so far been published by ANSM mentions symptoms related strictly to cerebral functions, with “very fast appearance”, with homogeneous aspects but where there are between-subject differences in severity and progression. MRI observations showed the presence of “micro-cerebral tissue damage of varying severity and with unusual topography”. On the other hand, this report has not provided a credible lead concerning possible mechanisms involved.

The fact that the adverse events were observed in 5 out of the 6 subjects who received 50 mg per day of BIA 10-2474 excludes the possibility of an idiosyncratic reaction in an individual with a particular phenotype. Thus, a concentration-dependent effect, with some between-subject variability, seems more likely. In animals, the pharmacokinetic half-life is relatively long and the fact that the accident happened after several administrations, argues for a product accumulation in which toxic levels are reached within a few days. This may result from low clearance rates and elimination pathways in humans that differ from those in the experimental animals used during product development. Although this hypothesis cannot be excluded, it does not appear to be a likely explanation because BIA 10-2474 does not interact directly with receptors, but rather inhibits an enzyme involved in the catabolism of endocannabinoids. In biological systems, inhibition of one metabolic pathway is often compensated by activation of alternative pathways in order to restore or at least partially restore the substrate balance and status quo. In support of this viewpoint is the fact that phenomena resembling those seen in humans with BIA 10-2474 have not been reported in the literature in any of the numerous animal studies with various inhibitors of FAAH, even when high doses have been used or, as in the case of rodents, when endocannabinoids have been administered by intracerebral injection. That FAAH-deficient mice (knock-out) thrive and exhibit no abnormal behavioral symptoms supports the hypothesis even though not yet conclusive, that BIA 10-2474 itself or a

metabolite thereof and not the inhibition of FAAH is the cause of the adverse reactions. The analysis of BIA 10-2474 plasma concentrations in the healthy subjects who were included in the repeated-dose study should provide information important for answering two fundamental questions, namely: (i) Does drug or metabolite accumulation occurring in humans result from low clearance or the presence of saturable metabolic pathways? (ii) Is there a relationship between the magnitude of blood concentrations and the severity of neurological symptoms in subjects who received the highest dose of 50 mg? In the course of the coming weeks or months it is hoped that information will come to light enabling a distinction to be made between the possible target specificities of BIA 10-2474 and those of other FAAH inhibitors. This information is vitally important since the fate of this class of drug and the marketing potential of FAAH-inhibitors depend on it.

## Conflict of interest

The authors have no conflict of interests to declare.

## References

- [1] Battista N, Di Tommaso M, Bari M, Maccarrone M. The endocannabinoid system: an overview. *Front Behav Neurosci.* 2012; 6: 9. [CrossRef PubMed](#)
- [2] Pertwee RG. Elevating endocannabinoid levels: pharmacological strategies and potential therapeutic applications. *Proc Nutr Soc.* 2014; 73: 96-105. [CrossRef PubMed](#)
- [3] Fowler CJ. The potential of inhibitors of endocannabinoid metabolism as anxiolytic and antidepressive drugs – A practical view. *Eur Neuropsychopharmacol.* 2015; 25: 749-762. [CrossRef PubMed](#)
- [4] Mazier W, Saucisse N, Gatta-Cherifi B, Cota D. The endocannabinoid system: pivotal orchestrator of obesity and metabolic disease. *Trends Endocrinol Metab.* 2015; 26: 524-537. [CrossRef PubMed](#)
- [5] Maccarrone M, Bab I, Biró T, Cabral GA, Dey SK, Di Marzo V, Konje JC, Kunos G, Mechoulam R, Pacher P, Sharkey KA, Zimmer A. Endocannabinoid signaling at the periphery: 50 years after THC. *Trends Pharmacol Sci.* 2015; 36: 277-296. [CrossRef PubMed](#)
- [6] Rice AS. Should cannabinoids be used as analgesics for neuropathic pain? *Nat Clin Pract Neurol.* 2008; 4: 654-655. [CrossRef PubMed](#)
- [7] Martín-Sánchez E, Furukawa TA, Taylor J, Martin JL. Systematic review and meta-analysis of

- cannabis treatment for chronic pain. *Pain Med.* 2009; *10*: 1353-1368. [CrossRef PubMed](#)
- [8] Cravatt BF, Giang DK, Mayfield SP, Boger DL, Lerner RA, Gilula NB. Molecular characterization of an enzyme that degrades neuromodulatory fatty-acid amides. *Nature.* 1996; *384*: 83-87. [Cross-Ref PubMed](#)
- [9] Ahn K, McKinney MK, Cravatt BF. Enzymatic pathways that regulate endocannabinoid signaling in the nervous system. *Chem Rev.* 2008; *108*: 1687-1707. [CrossRef PubMed](#)
- [10] Cravatt BF, Demarest K, Patricelli MP, Bracey MH, Giang DK, Martin BR, Lichtman AH. Supersensitivity to anandamide and enhanced endogenous cannabinoid signaling in mice lacking fatty acid amide hydrolase. *Proc Natl Acad Sci USA.* 2001; *98*: 9371-9376. [CrossRef PubMed](#)
- [11] Lichtman AH, Shelton CC, Advani T, Cravatt BF. Mice lacking fatty acid amide hydrolase exhibit a cannabinoid receptor-mediated phenotypic hypoaesthesia. *Pain.* 2004; *109*: 319-327. [CrossRef PubMed](#)
- [12] Ahn K, Smith SE, Liimatta MB, Beidler D, Sadagopan N, Dudley DT, Young T, Wren P, Zhang Y, Swaney S, Van Becelaere K, Blankman JL, Nomura DK, Bhattachar SN, Stiff C, Nomanbhoy TK, Weerapana E, Johnson DS, Cravatt BF. Mechanistic and pharmacological characterization of PF-04457845: a highly potent and selective fatty acid amide hydrolase inhibitor that reduces inflammatory and noninflammatory pain. *J Pharmacol Exp Ther.* 2011; *338*: 114-124. [Cross-Ref PubMed](#)
- [13] Li GL, Winter H, Arends R, Jay GW, Le Y, Young T, Huggins JP. Assessment of the pharmacology and tolerability of PF-04457845, an irreversible inhibitor of fatty acid amide hydrolase-1, in healthy subjects. *Br J Clin Pharmacol.* 2012; *73*: 706-716. [CrossRef PubMed](#)
- [14] Huitron-Resendiz S, Sanchez-Alavez M, Wills DN, Cravatt BF, Henriksen SJ. Characterization of the sleep-wake patterns in mice lacking fatty acid amide hydrolase. *Sleep.* 2004; *27*: 857-865. [PubMed](#)
- [15] Lichtman AH, Leung D, Shelton CC, Saghatelian A, Hardouin C, Boger DL, Cravatt BF. Reversible inhibitors of fatty acid amide hydrolase that promote analgesia: evidence for an unprecedented combination of potency and selectivity. *J Pharmacol Exp Ther.* 2004; *311*: 441-448. [CrossRef PubMed](#)
- [16] Chang L, Luo L, Palmer JA, Sutton S, Wilson SJ, Barbier AJ, Breitenbucher JG, Chaplan SR, Webb M. Inhibition of fatty acid amide hydrolase produces analgesia by multiple mechanisms. *Br J Pharmacol.* 2006; *148*: 102-113. [CrossRef PubMed](#)
- [17] Huggins JP, Smart TS, Langman S, Taylor L, Young T. An efficient randomised, placebo-controlled clinical trial with the irreversible fatty acid amide hydrolase-1 inhibitor PF-04457845, which modulates endocannabinoids but fails to induce effective analgesia in patients with pain due to osteoarthritis of the knee. *Pain.* 2012; *153*: 1837-1846. [CrossRef PubMed](#)
- [18] Zannikos P, Wennerholm M, Van der Ark P, Vaccaro N, Schmidt M, Rassnick S, et al. The pharmacokinetics (PK), pharmacodynamics (PD), safety, and tolerability of JNJ-42165279, a potent and selective inhibitor of fatty acid amide hydrolase (FAAH) in healthy subjects. *Neuropsychopharmacology.* 2014; *39*: S43.
- [19] Sanofi. [http://en.sanofi.com/img/content/study/DFI10560\\_summary.pdf](http://en.sanofi.com/img/content/study/DFI10560_summary.pdf). 2016.