



HAL
open science

Identifying key predictors of deep brain stimulation outcomes – A feature importance analysis

Vittoria Bucciarelli, Dorian Vogel, Teresa Nordin, Marc Stawiski, Jerome Coste, Jean-Jacques Lemaire, Karin Wårdell, Raphaël Guzman, Simone Hemm

► To cite this version:

Vittoria Bucciarelli, Dorian Vogel, Teresa Nordin, Marc Stawiski, Jerome Coste, et al.. Identifying key predictors of deep brain stimulation outcomes – A feature importance analysis. BMT 2025 | Joint Conference of the Swiss (SSBE), Austrian (ÖGBMT) and German (VDE DGBMT) Societies for Biomedical Engineering, SSBE Swiss Society for Biomedical Engineering; ÖGBMT Austrian Society for Biomedical Engineering; VDE DGBMT German Society for Biomedical Engineering, Sep 2025, Bâle, Switzerland. pp.233, Id 065, <10.1515/bmt-2025-1001>. <hal-05141855>

HAL Id: hal-05141855

<https://uca.hal.science/hal-05141855v1>

Submitted on 3 Jul 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

Identifying Key Predictors of Deep Brain Stimulation Outcomes: a Feature Importance Analysis

1st author Vittoria Bucciarelli (vittoria.bucciarelli@fhnw.ch), **2nd author Dorian Vogel** (dorian.vogel@fhnw.ch), **4th author Marc Stawiski** (marc.jermann@fhnw.ch), **7th author Karin Wårdell** (karin.wardell@liu.se), **9th author Simone Hemm** (simone.hemm@fhnw.ch): Institute for Medical Engineering and Medical Informatics, School of Life Sciences, University of Applied Sciences and Arts Northwestern Switzerland, Muttenz, Switzerland

3rd author Teresa Nordin (teresa.nordin@liu.se), **7th author Karin Wårdell** (karin.wardell@liu.se), **9th author Simone Hemm** (simone.hemm@fhnw.ch): Department of Biomedical Engineering, Linköping University, Linköping, Sweden

5th author Jérôme Coste (jerome.coste@uca.fr), **6th author Jean-Jacques Lemaire** (jjlemaire@chu-clermontferrand.fr): Université Clermont Auvergne, Clermont Auvergne INP, CHU Clermont-Ferrand, CNRS, Institut Pascal, F-63000 Clermont-Ferrand, France

8th author Raphael Guzman (Raphael.Guzman@usb.ch): Department of Neurosurgery, University Hospital Basel, Basel, Switzerland

Introduction

Data-driven prediction of Deep Brain Stimulation (DBS) effects represents the foundation for algorithms automating DBS programming. Several models aiming at predicting symptom improvement in correspondence with specific stimulation parameters have been presented in literature. They exploit features ranging from patients' clinical characteristics, to measures extracted from fMRI scans or probabilistic mapping. However, the relationships between predictive features and clinical improvement are often underexplored. Therefore, this study aims to conduct a feature importance analysis to identify the most influential predictors of DBS outcomes.

Methods

2112 intra-operative stimulation tests of 65 patients were used to simulate the Volumes of Tissue Activated (VTAs) and to compute probabilistic volumes of therapeutic (Probabilistic Sweet Spots, PSS) and side effects (SE). Clinical, morphological, stimulation and probabilistic mapping-related features were calculated and DBS effects were categorized in 3 classes (side effects, low and high improvement). Feature importance was assessed with: univariate feature selection with ANOVA F-value, permutation importance and SHAP values.

Results

Despite being based on different principles, all three approaches consistently identified the same top three most influential predictors. These were: "VTA volume" ($F=269.5$), "PSS-VTA centroid distance" ($F=123.3$) and "VTA-SE volume overlap" ($F=115.9$). SHAP analysis revealed that high values of these features drive the prediction towards side effects class, whereas low values shift it towards high improvement class. Low improvement is associated with high VTA volumes and distances to PSS, coupled with low overlap with SE.

Conclusion

The findings highlight the predictive value of features derived from probabilistic mapping results in determining DBS outcomes. Clinical and anatomical features demonstrated limited predictive power. The high importance of the "VTA volume" may be attributed to the characteristics of the collected data: tests at the minimum amplitude yielding a therapeutic effect and maximum amplitude before onset of adverse effects.

This work was financed by the Swiss National Science Foundation (205320_207491).